

①

報告番号 甲 第 3130 号


学位論文

Zero-mode, winding number and quantization
of abelian sigma model in (1+1) dimensions

((1+1) 次元の可換シグマ模型の
トポロジーと量子化)

1994 年

谷村 省吾



Zero-mode, winding number and quantization of abelian sigma model in (1+1) dimensions

Shogo TANIMURA

Department of Physics, Nagoya University, Nagoya 464-01, Japan

Abstract

We consider the $U(1)$ sigma model in the two dimensional space-time $S^1 \times R$, which is a field-theoretical model possessing a non-trivial topology. It is pointed out that its topological structure is characterized by the zero-mode and the winding number. A new type of commutation relations is proposed to quantize the model respecting the topological nature. Hilbert spaces are constructed to be representation spaces of quantum operators. It is shown that there are an infinite number of inequivalent representations as a consequence of the nontrivial topology. The algebra generated by quantum operators is deformed by the central extension. When the central extension is introduced, it is shown that the zero-mode variables and the winding variables obey a new commutation relation, which we call twist relation. In addition, it is shown that the central extension makes momenta operators obey anomalous commutators. We demonstrate that topology enriches the structure of quantum field theories.

1 Introduction

All models in high energy physics proposed so far, for example, the standard model, grand unification models, supersymmetric models and technicolor models have large symmetries. We believe that such a large symmetry is spontaneously broken by some mechanism to a smaller symmetry, which is realized in the observed particle spectra. Symmetry and its breaking play an essential role in present understanding of high energy physics.

When a continuous symmetry is broken, degenerate vacua form a continuous manifold. The vacuum manifold possesses a nontrivial topology which is determined by the pattern of symmetry breaking. It is well known that massless excitations appear on the symmetry-breaking vacuum. These massless particles are called Nambu-Goldstone (NG) bosons. The number of species of NG bosons is equal to the dimension of the vacuum manifold. Threshold behavior of NG bosons is determined by the low energy theorems which follow as a consequence of symmetry.

To describe the dynamics of NG bosons in the language of field theory, a nonlinear sigma model was invented. In this model NG bosons are described by a field variable which is a mapping from the space-time to the vacuum manifold. This model is usually studied by the conventional canonical quantization and the perturbative method. However, in such a treatment, only a local property of vacuum is considered and no attention is paid to the global one. Namely, although the vacuum manifold has a nontrivial topology, the global nature cannot be seen by the perturbative approach. Then we expect that the nonperturbative behavior reflects the global nature of the vacuum. So we would like to have a new framework to study a quantum field theory with a nontrivial topology. If such a framework is successfully constructed, we expect to obtain a new insight into nonperturbative properties of the models which exhibit spontaneous symmetry breaking (SSB). These are the motivations for this investigation.

In this paper we would like to develop quantum theory in the direction to seek for another possibility which inheres in the theory itself. In the following we shall explain the well-known feature of quantum theory and next we shall show the direction to pursue.

To formulate a quantum theory we start from commutation relations among generators. Generators define an algebra, which is associative but not commutative.

When we have a classical theory, we ordinarily bring commutators among canonical variables by replacing the Poisson brackets. Then we construct a Hilbert space to be an irreducible representation space of the algebra. An element of the algebra is chosen and called a hamiltonian. Choice of a hamiltonian is restricted by physical requirements like self-adjointness, positivity and symmetries. The set formed by the algebra, the representation space and the hamiltonian is called a quantum theory. Then formulation itself is finished.

Next tasks are to solve it; we want to know eigenvalues of various observables, especially we are interested in spectrum of the hamiltonian, we want to know probability amplitude of transition between an initial state and a final one. For physically interesting theories, those are difficult tasks. Of course, we appreciate that they are worth hard work. However it is not the direction in which we would like to proceed in this paper. We are interested in a rather formal aspect.

A feature of quantum theory is that it requires a representation space; operators must be provided with operands. A commutator in quantum theory corresponds to a Poisson bracket in classical theory. But the Hilbert space in quantum theory has no correspondence in classical theory. State vectors, the superposition principle and amplitudes are characteristic concepts of quantum theory. We begin with an algebra of operators, then we construct a representation space. That is a unique procedure of quantum theory.

Here arise two questions; does a representation space exist? Is it unique? If there are inequivalent representations, calculation based on a different space gives a different answer for a physical quantity, for example, spectrum or amplitude.

When we consider a particle in a Euclidean space, we begin with the usual canonical commutation relations;

$$[\hat{x}_i, \hat{x}_j] = 0, \quad (i, j = 1, \dots, n) \quad (1.1)$$

$$[\hat{x}_i, \hat{p}_j] = i \delta_{ij}, \quad (1.2)$$

$$[\hat{p}_i, \hat{p}_j] = 0. \quad (1.3)$$

According to von Neumann's theorem the irreducible representation of the above algebra exists uniquely within a unitary equivalence class. Therefore there is no problem in choice of a Hilbert space. Although one may use the wave function representation and another may use the harmonic oscillator representation, both obtain a same result for calculation of a physical quantity.

Is there no need to worry about existence and uniqueness of a representation? Actually it is needed. We have encountered a situation in which the uniqueness is violated, when we consider a quantum field theory. In a quantum field theory we construct a representation space by defining a vacuum state and a Fock space. It was found that in several models there exist inequivalent vacuum states and they result in inequivalent Fock spaces. The different vacua are characterized by its transformation property under a certain symmetry operation. Such a situation is called spontaneous symmetry breaking (SSB). The discovery of SSB opened rich aspects of quantum field theories and led to deep understanding of the nature.

A field theory deals with a system which has infinite degrees of freedom, namely it is defined with a infinite number of generators which are called field variables. It is known that SSB is related to the infinity of degrees of freedom. On the other hand a particle has only finite degrees of freedom. Is there no occurrence of inequivalent representations in a quantum theory of a particle? (Usually a quantum theory with finite degrees of freedom is called a quantum mechanics.)

A strange result was found; when Ohnuki and Kitakado [1] investigated a quantum mechanics of a particle on a circle S^1 , they showed that there are a infinite number of inequivalent Hilbert spaces. Those spaces are parametrized by a continuous parameter α ranging from 0 to 1. What they have shown is that even a system with finite degrees of freedom can possess inequivalent representations when topology of the system is nontrivial. After that work, they studied a quantum mechanics on a sphere $S^n (n \geq 2)$ and showed existence of an infinite number of inequivalent Hilbert spaces specified by a discrete index.

Let us turn to field theories. The scalar field theory is a field-theoretical correspondence to the quantum mechanics of a particle in a Euclidean space. This field theory is quantized by requiring the canonical commutators and constructing the Fock space. There is also a correspondence in field theories to a quantum mechanics on a nontrivial manifold. It is a nonlinear sigma model because it has a manifold-valued field. For a review on nonlinear sigma models, see the reference [2].

Originally the nonlinear sigma model is designed to describe behavior of NG bosons at low energy scale. NG bosons are massless excitations associated with SSB. When a continuous symmetry specified by a group G is broken to a smaller symmetry specified by a subgroup H , vacua form a manifold which is called a homogeneous space G/H . In this model NG bosons are described by a field taking values in G/H .

It is already known [3] that even the quantum mechanics on G/H has inequivalent Hilbert spaces. Therefore it is naturally expected that a quantum field theory with a manifold-valued field may possess inequivalent Hilbert spaces. However in a usual approach, the nonlinear sigma model is quantized by the canonical quantization as the scalar field theory and is solved by the perturbative method. Thus the topological nature of the theory is missed and only a Fock space provides a representation.

If topological properties of field theories do not play an important role in physical application, we could neglect them. However we know several models which have nontrivial topology and in which topology plays an important role. For instance, the sine-Gordon model has topological kinks [4]; some gauge theories have topologically nontrivial vacua, so-called θ -vacua [5]; some nonlinear sigma models have the Wess-Zumino-Witten term, which reflects anomaly and topology of vacua [6]; the configuration space of nonabelian gauge fields modulo gauge transformations has an extremely complicated topology and causes the Gribov problem [7], and so on. Hence we would like to develop quantum field theories respecting topological nature.

One of the aims of this paper is to demonstrate that there are a lot of possibilities in constructing of quantum field theories, even if they are identical as classical theories. The second aim is to clarify the relation between topology and quantization. In the section 2 we will give a review of the quantum mechanics on S^1 . We will discuss physical implication of the existence of inequivalent representations. The section 3 is a main part of the present paper. There we consider a simple but nontrivial field-theoretical model, the abelian sigma model in (1+1) dimensions. We propose a definition of an algebra, which is quite different from the canonical one. Then we will construct Hilbert spaces and classify inequivalent ones. The section 4 is devoted to discussion on the results and directions for future development. This paper is a detailed and extended sequel to the previous work [8].

2 Quantum mechanics on S^1

Here we shall give a review of the quantum mechanics on S^1 [1]. Idea in it is useful for the next step in the field theory. More detailed consideration is found in the reference [9].

2.1 Motivation

Let us consider a particle moving on a circle S^1 . Its position is indicated by the angle coordinate θ . The coordinate $\theta + 2\pi$ indicates the same point as θ does. Hence θ is multivalued function on S^1 . Because of its nontrivial topology, a continuous and single-valued coordinate does not exist on S^1 .

Now let us quantize it. We should define an algebra. If we take the canonical approach, assume that $\hat{\theta}$ and \hat{P} are self-adjoint operators satisfying

$$[\hat{\theta}, \hat{P}] = i. \quad (2.1)$$

Next we should construct a representation space. If we define eigenstates of $\hat{\theta}$ by

$$\hat{\theta}|\theta\rangle = \theta|\theta\rangle, \quad (2.2)$$

we deduce that

$$\langle\theta|\hat{P}|\psi\rangle = -i\frac{\partial}{\partial\theta}\langle\theta|\psi\rangle \quad (2.3)$$

for an arbitrary state $|\psi\rangle$. Notice that the eigenvalue of $\hat{\theta}$ ranges over from $-\infty$ to $+\infty$; it is *not true* that $|\theta+2\pi\rangle = |\theta\rangle$. What we have constructed is just the quantum mechanics on \mathbf{R} , not S^1 .

What is wrong? It is wrong to use the multivalued coordinate $\hat{\theta}$ as a generator of the algebra. We must use a well-defined single-valued generator from the beginning.

As a substitute for θ , Ohnuki and Kitakado proposed to use $U = e^{i\hat{\theta}}$, which is complex-valued in the classical theory and a unitary operator in the quantum theory. From (2.1) we can deduce

$$[e^{i\hat{\theta}}, \hat{P}] = -e^{i\hat{\theta}}. \quad (2.4)$$

But they did not regard $\hat{\theta}$ as a generator. Instead they took existence of a unitary operator \hat{U} and a self-adjoint operator \hat{P} satisfying

$$[\hat{U}, \hat{P}] = -\hat{U} \quad (2.5)$$

as an assumption.

2.2 Representations

Now we see their construction of representations of the algebra defined by the above commutation relation (2.5). Since \hat{P} is a self-adjoint operator, it has an eigenvector with a real eigenvalue α ;

$$\hat{P}|\alpha\rangle = \alpha|\alpha\rangle, \quad \langle\alpha|\alpha\rangle = 1. \quad (2.6)$$

\hat{U} raises the eigenvalues of \hat{P} ;

$$\begin{aligned} \hat{P}\hat{U}|\alpha\rangle &= ([\hat{P}, \hat{U}] + \hat{U}\hat{P})|\alpha\rangle \\ &= (\hat{U} + \hat{U}\alpha)|\alpha\rangle \\ &= (1 + \alpha)\hat{U}|\alpha\rangle. \end{aligned} \quad (2.7)$$

Inversely, $\hat{U}^{-1} = \hat{U}^\dagger$ lowers them. Then we define

$$|n + \alpha\rangle = \hat{U}^n|\alpha\rangle, \quad (n = 0, \pm 1, \pm 2, \dots) \quad (2.8)$$

which have the following properties

$$\hat{P}|n + \alpha\rangle = (n + \alpha)|n + \alpha\rangle, \quad (2.9)$$

$$\langle m + \alpha | n + \alpha \rangle = \delta_{mn}. \quad (2.10)$$

The latter follows from self-adjointness of \hat{P} and unitarity of \hat{U} . With a fixed real number α , we define a Hilbert space H_α by completing the vector space of linear combinations of $|n + \alpha\rangle$ (n : integer). Equation (2.9) with

$$\hat{U}|n + \alpha\rangle = |n + 1 + \alpha\rangle \quad (2.11)$$

defines an irreducible representation of the algebra (2.5) on H_α .

H_α and H_β are unitary equivalent if and only if the difference $(\alpha - \beta)$ is an integer. Therefore the classification of irreducible representations of the algebra (2.5) has been completed; the whole of inequivalent irreducible representation spaces is $\{H_\alpha\}$ ($0 \leq \alpha < 1$). We call them Ohnuki-Kitakado representations.

2.3 Physical implications

Wave function

The physical meaning of the parameter α seems obscure. To clarify it they [1] studied eigenstates of the position operator \hat{U} . If we put

$$|\theta\rangle = \sum_{n=-\infty}^{\infty} e^{-in\theta} |n + \alpha\rangle, \quad (2.12)$$

it follows that

$$\hat{U}|\theta\rangle = e^{i\theta}|\theta\rangle, \quad (2.13)$$

$$|\theta + 2\pi\rangle = |\theta\rangle, \quad (2.14)$$

$$\langle\theta|\theta'\rangle = 2\pi\delta(\theta - \theta'). \quad (2.15)$$

In the last equation it is assumed that the δ -function is periodic with periodicity 2π . Eq. (2.14) is a desired property for the quantum mechanics on S^1 . It is natural to call \hat{U} a position operator due to this property. On the other hand, if we define $\hat{V}(\mu) = \exp(-i\mu\hat{P})$ for a real number μ , (2.5) implies that

$$\hat{V}^\dagger(\mu)\hat{U}\hat{V}(\mu) = e^{i\mu}\hat{U}. \quad (2.16)$$

Thus $\hat{U}\hat{V}(\mu)|\theta\rangle = e^{i(\theta+\mu)}\hat{V}(\mu)|\theta\rangle$ is an immediate consequence. A direct calculation shows that

$$\hat{V}(\mu)|\theta\rangle = e^{-i\alpha\mu}|\theta + \mu\rangle, \quad (2.17)$$

which says that \hat{P} is a generator of translation along S^1 . It should be noticed that an extra phase factor $e^{-i\alpha\mu}$ is multiplied. These states $|\theta\rangle$ ($0 \leq \theta < 2\pi$) define a wave function $\psi(\theta)$ for an arbitrary state $|\psi\rangle \in H_\alpha$ by $\psi(\theta) = \langle\theta|\psi\rangle$. This definition gives an isomorphism between H_α and $L^2(S^1)$ that is a space of square-integrable functions on S^1 . A bit calculation shows that the operators act on the wave function as

$$\hat{U}\psi(\theta) = \langle\theta|\hat{U}|\psi\rangle = e^{i\theta}\psi(\theta), \quad (2.18)$$

$$\hat{V}(\mu)\psi(\theta) = \langle\theta|\hat{V}(\mu)|\psi\rangle = e^{-i\alpha\mu}\psi(\theta - \mu), \quad (2.19)$$

$$\hat{P}\psi(\theta) = \langle\theta|\hat{P}|\psi\rangle = \left(-i\frac{\partial}{\partial\theta} + \alpha\right)\psi(\theta). \quad (2.20)$$

In the last expression the parameter α looks like the vector potential for magnetic flux $\Phi = 2\pi\alpha$ surrounded by S^1 . It should be noticed that α cannot be removed by gauge transformation $\psi(\theta) \rightarrow \psi'(\theta) = \kappa(\theta)\psi(\theta)$, where κ is a function from S^1 to $U(1)$. If we could chose $\kappa(\theta) = e^{i\alpha\theta}$,

$$\left(-i\frac{\partial}{\partial\theta} + \alpha\right)\psi(\theta) = e^{-i\alpha\theta}\left(-i\frac{\partial}{\partial\theta}\right)\psi'(\theta) \quad (2.21)$$

thus α disappeared. However, because $\psi'(2\pi) = e^{i\alpha 2\pi}\psi'(0)$, ψ' is not a periodic function. Hence ψ' does not remain in $L^2(S^1)$. In picture of wave functions, it is the boundary condition $\psi(2\pi) = \psi(0)$ what obstructs elimination of α and therefore

causes inequivalent representations. In the quantum mechanics on \mathbb{R} , there is no such boundary condition, thus such an extra term can be wiped away by gauge transformation.

Spectrum

To see a physical effect of the parameter α let us consider a free particle on S^1 . A free particle is defined by the hamiltonian

$$\hat{H} = \frac{1}{2}\hat{P}^2. \quad (2.22)$$

Its eigenvalue problem is trivially solved by

$$\hat{H}|n + \alpha\rangle = \frac{1}{2}(n + \alpha)^2|n + \alpha\rangle. \quad (2.23)$$

Apparently, the spectrum depends on the parameter α . For $\alpha = m$ (m : integer), all the eigenvalues but one of the ground state are doubly degenerate. While for $\alpha = m + \frac{1}{2}$, the all eigenvalues are doubly degenerate. For other values of α , there is no degeneracy. It is shown in the previous work [9] that these degeneracies reflect the parity symmetry. As $n + \alpha = (n - 1) + (\alpha + 1)$, the spectrum on the Hilbert space H_α is same as that on $H_{\alpha+1}$. Moreover, as $(n + \alpha)^2 = (-n - \alpha)^2$, the spectrum of on H_α is same to that on $H_{-\alpha}$, too. Therefore distinguishable values of α range over $0 \leq \alpha \leq \frac{1}{2}$.




Fig.1 Spectrum of a free particle on S^1

Path integral

As another example to show a physical effect of the parameter α , we shall see a path integral expression of the quantum mechanics on S^1 . We borrow a result from [10]. When the hamiltonian is of the form

$$\hat{H} = \frac{1}{2}\hat{P}^2 + V(\hat{U}, \hat{U}^\dagger), \quad (2.24)$$

they derived a path integral expression of transition amplitude as

$$\begin{aligned} K(\theta', \theta; t) &= \langle \theta' | \exp(-i\hat{H}t) | \theta \rangle \\ &= \sum_{n=-\infty}^{\infty} \int_{\text{winding } n \text{ times}} \mathcal{D}\theta \exp(iS_{\text{eff}}), \end{aligned} \quad (2.25)$$

where the effective action is defined as

$$S_{\text{eff}} = \int dt \left[\frac{1}{2} \left(\frac{d\theta}{dt} \right)^2 - V(\theta) - \alpha \frac{d\theta}{dt} \right]. \quad (2.26)$$

In (2.25), the integration is performed over paths winding n times around S^1 and the summation is performed with respect to the winding number. We would like to emphasize that the above path integral expression is derived from the operator formalism. It should be noticed that the global property—winding number—appears from the operator formalism alone.




Fig.2 Path integral on S^1

The last term in (2.26), $\int \alpha d\theta$ has no influence on the equation of motion but has an observable effect on the amplitude. To see the role of this term we rewrite (2.25) as

$$K(\theta', \theta; t) = e^{-i\alpha(\theta' - \theta)} \sum_{n=-\infty}^{\infty} e^{-i\alpha 2\pi n} \int_{\text{winding } n \text{ times}} \mathcal{D}\theta \exp(iS_0), \quad (2.27)$$

where $S_0 = \int dt [\frac{1}{2}\dot{\theta}^2 - V(\theta)]$. An amplitude for a path winding n times is weighted by the phase factor $\omega_n = \exp(-i\alpha 2\pi n)$. This phase factor causes observable interference effect; this phenomenon is analogous to the Aharonov-Bohm effect. Furthermore, ω_n 's obey composition rule; $\omega_m \omega_n = \omega_{m+n}$, which says that m -times winding followed by n -times one is equal to $(m+n)$ -times one. According to [11], ω_n can be interpreted as a unitary representation of the first homotopy group $\pi^1(S^1)$.

We conclude this section by repeating what has been shown. To formulate the quantum mechanics on S^1 we should choose suitable generators to define the algebra.

We recognize that θ is not suitable but U is suitable. The algebra is defined by the commutation relation (2.5). Representation spaces are constructed and inequivalent ones are parametrized by a continuous parameter α ($0 \leq \alpha < 1$). Inequivalent ones give different solutions to physical problems. The role of α resembles that of the vector potential. Topology of S^1 is an obstruction against elimination of such a vector potential. In path integral picture, α characterizes homotopy of a path of the particle on S^1 . Accordingly, existence of α reflects topology of S^1 .

3 Abelian sigma model

Here we shall consider the abelian sigma model as a generalization of the quantum mechanics on S^1 . The abelian sigma model has a field which takes values in S^1 . This model is designed to describe an NG boson associated with spontaneous breaking of $U(1)$ symmetry. We shall define two classes of algebras of the field theory; the first one is a natural generalization of the quantum mechanics on S^1 , which is called an algebra without central extension; the second one is its nontrivial generalization, which is called an algebra with central extension. For both classes we shall construct representation spaces combining the usual Fock representation with the Ohnuki-Kitakado representation. Topology of the model is carefully treated during the construction.

3.1 Algebra

Definition

To motivate definition of the algebra we will take three steps. First we start from the quantum mechanics on a Euclidean space \mathbf{R}^n . We shall give another expression to the canonical commutation relations (1.1)-(1.3). If we put $\hat{V}(a) = \exp(-i \sum_j a_j \hat{p}_j)$ for real numbers $a = (a_1, \dots, a_n)$, $\hat{V}(a)$ is a unitary operator and satisfies

$$\hat{x}_j \hat{x}_k = \hat{x}_k \hat{x}_j, \quad (3.1)$$

$$\hat{V}^\dagger(a) \hat{x}_j \hat{V}(a) = \hat{x}_j + a_j, \quad (3.2)$$

$$\hat{V}(a) \hat{V}(b) = \hat{V}(a+b). \quad (3.3)$$

Geometrical meaning of the above algebra is obvious. (3.1) says that coordinates \hat{x} 's of configuration are simultaneously measurable. (3.2) implies that configuration is movable by the displacement operator $\hat{V}(a)$. (3.3) says that displacement operators satisfy the composition law. An irreducible representation is uniquely given by $L^2(\mathbf{R}^n)$.

Second we turn to the scalar field theory in (1+1) dimensions. The usual canonical commutation relations are

$$[\hat{\phi}(\sigma), \hat{\phi}(\sigma')] = 0, \quad (-\infty < \sigma, \sigma' < +\infty) \quad (3.4)$$

$$[\hat{\phi}(\sigma), \hat{\pi}(\sigma')] = i\delta(\sigma - \sigma'), \quad (3.5)$$

$$[\hat{\pi}(\sigma), \hat{\pi}(\sigma')] = 0, \quad (3.6)$$

where σ is a coordinate of the space. $\hat{\phi}$ and $\hat{\pi}$ are distributions valued in hermite operators. We introduce a unitary operator

$$\hat{V}(f) = \exp \left[-i \int_{-\infty}^{\infty} f(\sigma) \hat{\pi}(\sigma) d\sigma \right] \quad (3.7)$$

for a real-valued test function f . Then they satisfy

$$\hat{\phi}(\sigma) \hat{\phi}(\sigma') = \hat{\phi}(\sigma') \hat{\phi}(\sigma), \quad (3.8)$$

$$\hat{V}^\dagger(f) \hat{\phi}(\sigma) \hat{V}(f) = \hat{\phi}(\sigma) + f(\sigma), \quad (3.9)$$

$$\hat{V}(f) \hat{V}(g) = \hat{V}(f+g), \quad (3.10)$$

where f and g are arbitrary real-valued test functions. The above algebra is interpreted as follows. (3.8) means that the field configurations at separated points are simultaneously measurable. (3.9) implies that a field configuration is movable arbitrarily by displacement operators. (3.10) is nothing but the composition property of displacements. A representation is usually given in terms of the Fock space.

As the third step we generalize the quantum mechanics on S^1 to the one on n -dimensional torus $T^n = (S^1)^n$. We introduce unitary operators \hat{U}_j and self-adjoint operators \hat{P}_j ($j = 1, \dots, n$). Put $\hat{V}(\mu) = \exp(-i \sum_j \mu_j \hat{P}_j)$ for $\mu = (\mu_1, \dots, \mu_n) \in \mathbf{R}^n$. Naive generalization of (2.5) or (2.16) leads the following relations

$$\hat{U}_j \hat{U}_k = \hat{U}_k \hat{U}_j, \quad (3.11)$$

$$\hat{V}^\dagger(\mu) \hat{U}_j \hat{V}(\mu) = e^{i\mu_j} \hat{U}_j, \quad (3.12)$$

$$\hat{V}(\mu) \hat{V}(\nu) = \hat{V}(\mu + \nu). \quad (3.13)$$

Geometrical meaning is so obvious that explanation is not repeated. We only point out that (3.12) expresses action of \mathbf{R}^n on T^n by displacement. Representations of this algebra are constructed by tensor products of Ohnuki-Kitakado representations $H_{\alpha_1} \otimes \dots \otimes H_{\alpha_n}$. Therefore irreducible representations are parametrized by n -tuple parameter $\alpha = (\alpha_1, \dots, \alpha_n)$.

Finally we turn to the abelian sigma model in (1+1) dimensions. The space-time is assumed to be $S^1 \times \mathbf{R}$. On an equal-time space-slice, the classical field variable is a map from S^1 to S^1 . So the configuration space of the model is $Q = \text{Map}(S^1; S^1)$. On the other hand $\Gamma = \text{Map}(S^1; U(1))$ becomes a group by pointwise multiplication. The group $U(1)$ acts on S^1 by displacement. Thus the group Γ acts on the configuration space Q by pointwise action, that is to say, for $\gamma \in \Gamma$ and $\phi \in Q$ let us define $\gamma \cdot \phi \in Q$

by

$$(\gamma \cdot \phi)(\sigma) = \gamma(\sigma) \cdot \phi(\sigma) \quad (\sigma \in S^1), \quad (3.14)$$

where σ denotes a point of the base space. In the right-hand side the multiplication indicates the action of $U(1)$ on S^1 .

To clarify geometry of the classical theory, we shall decompose the degrees of freedom of $\phi \in Q$ and $\gamma \in \Gamma$. In the classical sense we may rewrite $\phi : S^1 \rightarrow S^1 \cong U(1)$ by

$$\phi(\sigma) = U e^{i(N\sigma + \varphi(\sigma))}, \quad (3.15)$$

where $U \in U(1)$, $N \in \mathbb{Z}$. φ satisfies the no zero-mode condition;

$$\text{Map}_0(S^1; \mathbb{R}) = \{\varphi : S^1 \rightarrow \mathbb{R} \mid C^\infty, \int_0^{2\pi} \varphi(\sigma) d\sigma = 0\}. \quad (3.16)$$

The decomposition (3.15) says that $Q \cong S^1 \times \mathbb{Z} \times \text{Map}_0(S^1; \mathbb{R})$. Geometrical meaning of this decomposition is apparent; U describes the zero-mode or collective motion of the field ϕ ; N is nothing but the winding number; φ describes fluctuation or local degrees of freedom of ϕ . Topologically nontrivial parts are U and N .




Fig.3 Decomposition of degrees of freedom of $\text{Map}(S^1; S^1)$

Similarly $\gamma : S^1 \rightarrow U(1)$ is also rewritten as

$$\gamma(\sigma) = e^{i(\mu + m\sigma + f(\sigma))}, \quad (3.17)$$

where $\mu \in \mathbb{R}$, $m \in \mathbb{Z}$ and $f \in \text{Map}_0(S^1; \mathbb{R})$. The action (3.14) of γ on ϕ is decomposed into

$$U \rightarrow e^{i\mu} U, \quad (3.18)$$

$$N \rightarrow N + m, \quad (3.19)$$

$$\varphi(\sigma) \rightarrow \varphi(\sigma) + f(\sigma), \quad (3.20)$$

according to (3.15) and (3.17). Thus the first component of γ (3.17) translates the zero-mode; the second one changes the winding number; the third one gives a homotopic deformation.

To quantize this system let us assume that $\hat{\phi}(\sigma)$ is a unitary operator for each point $\sigma \in S^1$ and $\hat{V}(\gamma)$ is a unitary operator for each element $\gamma \in \Gamma$. Moreover we define an algebra generated by $\hat{\phi}(\sigma)$ and $\hat{V}(\gamma)$ with the following relations

$$\hat{\phi}(\sigma) \hat{\phi}(\sigma') = \hat{\phi}(\sigma') \hat{\phi}(\sigma), \quad (\sigma, \sigma' \in S^1) \quad (3.21)$$

$$\hat{V}^\dagger(\gamma) \hat{\phi}(\sigma) \hat{V}(\gamma) = \gamma(\sigma) \hat{\phi}(\sigma), \quad (3.22)$$

$$\hat{V}(\gamma_1) \hat{V}(\gamma_2) = e^{-ic(\gamma_1, \gamma_2)} \hat{V}(\gamma_1 \gamma_2) \quad (\gamma_1, \gamma_2 \in \Gamma). \quad (3.23)$$

At the last line a function $c : \Gamma \times \Gamma \rightarrow \mathbb{R}$ is called a central extension, which satisfies the cocycle condition

$$c(\gamma_1, \gamma_2) + c(\gamma_1 \gamma_2, \gamma_3) = c(\gamma_1, \gamma_2 \gamma_3) + c(\gamma_2, \gamma_3) \pmod{2\pi} \quad (3.24)$$

to ensure associativity $(\hat{V}(\gamma_1) \hat{V}(\gamma_2)) \hat{V}(\gamma_3) = \hat{V}(\gamma_1) (\hat{V}(\gamma_2) \hat{V}(\gamma_3))$. If $c \equiv 0$, the algebra (3.21)-(3.23) is a straightforward generalization of (3.11)-(3.13) to a system with infinite degrees of freedom. We call the algebra defined by (3.21)-(3.23) the fundamental algebra of the abelian sigma model.

We should explain why such an extra phase factor $e^{-ic(\gamma_1, \gamma_2)}$ is introduced. $\hat{V}(\gamma)$ acts on $\hat{\phi}(\sigma)$ by adjoint action as shown in (3.22). This action expresses the action of the group Γ on Q in terms of quantum operators. It satisfies the composition law

$$\hat{V}^\dagger(\gamma_2) \hat{V}^\dagger(\gamma_1) \hat{\phi}(\sigma) \hat{V}(\gamma_1) \hat{V}(\gamma_2) = \hat{V}^\dagger(\gamma_1 \gamma_2) \hat{\phi}(\sigma) \hat{V}(\gamma_1 \gamma_2), \quad (3.25)$$

which is demanded from geometrical viewpoint. However the above composition law does *not* imply that $\hat{V}(\gamma_1) \hat{V}(\gamma_2) = \hat{V}(\gamma_1 \gamma_2)$. In other words, \hat{V} is not necessarily a genuine unitary representation of the group Γ . We always have a possibility to insert a phase factor as done in (3.23). In other words, \hat{V} may be a projective unitary representation. If we can find a function $b : \Gamma \rightarrow \mathbb{R}$ such that

$$c(\gamma_1, \gamma_2) = b(\gamma_1) + b(\gamma_2) - b(\gamma_1 \gamma_2), \pmod{2\pi} \quad (3.26)$$

by defining $\tilde{V}(\gamma) = e^{ib(\gamma)} \hat{V}(\gamma)$, (3.23) results in a genuine unitary representation $\tilde{V}(\gamma_1) \tilde{V}(\gamma_2) = \tilde{V}(\gamma_1 \gamma_2)$. In that case c is called a coboundary of b and denoted by $c = \delta b$. A coboundary $c = \delta b$ identically satisfies the condition (3.24), namely a coboundary is always a cocycle. We usually demand both b and c to be continuous

functions from Γ to $\mathbf{R}/2\pi\mathbf{Z}$, since Γ is continuous. A class of cocycles modulo coboundaries is called a cohomology. Existence of a nontrivial cohomology depends on topology of the group Γ . The group considered now is a so-called loop group $\Gamma = \text{Map}(S^1; G)$ with $G = U(1)$. Its nontrivial topology allows for existence of a nontrivial cohomology. On the other hand, when $\Gamma = \mathbf{R}^n$, all cocycles are coboundaries. Thus there is no need to insert such a central extension into (3.3) when we considered the quantum mechanics on \mathbf{R}^n .

We add a comment. In quantum field theories, we have met such extensions of algebras when we study anomalous gauge theories and conformal field theories. In a gauge theory in (3+1) dimensions with a gauge group G , gauge transformations form a group $\Gamma = \text{Map}(S^3, G)$. Anomaly is deeply related to topology of Γ . When a nontrivial cohomology exists, extra terms are added to the commutators of the Gauss law constraints [12]. Similarly in two dimensional conformal field theories [13], the algebra of energy-momentum tensor is deformed by a central extension due to the conformal anomaly and becomes the Virasoro algebra. In the two dimensional Wess-Zumino-Witten (WZW) model [6], the current algebra is also deformed by a central extension and becomes the Kac-Moody algebra. Its deformation is caused by the WZW term brought into the lagrangian of a nonlinear sigma model. A necessary condition for existence of the WZW term is that G has nontrivial cohomology $H^3(G)$. But $U(1)$ does not satisfy it. We do not yet know a physical reason why we must introduce a central extension into the abelian sigma model. Even a reason to chose a specific central extension is not clear. In the case of the WZW model, QCD as an underlying theory tells the reason to bring the anomaly. But the physical meaning of the central extension in our model still remains obscure.

Now let us return to the fundamental algebra (3.21)-(3.23). As a nontrivial central extension for γ_1 given by (3.17) and

$$\gamma_2(\sigma) = e^{i(\nu + n\sigma + g(\sigma))}, \quad (3.27)$$

we define

$$c(\gamma_1, \gamma_2) = k \left\{ m\nu - n\mu + \frac{1}{4\pi} \int_0^{2\pi} \left(\frac{df}{d\sigma} g(\sigma) - f(\sigma) \frac{dg}{d\sigma} \right) d\sigma \right\}, \quad (3.28)$$

where k is an integer. This central extension is the simplest but nontrivial one which is invariant under the action of the group of orientation-preserving diffeomorphisms $\text{Diff}^+(S^1)$; $c(\gamma_1 \circ \omega, \gamma_2 \circ \omega) = c(\gamma_1, \gamma_2)$ for any $\omega \in \text{Diff}^+(S^1)$. The group Γ associated

with such an invariant central extension is called a Kac-Moody group of rank k . The relation (3.23) says that \hat{V} is a unitary representation of the Kac-Moody group. For classification of central extensions see the literature [14].

Algebra without central extension

According to decomposition of classical variables (3.15) and (3.17), quantum operators are also to be decomposed. For simplicity we consider the fundamental algebra (3.21)-(3.23) without the central extension, that is, here we restrict $c \equiv 0$.

Corresponding to (3.15), we introduce a unitary operator \hat{U} , a self-adjoint operator \hat{N} satisfying

$$\exp(2\pi i \hat{N}) = 1, \quad (3.29)$$

which is called the integer condition for \hat{N} , and a distribution $\hat{\phi}(\sigma)$ valued in hermite operators and constrained by

$$\int_0^{2\pi} \hat{\phi}(\sigma) d\sigma = 0. \quad (3.30)$$

We demand that the quantum field $\hat{\phi}(\sigma)$ is expressed in terms of these as

$$\hat{\phi}(\sigma) = \hat{U} e^{i(\hat{N}\sigma + \hat{\phi}(\sigma))}. \quad (3.31)$$

Actually this equation is too naive. Because $\hat{\phi}(\sigma)$ is an operator-valued distribution, its exponentiation $\exp(i\hat{\phi}(\sigma))$ is ill-defined. To make it well-defined we should regularize its divergence. This problem is postponed until discussion on the normal ordering.

Next, corresponding to (3.17) we introduce a self-adjoint operator \hat{P} , a unitary operator \hat{W} , and a distribution $\hat{\pi}(\sigma)$ valued in hermite operators and constrained by

$$\int_0^{2\pi} \hat{\pi}(\sigma) d\sigma = 0, \quad (3.32)$$

When γ is given by (3.17), the operator $\hat{V}(\gamma)$ is defined by

$$\hat{V}(\gamma) = e^{-i\mu\hat{P}} \hat{W}^m \exp \left[-i \int_0^{2\pi} f(\sigma) \hat{\pi}(\sigma) d\sigma \right]. \quad (3.33)$$

Using these operators the relation (3.22) is now rewritten as

$$e^{i\mu\hat{P}} \hat{U} e^{-i\mu\hat{P}} = e^{i\mu} \hat{U}, \quad (3.34)$$

$$\hat{W}^\dagger \hat{N} \hat{W} = \hat{N} + 1, \quad (3.35)$$

$$\exp \left[i \int_0^{2\pi} f(\sigma) \hat{\pi}(\sigma) d\sigma \right] \hat{\phi}(\sigma) \exp \left[-i \int_0^{2\pi} f(\sigma) \hat{\pi}(\sigma) d\sigma \right] = \hat{\phi}(\sigma) + f(\sigma) \quad (3.36)$$

These represent the action of Γ on Q by (3.18)-(3.20). Observing the relation (3.35), we call \hat{N} and \hat{W} the winding number and the winding operator, respectively. They are also rewritten in terms of commutation relations

$$[\hat{P}, \hat{U}] = \hat{U}, \quad (3.37)$$

$$[\hat{N}, \hat{W}] = \hat{W}, \quad (3.38)$$

$$[\hat{\phi}(\sigma), \hat{\pi}(\sigma')] = i\left(\delta(\sigma - \sigma') - \frac{1}{2\pi}\right), \quad (3.39)$$

with all other vanishing commutators. In (3.39) it is understood that the δ -function is defined on S^1 .

Algebra with central extension

Before constructing representations, we reexpress the fundamental algebra with the central extension (3.28) respecting the decomposition (3.15) and (3.17). The decomposition (3.31) of $\hat{\phi}$ does not need to be changed. On the other hand the decomposition (3.33) of \hat{V} should be modified a little. We formally introduce an operator $\hat{\Omega}$ by

$$\hat{W} = e^{-i\hat{\Omega}}. \quad (3.40)$$

Although \hat{W} itself is well-defined, $\hat{\Omega}$ is ill-defined. If $\hat{\Omega}$ exists, (3.38) would imply $[\hat{N}, \hat{\Omega}] = i$, which is nothing but the canonical commutation relation. Therefore \hat{N} should have a continuous spectrum, that contradicts the integer condition (3.29). Consequently $\hat{\Omega}$ must be eliminated after calculation. Bearing the above remark in mind, we replace (3.33) by

$$\hat{V}(\gamma) = \exp\left[-i\left(\mu\hat{P} + m\hat{\Omega} + \int_0^{2\pi} f(\sigma)\hat{\pi}(\sigma)d\sigma\right)\right]. \quad (3.41)$$

For the central extension (3.28) of rank k , addition of the following commutation relations to (3.37)-(3.39) is enough to satisfy the fundamental algebra;

$$[\hat{P}, \hat{\Omega}] = -2ik, \quad (3.42)$$

$$[\hat{\pi}(\sigma), \hat{\pi}(\sigma')] = -\frac{ik}{\pi}\delta'(\sigma - \sigma'). \quad (3.43)$$

We assume that all other commutators vanish.

Below we verify that they are sufficient for the fundamental algebra. A useful formula is

$$e^{\hat{X}}e^{\hat{Y}} = e^{\frac{1}{2}[\hat{X}, \hat{Y}]}e^{\hat{X}+\hat{Y}}, \quad (3.44)$$

which is valid when $[\hat{X}, [\hat{X}, \hat{Y}]] = [\hat{Y}, [\hat{X}, \hat{Y}]] = 0$. This formula yields

$$e^{-i(\mu\hat{P}+m\hat{\Omega})}e^{-i(\nu\hat{P}+n\hat{\Omega})} = e^{-\frac{1}{2}(m\nu-n\mu)[\hat{\Omega}, \hat{P}]}e^{-i((\mu+\nu)\hat{P}+(m+n)\hat{\Omega})}. \quad (3.45)$$

If we substitute (3.42), we obtain

$$e^{-i(\mu\hat{P}+m\hat{\Omega})}e^{-i(\nu\hat{P}+n\hat{\Omega})} = e^{-ik(m\nu-n\mu)}e^{-i((\mu+\nu)\hat{P}+(m+n)\hat{\Omega})}, \quad (3.46)$$

which is coincident with (3.23) for the central extension (3.28). Similarly the formula (3.44) implies

$$\begin{aligned} & e^{-i\int_0^{2\pi} f(\sigma)\hat{\pi}(\sigma)d\sigma}e^{-i\int_0^{2\pi} g(\sigma')\hat{\pi}(\sigma')d\sigma'} \\ &= e^{-\frac{1}{2}\int_0^{2\pi}\int_0^{2\pi} f(\sigma)g(\sigma')[\hat{\pi}(\sigma), \hat{\pi}(\sigma')]d\sigma d\sigma'}e^{-i\int_0^{2\pi} (f(\sigma)+g(\sigma))\hat{\pi}(\sigma)d\sigma}. \end{aligned} \quad (3.47)$$

Substitution of (3.43) yields

$$\begin{aligned} & -\frac{1}{2}\int_0^{2\pi}\int_0^{2\pi} f(\sigma)g(\sigma')[\hat{\pi}(\sigma), \hat{\pi}(\sigma')]d\sigma d\sigma' \\ &= \frac{ik}{2\pi}\int_0^{2\pi}\int_0^{2\pi} f(\sigma)g(\sigma')\delta'(\sigma - \sigma')d\sigma d\sigma' \\ &= \frac{ik}{4\pi}\int_0^{2\pi} (-f'(\sigma)g(\sigma) + f(\sigma)g'(\sigma))d\sigma. \end{aligned} \quad (3.48)$$

Hence (3.47) coincides with (3.23) for the central extension (3.28).

Using (3.40) Eq. (3.42) implies

$$[\hat{P}, \hat{W}] = -2k\hat{W}, \quad (3.49)$$

which says that the zero-mode momentum \hat{P} is decreased by $2k$ units when the winding number \hat{N} is increased by one unit under the operation of \hat{W} . This is an inevitable consequence of the central extension. We call this phenomenon "twist". Using (3.42) and (3.44), the decomposition (3.41) results in

$$\hat{V}(\gamma) = e^{-ikm\mu} \exp\left[-i\left(\mu\hat{P} + \int_0^{2\pi} f(\sigma)\hat{\pi}(\sigma)d\sigma\right)\right]\hat{W}^m. \quad (3.50)$$

Here we summarize a temporal result. Generators of the fundamental algebra are decomposed as (3.31) and (3.50) considering topological nature of the model. They are constrained by the no-zero-mode condition (3.30), (3.32) and the integer condition (3.29). The commutation relations are also decomposed into (3.37), (3.38), (3.39), (3.43) and (3.49). Noticeable effects of the central extension are the anomalous commutator (3.43) and the twist relation (3.49). These features also affect representation of the algebra as seen in the following sections.

3.2 Representations

Without the central extension

Now we proceed to construct representations of the algebra defined by (3.37)-(3.39) and other vanishing commutators with the constraints (3.29), (3.30) and (3.32).

Remember that \hat{P} and \hat{N} are self-adjoint and that \hat{U} and \hat{W} are unitary. Both of the relations (3.37) and (3.38) are isomorphic to (2.5). Hence the Ohnuki-Kitakado representations provide representations for them, too. \hat{P} and \hat{U} act on the Hilbert space H_α via (2.9) and (2.11). \hat{N} and \hat{W} act on another Hilbert space H_β via

$$\hat{N}|n+\beta\rangle = (n+\beta)|n+\beta\rangle, \quad (3.51)$$

$$\hat{W}|n+\beta\rangle = |n+1+\beta\rangle. \quad (3.52)$$

The value of α is arbitrary. However β is restricted to be an integer if we impose the condition (3.29).

For $\hat{\varphi}$ and $\hat{\pi}$ the Fock representation works. We define operators \hat{a}_n and \hat{a}_n^\dagger by

$$\hat{\varphi}(\sigma) = \frac{1}{2\pi} \sum_{n \neq 0} \sqrt{\frac{\pi}{|n|}} (\hat{a}_n e^{in\sigma} + \hat{a}_n^\dagger e^{-in\sigma}), \quad (3.53)$$

$$\hat{\pi}(\sigma) = \frac{i}{2\pi} \sum_{n \neq 0} \sqrt{\pi|n|} (-\hat{a}_n e^{in\sigma} + \hat{a}_n^\dagger e^{-in\sigma}). \quad (3.54)$$

In the Fourier series the zero-mode $n=0$ is excluded because of the constraints (3.30) and (3.32). It is easily verified that the commutator (3.39) is equivalent to

$$[\hat{a}_m, \hat{a}_n^\dagger] = \delta_{m,n} \quad (m, n = \pm 1, \pm 2, \dots) \quad (3.55)$$

with the other vanishing commutators. Hence the ordinary Fock space F gives a representation of \hat{a}^\dagger 's and \hat{a} 's, which are called creation operators and annihilation operators, respectively.

Consequently the tensor product space $H_\alpha \otimes H_0 \otimes F$ gives an irreducible representation of the fundamental algebra without the central extension. The inequivalent ones are parametrized by α ($0 \leq \alpha < 1$).

A remark is in order here; the coefficients in front of \hat{a} 's in (3.53) and (3.54) are chosen to diagonalize the hamiltonian of free field

$$\begin{aligned} \hat{H} &= \frac{1}{2} \int_0^{2\pi} \left[\left(\frac{1}{2\pi} \hat{P} + \hat{\pi}(\sigma) \right)^2 + \left(\partial \hat{\varphi}(\sigma) + \hat{N} \right)^2 \right] d\sigma \\ &= \frac{1}{2} \left(\frac{1}{2\pi} \hat{P}^2 + 2\pi \hat{N}^2 \right) + \sum_{n \neq 0} |n| \left(\hat{a}_n^\dagger \hat{a}_n + \frac{1}{2} \right). \end{aligned} \quad (3.56)$$

This hamiltonian corresponds to the lagrangian density

$$\mathcal{L} = \frac{1}{2} \partial_\mu \phi^\dagger \partial^\mu \phi, \quad (3.57)$$

where ϕ is given by (3.15). This lagrangian describes a massless boson. Although Coleman's theorem [15] forbids existence of massless bosons in (1+1) dimensions due to infrared catastrophe, our model is still permitted. In our model the base space S^1 is compact, hence there is no infrared divergence. Interacting field theory will be briefly discussed later.

Another remark is added. The Kamefuchi-O'Raifeartaigh-Salam theorem [16] states that the S -matrix in quantum field theories remains unchanged under any point transformation of field variables. Their theorem is proved within the framework of the conventional canonical formalism. Thus this theorem is not applicable to our model. If we take a real scalar field φ such that $\phi(\sigma) = e^{i\varphi(\sigma)}$, the lagrangian (3.57) becomes a genuine free scalar theory,

$$\mathcal{L} = \frac{1}{2} \partial_\mu \varphi \partial^\mu \varphi. \quad (3.58)$$

In this case the Fock space is a unique representation, thus there is no room for such an undetermined parameter α . They considered only field theories with a trivial topology, then they derived the equivalence theorem for S -matrix. On the other hand, we considered a field theory with a nontrivial topology, then we reached existence of inequivalent representations. However the S -matrix of our model is not yet calculated, hence it is left undetermined whether S -matrix does depend on the parameter α or not.

With the central extension

Next we shall construct representations of the algebra defined by (3.37), (3.38), (3.39), (3.43), (3.49) and other vanishing commutators with the constraints (3.29), (3.30) and (3.32). The way of construction is a bit modified from the previous one.

Taking account of the twist relation (3.49), the representation of \hat{P} , \hat{U} , \hat{N} and \hat{W} are given by

$$\hat{P}|p+\alpha; n\rangle = (p+\alpha)|p+\alpha; n\rangle, \quad (3.59)$$

$$\hat{U}|p+\alpha; n\rangle = |p+1+\alpha; n\rangle, \quad (3.60)$$

$$\hat{N} |p + \alpha; n\rangle = n |p + \alpha; n\rangle, \quad (3.61)$$

$$\hat{W} |p + \alpha; n\rangle = |p - 2k + \alpha; n + 1\rangle. \quad (3.62)$$

The inner product is defined by

$$\langle p + \alpha; m | q + \alpha; n \rangle = \delta_{pq} \delta_{mn} \quad (p, q, m, n \in \mathbb{Z}). \quad (3.63)$$

The Hilbert space formed by completing the space of linear combinations of $|p + \alpha; n\rangle$ is denoted by T_α . (T indicates "twist".)

Let us turn to $\hat{\varphi}$ and $\hat{\pi}$. Considering the anomalous commutator (3.43), after a tedious calculation we obtain a Fourier expansion

$$\hat{\varphi}(\sigma) = \sum_{n \neq 0} \frac{1}{\sqrt{2|kn|}} (\hat{a}_n e^{in\sigma} + \hat{a}_n^\dagger e^{-in\sigma}), \quad (3.64)$$

$$\hat{\pi}(\sigma) = \frac{i}{2\pi} \left\{ \begin{array}{l} \sum_{n=1}^{\infty} \quad (k > 0) \\ \sum_{n=-1}^{-\infty} \quad (k < 0) \end{array} \right\} \sqrt{2|kn|} (-\hat{a}_n e^{in\sigma} + \hat{a}_n^\dagger e^{-in\sigma}), \quad (3.65)$$

where \hat{a} 's and \hat{a}^\dagger 's obey the same commutation relations (3.55). The derivation of the above expansion is shown in the appendix. It should be noticed that only positive n 's appear in the expansion of $\hat{\pi}$ when $k > 0$, while only negative n 's appear when $k < 0$. However both positive and negative n 's appear in $\hat{\varphi}$. Physical implication of lack of half of modes in $\hat{\pi}(\sigma)$ is still unclear but will be discussed later.

The algebra defined by (3.55) is also represented by the Fock space F . Hence the tensor product space $T_\alpha \otimes F$ gives an irreducible representation of the fundamental algebra with the central extension for each value of α ($0 \leq \alpha < 1$).

3.3 Normal ordering

Although most of our main subjects are finished, a subtle problem is still left. At (3.31) exponential of the local operator $\hat{\varphi}(\sigma)$ is introduced. As mentioned there, it contains divergence thus it is ill-defined. Now we shall consider this problem closely.

Without the central extension

$e^{i\hat{\varphi}(\sigma)}$ should be expressed in terms of creation and annihilation operators to act on the Fock space. First we study the case without central extension. If we define

$$\hat{\chi}(\sigma) = \frac{1}{2\pi} \sum_{n \neq 0} \sqrt{\frac{\pi}{|n|}} \hat{a}_n e^{in\sigma}, \quad (3.66)$$

(3.53) is decomposed into $\hat{\varphi}(\sigma) = \hat{\chi}(\sigma) + \hat{\chi}^\dagger(\sigma)$. A bit calculation shows that

$$\begin{aligned} [\hat{\chi}(\sigma), \hat{\chi}^\dagger(\sigma')] &= \frac{1}{4\pi} \left\{ \sum_{n=1}^{\infty} \frac{1}{n} e^{in(\sigma-\sigma')} + \sum_{n=1}^{\infty} \frac{1}{n} e^{-in(\sigma-\sigma')} \right\} \\ &= \frac{1}{4\pi} \{ \log(1 - e^{i(\sigma-\sigma')}) + \log(1 - e^{-i(\sigma-\sigma')}) \}, \end{aligned} \quad (3.67)$$

which diverges when σ' closes to σ ,

$$\lim_{\sigma' \rightarrow \sigma} [\hat{\chi}(\sigma), \hat{\chi}^\dagger(\sigma')] = -\infty. \quad (3.68)$$

Thus application of the formula (3.44) tells that

$$e^{i\hat{\varphi}(\sigma)} = e^{i\hat{\chi}(\sigma) + i\hat{\chi}^\dagger(\sigma)} = e^{-\frac{1}{2}[\hat{\chi}(\sigma), \hat{\chi}^\dagger(\sigma)]} e^{i\hat{\chi}^\dagger(\sigma)} e^{i\hat{\chi}(\sigma)} \quad (3.69)$$

diverges. To eliminate the divergence we introduce the normal ordering procedure, which is a rule to rearrange creation operators to the left and annihilation operators to the right for each term. This procedure is denoted by sandwiching by double colons, for example

$$: e^{i\hat{\varphi}(\sigma)} : = e^{i\hat{\chi}^\dagger(\sigma)} e^{i\hat{\chi}(\sigma)}. \quad (3.70)$$

As (3.44) implies that $e^{\hat{X}} e^{\hat{Y}} = e^{[\hat{X}, \hat{Y}]} e^{\hat{Y}} e^{\hat{X}}$, (3.67) gives

$$\begin{aligned} &: e^{i\hat{\varphi}(\sigma)} :: e^{i\hat{\varphi}(\sigma')} : \\ &= e^{i\hat{\chi}^\dagger(\sigma)} e^{i\hat{\chi}(\sigma)} e^{i\hat{\chi}^\dagger(\sigma')} e^{i\hat{\chi}(\sigma')} \\ &= e^{-[\hat{\chi}(\sigma), \hat{\chi}^\dagger(\sigma')]} e^{i\hat{\chi}^\dagger(\sigma)} e^{i\hat{\chi}^\dagger(\sigma')} e^{i\hat{\chi}(\sigma)} e^{i\hat{\chi}(\sigma')} \\ &= \exp \left[-\frac{1}{4\pi} \log(1 - e^{i(\sigma-\sigma')})(1 - e^{-i(\sigma-\sigma')}) \right] : e^{i\hat{\varphi}(\sigma)} e^{i\hat{\varphi}(\sigma')} : \end{aligned} \quad (3.71)$$

which is well-defined except for $\sigma = \sigma'$ and invariant under permutation of σ with σ' . Therefore we conclude that

$$: e^{i\hat{\varphi}(\sigma)} :: e^{i\hat{\varphi}(\sigma')} : = : e^{i\hat{\varphi}(\sigma')} :: e^{i\hat{\varphi}(\sigma)} :, \quad (3.72)$$

then (3.21) is satisfied.

The other relation (3.22) also must be satisfied after the normal ordering procedure. Let us check it. We substitute (3.54) and

$$f(\sigma) = \sum_{n \neq 0} f_n e^{in\sigma} \quad (f_{-n} = f_n^*) \quad (3.73)$$

into (3.33) to obtain

$$\hat{V}(f) = \exp \left[-i \int_0^{2\pi} f(\sigma) \hat{\pi}(\sigma) d\sigma \right] = \exp \left[\sum_{n \neq 0} \sqrt{\pi|n|} (-f_{-n} \hat{a}_n + f_n \hat{a}_n^\dagger) \right]. \quad (3.74)$$

Then it is easily seen that

$$\hat{V}^\dagger(f) \hat{a}_n \hat{V}(f) = \hat{a}_n + \sqrt{\pi|n|} f_n, \quad (3.75)$$

$$\hat{V}^\dagger(f) \hat{a}_n^\dagger \hat{V}(f) = \hat{a}_n^\dagger + \sqrt{\pi|n|} f_{-n}. \quad (3.76)$$

The definitions (3.66) and (3.73) with the above equations yield

$$\begin{aligned} \hat{V}^\dagger(f) \hat{\chi}(\sigma) \hat{V}(f) &= \frac{1}{2\pi} \sum_{n \neq 0} \sqrt{\frac{\pi}{|n|}} (\hat{a}_n + \sqrt{\pi|n|} f_n) e^{in\sigma} \\ &= \hat{\chi}(\sigma) + \frac{1}{2} f(\sigma), \end{aligned} \quad (3.77)$$

$$\begin{aligned} \hat{V}^\dagger(f) \hat{\chi}^\dagger(\sigma) \hat{V}(f) &= \frac{1}{2\pi} \sum_{n \neq 0} \sqrt{\frac{\pi}{|n|}} (\hat{a}_n^\dagger + \sqrt{\pi|n|} f_{-n}) e^{-in\sigma} \\ &= \hat{\chi}^\dagger(\sigma) + \frac{1}{2} f(\sigma), \end{aligned} \quad (3.78)$$

therefore

$$\begin{aligned} \hat{V}^\dagger(f) : e^{i\hat{\phi}(\sigma)} : \hat{V}(f) &= \hat{V}^\dagger(f) e^{i\hat{\chi}^\dagger(\sigma)} e^{i\hat{\chi}(\sigma)} \hat{V}(f) \\ &= e^{if(\sigma)} e^{i\hat{\chi}^\dagger(\sigma)} e^{i\hat{\chi}(\sigma)} \\ &= e^{if(\sigma)} : e^{i\hat{\phi}(\sigma)} :. \end{aligned} \quad (3.79)$$

This result coincides with (3.22). Consequently we have checked that the fundamental relations (3.21) and (3.22) are preserved by the normal ordering procedure. Although $\hat{\phi}(\sigma)$ is claimed to be a unitary operator above (3.21), $\hat{\phi}(\sigma) = : e^{i\hat{\phi}(\sigma)} :$ is not unitary. If it were unitary, $\hat{\phi}(\sigma) \hat{\phi}^\dagger(\sigma)$ must be equal to identity. But actually

$$\begin{aligned} \lim_{\sigma' \rightarrow \sigma} \hat{\phi}(\sigma) \hat{\phi}^\dagger(\sigma') &= \lim_{\sigma' \rightarrow \sigma} e^{i\hat{\chi}^\dagger(\sigma)} e^{i\hat{\chi}(\sigma)} e^{-i\hat{\chi}^\dagger(\sigma')} e^{-i\hat{\chi}(\sigma')} \\ &= \lim_{\sigma' \rightarrow \sigma} e^{[\hat{\chi}(\sigma), \hat{\chi}^\dagger(\sigma')]} e^{i\hat{\chi}^\dagger(\sigma)} e^{-i\hat{\chi}^\dagger(\sigma')} e^{i\hat{\chi}(\sigma)} e^{-i\hat{\chi}(\sigma')} \\ &= \lim_{\sigma' \rightarrow \sigma} e^{[\hat{\chi}(\sigma), \hat{\chi}^\dagger(\sigma)]} \hat{1} \\ &= 0 \end{aligned} \quad (3.80)$$

due to (3.68), hence $\hat{\phi}(\sigma) = : e^{i\hat{\phi}(\sigma)} :$ is not unitary.

With the central extension

Similarly we can verify the case of $k \neq 0$. The way of verification is almost identical to the previous one but a bit changed. We define

$$\hat{\chi}(\sigma) = \sum_{n \neq 0} \frac{1}{\sqrt{2|kn|}} \hat{a}_n e^{in\sigma}, \quad (3.81)$$

to decompose (3.64) into $\hat{\phi}(\sigma) = \hat{\chi}(\sigma) + \hat{\chi}^\dagger(\sigma)$. A bit calculation yields

$$\begin{aligned} [\hat{\chi}(\sigma), \hat{\chi}^\dagger(\sigma')] &= \frac{1}{2|k|} \left\{ \sum_{n=1}^{\infty} \frac{1}{n} e^{in(\sigma-\sigma')} + \sum_{n=1}^{\infty} \frac{1}{n} e^{-in(\sigma-\sigma')} \right\} \\ &= \frac{1}{2|k|} \{ \log(1 - e^{i(\sigma-\sigma')}) + \log(1 - e^{-i(\sigma-\sigma')}) \}, \end{aligned} \quad (3.82)$$

which is also divergent when $\sigma = \sigma'$. To eliminate the divergence we use the normal ordering procedure again;

$$: e^{i\hat{\phi}(\sigma)} : = e^{i\hat{\chi}^\dagger(\sigma)} e^{i\hat{\chi}(\sigma)}. \quad (3.83)$$

A calculation similar to (3.71) yields

$$: e^{i\hat{\phi}(\sigma)} :: e^{i\hat{\phi}(\sigma')} : = \exp \left[-\frac{1}{2|k|} \log(1 - e^{i(\sigma-\sigma')})(1 - e^{-i(\sigma-\sigma')}) \right] : e^{i\hat{\phi}(\sigma)} e^{i\hat{\phi}(\sigma')} :. \quad (3.84)$$

Thus the commutativity (3.21) is ensured again.

Verification of the other relation (3.22) is a bit complicated. We define

$$\begin{aligned} f^{(+)}(\sigma) &= \sum_{n=1}^{\infty} f_n e^{in\sigma}, \\ f^{(-)}(\sigma) &= \sum_{n=-1}^{-\infty} f_n e^{in\sigma}, \\ f(\sigma) &= f^{(+)}(\sigma) + f^{(-)}(\sigma). \quad (f_{-n} = f_n^*) \end{aligned} \quad (3.85)$$

Substitution of (3.65) into (3.50) gives

$$\begin{aligned} \hat{V}(f) &= \exp \left[-i \int_0^{2\pi} f(\sigma) \hat{\pi}(\sigma) d\sigma \right] \\ &= \exp \left[\left\{ \sum_{n=1}^{\infty} \frac{(k > 0)}{\sum_{n=-1}^{-\infty} (k < 0)} \right\} \sqrt{2|kn|} (-f_{-n} \hat{a}_n + f_n \hat{a}_n^\dagger) \right]. \end{aligned} \quad (3.86)$$

It follows that

$$\hat{V}^\dagger(f) \hat{a}_n \hat{V}(f) = \hat{a}_n + \theta(kn) \sqrt{2|kn|} f_n, \quad (3.87)$$

$$\hat{V}^\dagger(f) \hat{a}_n^\dagger \hat{V}(f) = \hat{a}_n^\dagger + \theta(kn) \sqrt{2|kn|} f_{-n}, \quad (3.88)$$

where $\theta(x)$ is 1 when $x > 0$ and 0 when $x < 0$. The definitions (3.81) and (3.85) with the above equations yield

$$\begin{aligned} \hat{V}^\dagger(f) \hat{\chi}(\sigma) \hat{V}(f) &= \sum_{n \neq 0} \frac{1}{\sqrt{2|kn|}} (\hat{a}_n + \theta(kn) \sqrt{2|kn|} f_n) e^{in\sigma} \\ &= \hat{\chi}(\sigma) + f^{(+)}(\sigma), \end{aligned} \quad (3.89)$$

$$\begin{aligned} \hat{V}^\dagger(f) \hat{\chi}^\dagger(\sigma) \hat{V}(f) &= \sum_{n \neq 0} \frac{1}{\sqrt{2|kn|}} (\hat{a}_n^\dagger + \theta(kn) \sqrt{2|kn|} f_{-n}) e^{-in\sigma} \\ &= \hat{\chi}^\dagger(\sigma) + f^{(-)}(\sigma), \end{aligned} \quad (3.90)$$

where the alternative sign is chosen according to the sign of k . Again we arrive at the same result

$$\begin{aligned}\hat{V}^\dagger(f) : e^{i\hat{\varphi}(\sigma)} : \hat{V}(f) &= \hat{V}^\dagger(f) e^{i\hat{\chi}^\dagger(\sigma)} e^{i\hat{\chi}(\sigma)} \hat{V}(f) \\ &= e^{if^{(+)}(\sigma)+if^{(-)}(\sigma)} e^{i\hat{\chi}^\dagger(\sigma)} e^{i\hat{\chi}(\sigma)} \\ &= e^{if(\sigma)} : e^{i\hat{\varphi}(\sigma)} : ,\end{aligned}\quad (3.91)$$

4 Summary and discussion

4.1 Summary

Now let us summarize what have been done in this paper. We have reviewed the quantum mechanics on S^1 originally formulated by Ohnuki and Kitakado. They chose generators respecting topology of S^1 . They defined the algebra and classified its irreducible representations. Inequivalent representations are characterized by a continuous parameter α ($0 < \alpha \leq 1$). Elimination of α is obstructed by nontrivial topology of S^1 .

As a generalization of the quantum mechanics on S^1 , we have proposed the definition of the algebra for the abelian sigma model in (1+1) dimensions. The central extensions are also introduced into the algebra.

The degrees of freedom of the field variable are separated as

$$\text{Map}(S^1; S^1) \cong S^1 \times \mathbb{Z} \times \text{Map}_0(S^1; \mathbb{R}). \quad (4.1)$$

The separation is done as follows. Identify $\phi : S^1 \rightarrow S^1$ with $\phi : S^1 \rightarrow U(1)$. Take a branch of its logarithm and put $\tilde{\varphi}(\sigma) = -i \log \phi(\sigma)$. Define $N \in \mathbb{Z}$ by $2\pi N = \tilde{\varphi}(2\pi) - \tilde{\varphi}(0)$. Next define $e^{i\lambda} \in S^1$ by $2\pi\lambda = \int_0^{2\pi} (\tilde{\varphi}(\sigma) - N\sigma) d\sigma$. Finally define $\varphi : S^1 \rightarrow \mathbb{R}$ by $\varphi(\sigma) = \tilde{\varphi}(\sigma) - \lambda - N\sigma$. Thus $\varphi(2\pi) = \varphi(0)$ and $\int_0^{2\pi} \varphi(\sigma) d\sigma = 0$. Then

$$\phi(\sigma) = e^{i(\lambda + N\sigma + \varphi(\sigma))} \quad (4.2)$$

is a decomposition according to (4.1) and results in (3.15) by putting $U = e^{i\lambda}$. What we have done is to define a coordinate system in the infinite dimensional manifold $Q = \text{Map}(S^1; S^1)$. In the same way we can define a coordinate in the group $\Gamma = \text{Map}(S^1; U(1))$ as given by (3.17). These coordinates are convenient; they are direct product decompositions of the manifold Q , the group Γ and the action of Γ on Q . In other words, these decompositions are preserved under group operation of Γ and are also preserved under the action of Γ on Q as shown in (3.18)-(3.20).

Existence of such coordinates is crucial for construction of the quantum theory. The fundamental relations (3.21)-(3.23) are easy to understand intuitively, however too complicated to construct its concrete representation. The coordinates reduce them to simpler relations (3.37)-(3.39). Even if the central extension exists, other complication is only addition of the anomalous commutator (3.43) and the twist

relation (3.49). Thus we have noticed that the Ohnuki-Kitakado representations and the Fock representation provide representations for our model.

We conclude that inequivalent irreducible representations are parametrized again by α ($0 \leq \alpha < 1$). When there is the central extension, the action of the operator \tilde{W} is changed as (3.62) and half of modes in $\hat{\pi}(\sigma)$ is removed as (3.65).

Exponential of the local operator $\hat{\phi}(\sigma)$ must be regularized by the normal ordering procedure. We have shown that the procedure preserves the fundamental relations but violates unitarity of $e^{i\hat{\phi}(\sigma)}$.

4.2 Discussion

For what kind of physics is our theory applicable? What we have done is just formulation of a rather ideal model. It gives a lesson; *when a model has nontrivial topology, it is possible to construct inequivalent quantum theories, even if they are equivalent as classical theories*. We would like to point out some models which have such possibilities.

The first example is still an ideal model in (1+1) dimensions; it is the sine-Gordon model, whose lagrangian is

$$\mathcal{L} = \frac{1}{2} \partial_\mu \tilde{\varphi}(x) \partial^\mu \tilde{\varphi}(x) + \kappa^2 \cos(\tilde{\varphi}(x)). \quad (4.3)$$

It is a model which has interaction. It can be rewritten by the variables of the abelian sigma model without the central extension by identifying ϕ with $e^{i\tilde{\varphi}}$. Then the corresponding hamiltonian is defined by

$$\begin{aligned} \hat{H} &= \frac{1}{2} \int_0^{2\pi} \left[\left(\frac{1}{2\pi} \hat{P} + \hat{\pi} \right)^2 + \partial \hat{\phi}^\dagger \partial \hat{\phi} - \kappa^2 (\hat{\phi} + \hat{\phi}^\dagger) \right] d\sigma \\ &= \frac{1}{2} \left(\frac{1}{2\pi} \hat{P}^2 + 2\pi \hat{N}^2 \right) + \sum_{n \neq 0} |n| \left(\hat{a}_n^\dagger \hat{a}_n + \frac{1}{2} \right) \\ &\quad - \frac{\kappa^2}{2} \int_0^{2\pi} \left(\hat{U} e^{i(\hat{N}\sigma + \hat{\varphi}(\sigma))} + \hat{U}^\dagger e^{-i(\hat{N}\sigma + \hat{\varphi}(\sigma))} \right) d\sigma. \end{aligned} \quad (4.4)$$

The last term contains highly nonlinear complicated interaction. It also contains interaction between the zero-mode \hat{U} and the fluctuation mode $\hat{\phi}$. This hamiltonian commutes with \hat{N} , hence the winding number is conserved. If the hamiltonian includes the winding operator \hat{W} , change of the winding number can occur. Such a jumping motion is not allowed in classical theory but is possible in quantum theory. The winding number is sometimes called soliton or kink number. It is known [4] that this model has a topological soliton, which behaves like a fermion. However it is still

Table 1: Possible extensions

	abelian sigma model	higher dimensions	nonabelian
base space X	S^1	S^n, T^n, \mathbf{R}^n	
target space M	S^1		$G, G/H, T^n/Z$
group G	$U(1)$		$SU(n), SO(n)$ etc

obscure whether our formulation is relevant to soliton physics or not. It is expected that our model may give an insight to quantum theory of solitons.

Other examples are found in both field theories and string theories. From both points of view, it is hoped to extend our model to nonabelian groups and to higher dimensions. Our model has the field configuration space $Q = \text{Map}(S^1; S^1)$. The group $\Gamma = \text{Map}(S^1; U(1))$ acts on Q simply transitively. The most general model has $Q = \text{Map}(X; M)$, in which X is called a base space and M is a target space. When $\dim X > 1$, we call it a higher dimensional model. If a group G acts on M transitively, M is called a homogeneous space G/H . Then an infinite dimensional group $\Gamma = \text{Map}(X; G)$ acts transitively on $Q = \text{Map}(X; M)$ by pointwise multiplication. When the group G is nonabelian, it is called a nonlinear sigma model. When the base space X is S^1 , it is called a bosonic string model. In addition, when the target space M is an orbifold, for example a toroidal orbifold T^n/Z , it is called an orbifold string model. Directions for extensions are summarized in the table.

We should refer to the known results on nonlinear sigma models in two dimensions. Some of them are exactly solved by the method of factorization theory and the Bethe ansatz [17]. Here "exactly solved" means that the exact S -matrix is obtained and therefore the mass spectrum defined by poles of the S -matrix is also calculated. Wiegmann *et al* [18] have already obtained exact solutions of nonlinear sigma models for the algebras $SO(n+2)$, $SU(n+1)$, $Sp(2n)$ ($n = 1, 2, \dots$). All of them exhibit massive spectra.

Their models are quite different from ours at two points. First, they considered nonabelian sigma models while we consider the abelian sigma model. Second, their models are defined in the noncompact space \mathbf{R} while our model is defined in the compact space S^1 . Hence infrared behavior of their models is utterly different from ours. Their models suffer from infrared divergence and massless particles are ill-defined as stated by Coleman's theorem. Thus only massive particles appear in their models which leads to the symmetric phase. In our model, infrared divergence is

absent because of finite size of the space. Thus massless boson remains well-defined and our model is in the broken phase. Therefore to find a more useful example, it is hoped to extend our consideration to higher dimensional space-times.

In addition, their approach to the problem is also different from ours. In fact they use neither field variables nor lagrangians. They demand some reasonable properties to be satisfied by the S -matrix; unitarity, factorizability, crossing symmetry, analyticity and other symmetries specified by a Lie algebra. In (1+1) dimensions such a requirement determines the S -matrix directly and unambiguously. Actually what they have constructed is a realization of symmetries in terms of S -matrix. But they do not pay attention to topology. Although the model which we have considered is quite simple, that is the $SO(2)$ sigma model, we have shown existence of inequivalent quantizations as a consequence of the nontrivial topology. At present we do not know how to incorporate topology into the algebraic approach of Zamolodchikov and Wiegmann *et al*.

We shall briefly comment upon the known results on orbifold string models [19]. Sakamoto *et al* [20] investigated theories of closed bosonic strings on orbifolds in operator formalism. He have shown that commutators among the zero-mode and winding-mode variables are left ambiguous and that these variables obey nontrivial quantization. The relation between his result and ours is still left obscure.

Finally we would like to suggest a way to explore nonabelian and higher dimensional theories. The decomposition (4.1) heavily relies on the abelian nature of the group $U(1)$. As a generalization to a nonabelian group G , we expect a decomposition

$$\Gamma = \text{Map}(S^n; G) \cong G \times \pi_n(G) \times \text{Map}_0(S^n; \text{Lie}(G)), \quad (4.5)$$

where π_n denotes the n -th homotopy group and

$$\text{Map}_0(S^n; \text{Lie}(G)) = \{g : S^n \rightarrow \text{Lie}(G) \mid C^\infty, \int_{S^n} g = 0\}. \quad (4.6)$$

In the decomposition (4.5), the first component describes the zero-mode, the second one does the topologically disconnected mode, and the third one does the fluctuation mode. Unfortunately such a decomposition does not exist, because the nonabelian nature severely entangles degrees of freedom of Γ . It seems hopeless to find a convenient coordinate in Γ generally. We should take a rather abstract approach to construct a representation for such a complicated group. Gelfand *et al* [21] has investigated the representation theory of the group $\Gamma = \text{Map}(X; G)$ for a base space

$X(\dim X \geq 2)$ and a compact semisimple G . They do not rely on any coordinates but take a quite abstract approach. However they do not consider a manifold $Q = \text{Map}(X; G/H)$ on which Γ acts. We do not know how to incorporate such a configuration space Q into their representation theory.

Acknowledgments

I am indebted to so many people that I cannot record all of their names here. I gratefully acknowledge helpful discussions with T.Kashiwa and S.Sakoda (Kyushu University), K.Higashijima (Osaka University), M.Sakamoto and M.Tachibana (Kobe University), I.Tsutsui and S.Tsujimaru (Institute for Nuclear Study), K.Fujii (Yokohama City University) and Y.Igarashi (Niigata University). I wish to express my gratitude to Y.Ohnuki, S.Kitakado and H.Ikemori, who discussed with me and encouraged me continuously. I thank all members of E-laboratory, who provided me with encouraging environment during the whole period of my studying.

A Appendix

Here we give an explicit calculation of the Fourier expansion of (3.39) and (3.43) to derive (3.64), (3.65) and (3.55). We repeat the assumptions;

$$[\hat{\varphi}(\sigma), \hat{\varphi}(\sigma')] = 0, \quad (\text{A.1})$$

$$[\hat{\varphi}(\sigma), \hat{\pi}(\sigma')] = i\left(\delta(\sigma - \sigma') - \frac{1}{2\pi}\right), \quad (\text{A.2})$$

$$[\hat{\pi}(\sigma), \hat{\pi}(\sigma')] = -\frac{ik}{\pi}\delta'(\sigma - \sigma'), \quad (\text{A.3})$$

with constraints (3.30) and (3.32). k is assumed to be a non-zero integer. We define

$$\hat{\varphi}_m = \frac{1}{2\pi} \int_0^{2\pi} \hat{\varphi}(\sigma) e^{-im\sigma} d\sigma, \quad (\text{A.4})$$

$$\hat{\pi}_n = \frac{1}{2\pi} \int_0^{2\pi} \hat{\pi}(\sigma) e^{-in\sigma} d\sigma, \quad (\text{A.5})$$

for integers m, n . Obviously, $\hat{\varphi}_m^\dagger = \hat{\varphi}_{-m}$, $\hat{\pi}_n^\dagger = \hat{\pi}_{-n}$ and $\hat{\varphi}_0 = \hat{\pi}_0 = 0$. Multiplying $(2\pi)^{-2} e^{-im\sigma - in\sigma'}$ to (A.1)-(A.3) and integrating $\int_0^{2\pi} \int_0^{2\pi} d\sigma d\sigma'$, we obtain

$$[\hat{\varphi}_m, \hat{\varphi}_n] = 0, \quad (\text{A.6})$$

$$[\hat{\varphi}_m, \hat{\pi}_n] = \frac{i}{2\pi}(\delta_{m+n,0} - \delta_{m,0}\delta_{n,0}), \quad (\text{A.7})$$

$$[\hat{\pi}_m, \hat{\pi}_n] = \frac{k}{2\pi^2} m \delta_{m+n,0}. \quad (\text{A.8})$$

We put

$$\hat{b}_n = i\sqrt{\frac{2\pi^2}{|k|n}} \hat{\pi}_n, \quad \hat{b}_n^\dagger = -i\sqrt{\frac{2\pi^2}{|k|n}} \hat{\pi}_{-n}, \quad (n > 0) \quad (\text{A.9})$$

then (A.8) implies

$$[\hat{b}_m, \hat{b}_n] = 0, \quad (\text{A.10})$$

$$[\hat{b}_m, \hat{b}_n^\dagger] = \epsilon(k) \delta_{mn}, \quad (m, n > 0) \quad (\text{A.11})$$

where $\epsilon(k) = k/|k|$. By definition we have

$$\hat{\pi}_n = -i\sqrt{\frac{|k|n}{2\pi^2}} \hat{b}_n, \quad \hat{\pi}_{-n} = i\sqrt{\frac{|k|n}{2\pi^2}} \hat{b}_n^\dagger, \quad (n > 0) \quad (\text{A.12})$$

hence

$$\begin{aligned} \hat{\pi}(\sigma) &= \sum_{n \neq 0} \hat{\pi}_n e^{in\sigma} \\ &= \frac{i}{2\pi} \sum_{n=1}^{\infty} \sqrt{2|k|n} (-\hat{b}_n e^{in\sigma} + \hat{b}_n^\dagger e^{-in\sigma}). \end{aligned} \quad (\text{A.13})$$

On the other hand, for non-zero m and n , (A.6) and (A.7) yield

$$[\hat{\varphi}_m - \frac{i\pi}{km} \hat{\pi}_m, \hat{\pi}_n] = \frac{i}{2\pi} \delta_{m+n,0} - \frac{i\pi}{km} \frac{km}{2\pi^2} \delta_{m+n,0} = 0 \quad (\text{A.14})$$

and also

$$[\hat{\varphi}_m - \frac{i\pi}{km} \hat{\pi}_m, \hat{\varphi}_n - \frac{i\pi}{kn} \hat{\pi}_n] = \frac{1}{2kn} \delta_{m+n,0}. \quad (\text{A.15})$$

Therefore if we put

$$\begin{cases} \hat{b}_{-n} = \epsilon(k) \sqrt{2|k|n} (\hat{\varphi}_{-n} + \frac{i\pi}{kn} \hat{\pi}_{-n}), \\ \hat{b}_n^\dagger = \epsilon(k) \sqrt{2|k|n} (\hat{\varphi}_n - \frac{i\pi}{kn} \hat{\pi}_n), \end{cases} \quad (n > 0) \quad (\text{A.16})$$

they satisfy

$$[\hat{b}_m, \hat{b}_n] = 0, \quad (\text{A.17})$$

$$[\hat{b}_m, \hat{b}_n^\dagger] = \epsilon(k) \delta_{mn}. \quad (m, n \neq 0) \quad (\text{A.18})$$

It is easily seen that

$$\hat{\varphi}_n = \frac{\epsilon(k)}{\sqrt{2|k|n}} \hat{b}_{-n}^\dagger + \frac{i\pi}{kn} \hat{\pi}_n = \frac{\epsilon(k)}{\sqrt{2|k|n}} (\hat{b}_n + \hat{b}_{-n}^\dagger), \quad (\text{A.19})$$

$$\hat{\varphi}_{-n} = \frac{\epsilon(k)}{\sqrt{2|k|n}} \hat{b}_{-n} - \frac{i\pi}{kn} \hat{\pi}_{-n} = \frac{\epsilon(k)}{\sqrt{2|k|n}} (\hat{b}_{-n} + \hat{b}_n^\dagger), \quad (n > 0) \quad (\text{A.20})$$

using (A.16) with (A.12). Thus we have

$$\begin{aligned} \hat{\varphi}(\sigma) &= \sum_{n \neq 0} \hat{\varphi}_n e^{in\sigma} \\ &= \epsilon(k) \sum_{n \neq 0} \frac{1}{\sqrt{2|kn|}} (\hat{b}_n e^{in\sigma} + \hat{b}_n^\dagger e^{-in\sigma}). \end{aligned} \quad (\text{A.21})$$

Finally we define

$$\hat{a}_n = \begin{cases} \hat{b}_n & (k > 0) \\ -\hat{b}_{-n}^\dagger & (k < 0) \end{cases} \quad \hat{a}_n^\dagger = \begin{cases} \hat{b}_n^\dagger & (k > 0) \\ -\hat{b}_{-n} & (k < 0) \end{cases} \quad (\text{A.22})$$

to get

$$[\hat{a}_m, \hat{a}_n^\dagger] = \delta_{mn}, \quad (\text{A.23})$$

$$\hat{\varphi}(\sigma) = \sum_{n \neq 0} \frac{1}{\sqrt{2|kn|}} (\hat{a}_n e^{in\sigma} + \hat{a}_n^\dagger e^{-in\sigma}), \quad (\text{A.24})$$

$$\hat{\pi}(\sigma) = \frac{i}{2\pi} \left\{ \begin{array}{ll} \sum_{n=1}^{\infty} & (k > 0) \\ \sum_{n=-1}^{-\infty} & (k < 0) \end{array} \right\} \sqrt{2|kn|} (-\hat{a}_n e^{in\sigma} + \hat{a}_n^\dagger e^{-in\sigma}), \quad (\text{A.25})$$

which are the desired results (3.55), (3.64) and (3.65).

References

- [1] Y. Ohnuki and S. Kitakado, "On quantum mechanics on compact space", *Mod. Phys. Lett. A* **7** (1992) 2477; "Fundamental algebra for quantum mechanics on S^D and gauge potentials", *J. Math. Phys.* **34** (1993) 2827
- [2] M. Bando, T. Kugo and K. Yamawaki, "Nonlinear realization and hidden local symmetries", *Phys. Rep.* **164** (1988) 217
- [3] G. W. Mackey, "Induced representations of groups and quantum mechanics", W. A. Benjamin, INC. New York (1968);
H. D. Doebner and J. Tolar, "Quantum mechanics on homogeneous spaces", *J. Math. Phys.* **16** (1975) 975;
N. P. Landsman and N. Linden, "The geometry of inequivalent quantizations", *Nucl. Phys. B* **365** (1991) 121;
S. Tanimura, "Quantum mechanics on manifolds", Nagoya University preprint DPNU-93-21 (1993);
D. McMullan and I. Tsutsui, "On the emergence of gauge structures and generalized spin when quantizing on a coset space", Dublin Institute for Advanced Studies preprint DIAS-STP-93-14 (1993)
- [4] S. Coleman, "Quantum sine-Gordon equation as the massive Thirring model", *Phys. Rev. D* **11** (1975) 2088;
S. Mandelstam, "Soliton operators for the quantized sine-Gordon equation", *Phys. Rev. D* **11** (1975) 3026;
G. I. Japaridze, A. A. Nersesyan and P. B. Wiegmann, "Regularized integrable version of the one-dimensional quantum sine-Gordon model", *Phys. Scripta* **27** (1983) 5;
S. G. Rajeev, "Fermions from bosons in 3+1 dimensions through anomalous commutators", *Phys. Rev. D* **29** (1984) 2944
- [5] R. Jackiw and C. Rebbi, "Vacuum periodicity in a Yang-Mills quantum theory", *Phys. Rev. Lett.* **37** (1976) 172;
C. G. Callan Jr., R. F. Dashen and D. J. Gross, "The structure of the gauge theory vacuum", *Phys. Lett.* **63B** (1976) 334

- [6] E. Witten, "Global aspects of current algebra", *Nucl. Phys.* **B223** (1983) 422;
"Non-abelian bosonization in two dimensions", *Commun. Math. Phys.* **92** (1984) 455
- [7] V. N. Gribov, "Quantization of non-abelian gauge theories", *Nucl. Phys.* **B139** (1978) 1;
I. M. Singer, "Some remarks on the Gribov ambiguity", *Commun. Math. Phys.* **60** (1978) 7; "The geometry of the orbit space for non-abelian gauge theories", *Phys. Scripta* **24** (1981) 817
- [8] S. Tanimura, "Topology and quantization of abelian sigma model in (1+1) dimensions", *Phys. Lett.* **340B** (1994) 57
- [9] S. Tanimura, "Gauge field, parity and uncertainty relation of quantum mechanics on S^1 ", *Prog. Theor. Phys.* **90** (1993) 271
- [10] Y. Ohnuki, M. Suzuki and T. Kashiwa, "Method of path integrals" (in Japanese) Iwanami, Tokyo (1992)
- [11] L. Schulman, "A path integral for spin", *Phys. Rev.* **176** (1968), 1558;
M.G.G. Laidlaw and C.M. DeWitt, "Feynman functional integrals for systems of indistinguishable particles", *Phys. Rev.* **D3** (1971), 1375
- [12] L. D. Faddeev, "Operator anomaly for the Gauss law", *Phys. Lett.* **145B** (1984) 81;
Y-S. Wu, "Cohomology of gauge fields and explicit construction of Wess-Zumino lagrangians in nonlinear sigma models over G/H ", *Phys. Lett.* **153B** (1985) 70;
J. Mickelsson, "Kac-Moody groups, topology of the Dirac determinant bundle, and fermionization", *Commun. Math. Phys.* **110** (1987) 173;
K. Fujii and M. Tanaka, "Universal Schwinger cocycles of current algebras in $(D+1)$ -dimensions : geometry and physics", *Commun. Math. Phys.* **129** (1990) 267
- [13] P. Goddard and D. Olive, "Kac-Moody and Virasoro algebras in relation to quantum physics", *Int. J. Mod. Phys.* **A1** (1986) 303

- [14] G. Segal, "Unitary representations of some infinite dimensional groups", *Commun. Math. Phys.* **80** (1981) 301;
A. Pressley and G. Segal, "Loop groups", Oxford University Press, New York (1986)
- [15] S. Coleman, "There are no Goldstone bosons in two dimensions", *Commun. Math. Phys.* **31** (1973) 259
- [16] S. Kamefuchi, L. O'Raifeartaigh and A. Salam, "Change of variables and equivalence theorems in quantum field theories", *Nucl. Phys.* **28** (1961) 529
- [17] A. B. Zamolodchikov and Al. B. Zamolodchikov, "Factorized S -matrix in two dimensions as the exact solutions of certain relativistic quantum field theory models", *Ann. Phys.* **120** (1979) 253;
A. A. Belavin, "Exact solution of the two-dimensional model with asymptotic freedom", *Phys. Lett.* **87B** (1979) 117;
A. Polyakov and P. B. Wiegmann, "Theory of nonabelian Goldstone bosons in two dimensions", *Phys. Lett.* **131B** (1983) 121;
V. A. Fateev, V. A. Kazakov and P. B. Wiegmann, "Principal chiral field at large N ", *Nucl. Phys.* **B424** (1994) 505
- [18] E. Ogievetsky, N. Reshetikhin and P. Wiegmann, "The principal chiral field in two dimensions on classical Lie algebras", *Nucl. Phys.* **B280** (1987) 45;
- [19] K. S. Narain, M. H. Sarmadi and E. Witten, "A note on toroidal compactification of heterotic string theory", *Nucl. Phys.* **B279** (1987) 369
- [20] J. O. Madsen and M. Sakamoto, "Toroidal orbifold models with a Wess-Zumino term", *Phys. Lett.* **B322** (1994) 91;
M. Sakamoto, "Topological aspects of antisymmetric background field on orbifolds", *Nucl. Phys.* **B414** (1994) 267
- [21] I. M. Gelfand, M. I. Graev and A. M. Vershik, "Representations of the group of smooth mappings of a manifold X into a compact Lie group", *Compositio Math.* **35** (1977) 299; "Representations of the groups of functions taking values in a compact Lie group", *Compositio Math.* **42** (1981) 217;

副論文

1. S. Tanimura, "Gauge field, parity and uncertainty relation of quantum mechanics on S^1 ", *Prog. Theor. Phys.* **90** (1993) 271-291.
2. S. Tanimura, "Topology and quantization of abelian sigma model in (1+1) dimensions", *Phys. Lett.* **340B** (1994) 57-62.