

FUNCTIONAL RELATIONS AND SPECIAL VALUES OF MORDELL-TORNHEIM TRIPLE ZETA AND L-FUNCTIONS

KOHJI MATSUMOTO, TAKASHI NAKAMURA, AND HIROFUMI TSUMURA

ABSTRACT. In this paper, we prove the existence of meromorphic continuation of certain triple zeta-functions of Lerch's type. Based on this result, we prove some functional relations for triple zeta and L -functions of the Mordell-Tornheim type. Using these functional relations, we prove new explicit evaluation formulas for special values of these functions. These can be regarded as triple analogues of known results for double zeta and L -functions.

1. INTRODUCTION

Let \mathbb{N} be the set of natural numbers, $\mathbb{N}_0 = \mathbb{N} \cup \{0\}$, \mathbb{Z} the ring of rational integers, \mathbb{Q} the field of rational numbers, \mathbb{R} the field of real numbers, and \mathbb{C} the field of complex numbers.

In this paper, we study triple zeta and L -functions defined by

$$(1.1) \quad \zeta_{MT,3}(s_1, s_2, s_3, s_4; \alpha, \beta, \gamma, \delta) := \sum_{l,m,n=1}^{\infty} \frac{e^{2\pi i l \alpha} e^{2\pi i m \beta} e^{2\pi i n \gamma} e^{2\pi i(l+m+n)\delta}}{l^{s_1} m^{s_2} n^{s_3} (l+m+n)^{s_4}},$$

$$(1.2) \quad L_{MT,3}(s_1, s_2, s_3, s_4; \varphi, \chi, \psi, \omega) := \sum_{l,m,n=1}^{\infty} \frac{\varphi(l) \chi(m) \psi(n) \omega(l+m+n)}{l^{s_1} m^{s_2} n^{s_3} (l+m+n)^{s_4}},$$

where $i = \sqrt{-1}$, $\alpha, \beta, \gamma, \delta \in \mathbb{R}$ and $\varphi, \chi, \psi, \omega$ are primitive Dirichlet characters. These are convergent absolutely when $\Re s_j \geq 1$ ($1 \leq j \leq 4$). These are triple analogues of Tornheim's double series $T(a, b, c) = \sum_{m,n \geq 1} m^{-a} n^{-b} (m+n)^{-c}$ ([11]) which was also studied by Mordell independently ([6]).

The first-named author ([3]) analytically studied $T(s_1, s_2, s_3)$ for $(s_1, s_2, s_3) \in \mathbb{C}^3$, and defined more generally a multiple analogue of $T(s_1, s_2, s_3)$, what is called the Mordell-Tornheim multiple zeta-functions denoted by $\zeta_{MT,r}(s_1, \dots, s_r, s_{r+1})$ ([4]). In particular, $\zeta_{MT,2}(s_1, s_2, s_3) = T(s_1, s_2, s_3)$ and $\zeta_{MT,3}(s_1, s_2, s_3, s_4) = \zeta_{MT,3}(s_1, s_2, s_3, s_4; 1, 1, 1, 1)$.

The double series $T(a, b, c)$ and its χ -analogue have been studied extensively. For instance, evaluation formulas for special values of them have been studied by a lot of authors, and functional relations for them have been obtained in [7, 8, 17, 18]. However, there are very few papers in which properties of $\zeta_{MT,r}$ for $r \geq 3$ have been studied. Recently Ochiai and the authors studied $\zeta_{MT,3}(s_1, s_2, s_3, s_4)$ and relevant functions, and proved some functional relations for them ([5]).

In this paper we aim to study analytic properties of (1.1) and (1.2), and to prove certain functional relations for them. The results in this paper can be regarded as triple analogues of the known results for double zeta and L -functions proved in the papers quoted above and [9, 10, 13, 14, 15], and also be regarded as χ -analogues of the results in [5]. Therefore

2000 *Mathematics Subject Classification.* Primary 40B05; Secondly 11M35, 11M06, 33E20.

Key words and phrases. Triple zeta and L -functions, Tornheim's double series, Riemann's zeta-function, Dirichlet L -functions, functional relations.

this paper is a step toward the general theory of functional relations for Mordell-Tornheim r -ple zeta and L -functions including known results of r -ple zeta values given in [16].

2. ANALYTIC CONTINUATIONS OF CERTAIN TRIPLE ZETA AND L -FUNCTIONS

The aim of this section is to prove the following proposition by using the same method as in [3, 4].

Proposition 2.1. *The triple zeta-function (1.1) and the triple L -function (1.2) can be continued meromorphically to the whole \mathbb{C}^4 space.*

Proof. Since

$$e^{2\pi il\alpha} e^{2\pi im\beta} e^{2\pi in\gamma} e^{2\pi i(l+m+n)\delta} = e^{2\pi il(\alpha+\delta)} e^{2\pi im(\beta+\delta)} e^{2\pi in(\gamma+\delta)},$$

we may assume $\delta = 0$ without loss of generality in the proof of (1.1). We use the notation

$$\zeta_{MT,2}(s_1, s_2, s_3; \alpha, \beta, \gamma) := \sum_{m_1, m_2=1}^{\infty} \frac{e^{2\pi im_1\alpha} e^{2\pi im_2\beta} e^{2\pi i(m_1+m_2)\gamma}}{m_1^{s_1} m_2^{s_2} (m_1 + m_2)^{s_3}}.$$

First we assume $\Re s_j > 1$ ($1 \leq j \leq 4$). We recall the Mellin-Barnes formula

$$(2.1) \quad (1 + \lambda)^{-s} = \frac{1}{2\pi i} \int_{(c)} \frac{\Gamma(s+z)\Gamma(-z)}{\Gamma(s)} \lambda^z dz,$$

where s, λ are complex numbers with $\Re s > 0$, $|\arg \lambda| < \pi$, $\lambda \neq 0$, c is real with $-\Re s < c < 0$, and the path (c) of integration is the vertical line $\Re z = c$. Since

$$(l + m + n)^{-s_4} = (l + m)^{-s_4} \cdot \frac{1}{2\pi i} \int_{(c)} \frac{\Gamma(s_4+z)\Gamma(-z)}{\Gamma(s_4)} \left(\frac{n}{l+m}\right)^z dz,$$

where $-\Re s_4 < c < 0$, (1.1) with $\delta = 0$ can be written as

$$\frac{1}{2\pi i} \int_{(c)} \frac{\Gamma(s_4+z)\Gamma(-z)}{\Gamma(s_4)} \zeta_{MT,2}(s_1, s_2, s_4+z; \alpha, \beta, 0) \phi(s_3 - z, \gamma) dz,$$

where $\phi(s, \gamma) = \sum_{n \geq 1} e^{2\pi in\gamma} n^{-s}$. Furthermore, by using

$$(l + m)^{-s_4} = l^{-s_4} \cdot \frac{1}{2\pi i} \int_{(c')} \frac{\Gamma(s_4+z')\Gamma(-z')}{\Gamma(s_4)} \left(\frac{m}{l}\right)^{z'} dz',$$

we have

$$(2.2) \quad \begin{aligned} \zeta_{MT,2}(s_1, s_2, s_4; \alpha, \beta, 0) &= \frac{1}{2\pi i} \int_{(c')} \frac{\Gamma(s_4+z')\Gamma(-z')}{\Gamma(s_4)} \\ &\quad \times \phi(s_1 + s_4 + z', \alpha) \phi(s_2 - z', \beta) dz', \end{aligned}$$

where $-\Re s_4 < c' < 0$. We set $\nu(x) := 1$ (resp. 0) if $x \in \mathbb{Z}$ (resp. $x \notin \mathbb{Z}$). Then we can determine the poles of the integrand on the right-hand side of (2.2) as

$$z' = -s_4 - k, \quad z' = k \quad (k \in \mathbb{N}_0); \quad \nu(\alpha)(z' = 1 - s_1 - s_4), \quad \nu(\beta)(z' = s_2 - 1).$$

In fact, the former two expressions come from the Γ -factors, and the latter two expressions come from the ϕ -factors.

Now we shift the path $\Re z' = c'$ to $\Re z' = M - \varepsilon$ for a sufficiently large $M \in \mathbb{N}$ and a sufficiently small positive $\varepsilon \in \mathbb{R}$. Then the relevant poles through this shifting are $z' = s_2 - 1$ and $z' = k$ ($0 \leq k \leq M - 1$). Counting their residues, we can obtain

$$\begin{aligned} \zeta_{MT,2}(s_1, s_2, s_4; \alpha, \beta, 0) &= \nu(\beta) \frac{\Gamma(s_2 + s_4 - 1)\Gamma(1 - s_2)}{\Gamma(s_4)} \phi(s_1 + s_2 + s_4 - 1, \alpha) \\ &+ \sum_{k=0}^{M-1} \binom{-s_4}{k} \phi(s_1 + s_4 + k, \alpha) \phi(s_2 - k, \beta) \\ &+ \frac{1}{2\pi i} \int_{(M-\varepsilon)} \frac{\Gamma(s_4 + z')\Gamma(-z')}{\Gamma(s_4)} \phi(s_1 + s_4 + z', \alpha) \phi(s_2 - z', \beta) dz'. \end{aligned}$$

Similarly to our previous results [3, 4, 5], this gives the meromorphic continuation of $\zeta_{MT,2}(s_1, s_2, s_4; \alpha, \beta, 0)$ and determines its singularities as

$$\nu(\alpha)(s_1 + s_4 = 1 - k), \quad \nu(\beta)(s_2 + s_4 = 1 - k), \quad \nu(\alpha)\nu(\beta)(s_1 + s_2 + s_4 = 2).$$

Hence singularities of $\zeta_{MT,2}(s_1, s_2, s_4 + z; \alpha, \beta, 0)\phi(s_3 - z, \gamma)$ are

$$\begin{aligned} \nu(\alpha)(s_1 + s_4 + z = 1 - k), \quad \nu(\beta)(s_2 + s_4 + z = 1 - k), \\ \nu(\alpha)\nu(\beta)(s_1 + s_2 + s_4 + z = 2), \quad \nu(\gamma)(s_3 - z = 1). \end{aligned}$$

These expressions can be rewritten as $z = -s_1 - s_4 + 1 - k$, $z = -s_2 - s_4 + 1 - k$, $z = 2 - s_1 - s_2 - s_4$, $z = s_3 - 1$. Therefore, when we shift the path $\Re z = c$ to $\Re z = M - \varepsilon$ for a sufficiently large $M \in \mathbb{N}$ and a sufficiently small positive $\varepsilon \in \mathbb{R}$, we have only to consider the poles $z = k$ and $z = s_3 - 1$ in the case $\nu(\gamma) = 1$. Hence we have

$$\begin{aligned} \zeta_{MT,3}(s_1, s_2, s_3, s_4; \alpha, \beta, \gamma, 0) &= -\frac{\Gamma(s_3 + s_4 - 1)\Gamma(1 - s_3)}{\Gamma(s_4)} \times \text{Res}_{z=s_3-1}(\zeta_{MT,2}(s_1, s_2, s_4 + z; \alpha, \beta, 0)\phi(s_3 - z, \gamma)) \\ &+ \sum_{k=0}^{M-1} \binom{-s_4}{k} \zeta_{MT,2}(s_1, s_2, s_4 + k; \alpha, \beta, 0)\phi(s_3 - k, \gamma) \\ &+ \frac{1}{2\pi i} \int_{(M-\varepsilon)} \frac{\Gamma(s_4 + z)\Gamma(-z)}{\Gamma(s_4)} \zeta_{MT,2}(s_1, s_2, s_4 + z; \alpha, \beta, 0)\phi(s_3 - z, \gamma) dz. \end{aligned}$$

This gives the meromorphic continuation of (1.1).

As for (1.2), we denote the conductors of $\varphi, \chi, \psi, \omega$ by f_1, f_2, f_3, f_4 , respectively. It follows from [19, Lemma 4.7] that (1.2) can be written as

$$\begin{aligned} &\frac{1}{\tau(\bar{\varphi})\tau(\bar{\chi})\tau(\bar{\psi})\tau(\bar{\omega})} \sum_{j_1=1}^{f_1} \sum_{j_2=1}^{f_2} \sum_{j_3=1}^{f_3} \sum_{j_4=1}^{f_4} \bar{\varphi}(j_1)\bar{\chi}(j_2)\bar{\psi}(j_3)\bar{\omega}(j_4) \\ &\times \sum_{l,m,n=1}^{\infty} \frac{e^{2\pi il(j_1/f_1+j_4/f_4)} e^{2\pi im(j_2/f_2+j_4/f_4)} e^{2\pi in(j_3/f_3+j_4/f_4)}}{l^{s_1} m^{s_2} n^{s_3} (l+m+n)^{s_4}}, \end{aligned}$$

where $\tau(\cdot)$ is the Gauss sum. Hence, from the assertion for (1.1), we obtain the meromorphic continuation of (1.2). This completes the proof of Proposition 2.1. \square

3. FUNCTIONAL RELATIONS FOR TRIPLE ZETA AND L -FUNCTIONS

In this section, we use the same method as in [7]. We denote by $\{B_j(x)\}$ the Bernoulli polynomials defined by $te^{xt}/(e^t - 1) = \sum_{j \geq 0} B_j(x)t^j/j!$ ($|t| < 2\pi$). It is known (see [1, p.266, (22) and p.267, (24)]) that

$$(3.1) \quad B_{2j} := B_{2j}(0) = (-1)^{j+1} 2(2j)!(2\pi)^{-2j} \zeta(2j) \quad (j \in \mathbb{N}_0),$$

$$(3.2) \quad B_j(x - [x]) = -\frac{j!}{(2\pi i)^j} \lim_{K \rightarrow \infty} \sum_{k=-K, k \neq 0}^K \frac{e^{2\pi i k x}}{k^j} \quad (j \in \mathbb{N}),$$

where $\zeta(s)$ is the Riemann zeta-function and $[\cdot]$ is the integer part. Especially, we can see $\zeta(0) = -1/2$ by (3.1). For $k \in \mathbb{Z}$, $j \in \mathbb{N}$ we have

$$(3.3) \quad \int_0^1 e^{-2\pi i k x} B_j(x) dx = \begin{cases} 0 & (k = 0), \\ -(2\pi i k)^{-j} j! & (k \neq 0). \end{cases}$$

In fact the case of $k = 0$ is obvious, and in the case of $k \neq 0$, we have (3.3) by using (3.2). Next we quote [1, p.276 19.(b)], for $p, q \geq 1$, which is

$$(3.4) \quad B_p(x)B_q(x) = \sum_{k=0}^{\max(p,q)/2} \left\{ p \binom{q}{2k} + q \binom{p}{2k} \right\} \frac{B_{2k} B_{p+q-2k}(x)}{p+q-2k} - (-1)^p \frac{p!q!}{(p+q)!} B_{p+q}.$$

Definition 3.1. Let $\zeta(s)$ and $L(s, \chi)$ be the Riemann zeta-function and the Dirichlet L -function, respectively. Let $\alpha, \beta, \gamma, \delta \in \mathbb{R}$ and $\varphi, \chi, \psi, \omega$ be Dirichlet characters. For $s_1, s_2, s_3, s_4 \in \mathbb{C}$ with $\Re s_j \geq 1$ ($1 \leq j \leq 4$), we define

$$\begin{aligned} L_{MT,2}(s_1, s_2, s_3; \varphi, \chi, \psi) &:= \sum_{m_1, m_2=1}^{\infty} \frac{\varphi(m_1)\chi(m_2)\psi(m_1+m_2)}{m_1^{s_1} m_2^{s_2} (m_1+m_2)^{s_3}}; \\ S_j(s_1, s_2, s_3, s_4; \alpha, \beta, \gamma, \delta) &:= \sum_{\substack{m_1, m_2, m_3, m_4=1 \\ m_1+m_j=m_k+m_l}}^{\infty} \frac{e^{2\pi i m_1 \alpha} e^{2\pi i m_2 \beta} e^{2\pi i m_3 \gamma} e^{2\pi i m_4 \delta}}{m_1^{s_1} m_2^{s_2} m_3^{s_3} m_4^{s_4}}; \\ T_j(s_1, s_2, s_3, s_4; \varphi, \chi, \psi, \omega) &:= \sum_{\substack{m_1, m_2, m_3, m_4=1 \\ m_1+m_j=m_k+m_l}}^{\infty} \frac{\varphi(m_1)\chi(m_2)\psi(m_3)\omega(m_4)}{m_1^{s_1} m_2^{s_2} m_3^{s_3} m_4^{s_4}}, \end{aligned}$$

where $\{j, k, l\} = \{2, 3, 4\}$. Note that these functions are absolutely convergent in this region. In fact, putting $m_1 + m_2 = m_3 + m_4 = n$, we can see that

$$\begin{aligned} \sum_{\substack{m_1, m_2, m_3, m_4=1 \\ m_1+m_2=m_3+m_4}}^{\infty} \frac{1}{m_1 m_2 m_3 m_4} &= \sum_{n=1}^{\infty} \left(\sum_{m=1}^{n-1} \frac{1}{m(n-m)} \right)^2 \\ &= \sum_{n=1}^{\infty} \frac{1}{n^2} \left(\sum_{m=1}^{n-1} \left(\frac{1}{m} + \frac{1}{n-m} \right) \right)^2 = 8 \sum_{1 \leq l < m < n} \frac{1}{lmn^2} + 4 \sum_{1 \leq m < n} \frac{1}{m^2 n^2} < \infty. \end{aligned}$$

For $a, b \in \mathbb{N}$ and $s_3, s_4 \in \mathbb{C}$ with $\Re s_j \geq 1$ ($j = 3, 4$), we define

$$(3.5) \quad \begin{aligned} M_1(a, b, s_3, s_4; \gamma, \delta) &:= \frac{2}{a! b!} \sum_{k=0}^{\max(a,b)/2} \left\{ a \binom{b}{2k} + b \binom{a}{2k} \right\} (a+b-2k-1)!(2k)! \\ &\quad \times \zeta(2k) \zeta_{MT,2}(s_3, s_4, a+b-2k; \gamma, \delta, 1); \end{aligned}$$

$$\begin{aligned}
M_2(a, b, s_3, s_4; \gamma, \delta) &:= \frac{2}{a! b!} \sum_{k=0}^{\max(a,b)/2} \left\{ a \binom{b}{2k} + b \binom{a}{2k} \right\} (a+b-2k-1)! (2k)! \\
&\quad \times \zeta(2k) \zeta_{MT,2}(a+b-2k, s_4, s_3; 1, \delta, \gamma) \\
(3.6) \quad &+ \frac{2(-1)^{a+b}}{a! b!} \sum_{k=0}^{\max(a,b)/2} \left\{ a \binom{b}{2k} + b \binom{a}{2k} \right\} (a+b-2k-1)! (2k)! \\
&\quad \times \zeta(2k) \zeta_{MT,2}(a+b-2k, s_3, s_4; 1, \gamma, \delta) \\
&+ (-1)^{a+1} \frac{(2\pi i)^{a+b} B_{a+b}}{(a+b)!} \phi(s_3 + s_4, \gamma + \delta).
\end{aligned}$$

Similarly, for any ψ, ω , we define $N_1(a, b, s_3, s_4; \psi, \omega)$ as a function given by replacing $\zeta_{MT,2}(s_3, s_4, a+b-2k; \gamma, \delta, 1)$ by $L_{MT,2}(s_3, s_4, a+b-2k; \psi, \omega, \mathbf{1})$ on the right-hand side of (3.5), where $\mathbf{1}$ is the trivial character. Furthermore we define $N_2(a, b, s_3, s_4; \psi, \omega)$ as that given by replacing $\zeta_{MT,2}(a+b-2k, s_4, s_3; 1, \delta, \gamma)$, $\zeta_{MT,2}(a+b-2k, s_3, s_4; 1, \gamma, \delta)$ and $\phi(s_3 + s_4, \gamma + \delta)$ by $L_{MT,2}(a+b-2k, s_4, s_3; \mathbf{1}, \omega, \psi)$, $L_{MT,2}(a+b-2k, s_3, s_4; \mathbf{1}, \psi, \omega)$ and $L(s_3 + s_4, \psi\omega)$ on the right-hand side of (3.6), respectively.

Note that, similarly to the proof of Proposition 2.1, by using the Mellin-Barnes formula (2.1), we can prove the existence of meromorphic continuation of $\zeta_{MT,2}(s_1, s_2, s_3; \alpha, \beta, \gamma)$ and $L_{MT,2}(s_1, s_2, s_3; \varphi, \chi, \psi)$, and furthermore of M_1, M_2, N_1, N_2 in Definition 3.1.

Lemma 3.2. *For $a, b \in \mathbb{N}$, $\gamma, \delta \in \mathbb{R}$ and $s_3, s_4 \in \mathbb{C}$ with $\Re(s_3) \geq 1$, $\Re(s_4) \geq 1$, we have*

$$\begin{aligned}
(3.7) \quad &(-1)^b \zeta_{MT,3}(b, s_3, s_4, a; 1, \gamma, \delta, 1) + (-1)^a \zeta_{MT,3}(s_3, s_4, a, b; \gamma, \delta, 1, 1) \\
&+ S_2(a, b, s_3, s_4; 1, 1, \gamma, \delta) = M_1(a, b, s_3, s_4; \gamma, \delta)
\end{aligned}$$

and

$$\begin{aligned}
(3.8) \quad &\zeta_{MT,3}(s_4, a, b, s_3; \delta, 1, 1, \gamma) + (-1)^a S_3(a, b, s_3, s_4; 1, 1, \gamma, \delta) \\
&+ (-1)^b S_4(a, b, s_3, s_4; 1, 1, \gamma, \delta) + (-1)^{a+b} \zeta_{MT,3}(a, b, s_3, s_4; 1, 1, \gamma, \delta) \\
&= M_2(a, b, s_3, s_4; \gamma, \delta).
\end{aligned}$$

Proof. We have

$$\begin{aligned}
&\lim_{K \rightarrow \infty} \int_0^1 \sum_{k=1}^K \frac{e^{2\pi i k x}}{k^a} \sum_{l=1}^K \frac{e^{2\pi i l x}}{l^b} \sum_{m=1}^K \frac{e^{2\pi i m(x+\gamma)}}{m^{s_3}} \sum_{n=1}^K \frac{e^{2\pi i n(x+\delta)}}{n^{s_4}} dx = 0; \\
&\lim_{K \rightarrow \infty} \int_0^1 \sum_{k=-K}^{-1} \frac{e^{2\pi i k x}}{k^a} \sum_{l=1}^K \frac{e^{2\pi i l x}}{l^b} \sum_{m=1}^K \frac{e^{2\pi i m(x+\gamma)}}{m^{s_3}} \sum_{n=1}^K \frac{e^{2\pi i n(x+\delta)}}{n^{s_4}} dx \\
&= (-1)^a \zeta_{MT,3}(b, s_3, s_4, a; 1, \gamma, \delta, 1); \\
&\lim_{K \rightarrow \infty} \int_0^1 \sum_{k=1}^K \frac{e^{2\pi i k x}}{k^a} \sum_{l=-K}^{-1} \frac{e^{2\pi i l x}}{l^b} \sum_{m=1}^K \frac{e^{2\pi i m(x+\gamma)}}{m^{s_3}} \sum_{n=1}^K \frac{e^{2\pi i n(x+\delta)}}{n^{s_4}} dx \\
&= (-1)^b \zeta_{MT,3}(s_3, s_4, a, b; \gamma, \delta, 1, 1); \\
&\lim_{K \rightarrow \infty} \int_0^1 \sum_{k=-K}^{-1} \frac{e^{2\pi i k x}}{k^a} \sum_{l=-K}^{-1} \frac{e^{2\pi i l x}}{l^b} \sum_{m=1}^K \frac{e^{2\pi i m(x+\gamma)}}{m^{s_3}} \sum_{n=1}^K \frac{e^{2\pi i n(x+\delta)}}{n^{s_4}} dx \\
&= (-1)^{a+b} S_2(a, b, s_3, s_4; 1, 1, \gamma, \delta),
\end{aligned}$$

and

$$\begin{aligned}
& \lim_{K \rightarrow \infty} \int_0^1 \sum_{k=1}^K \frac{e^{2\pi ikx}}{k^a} \sum_{l=1}^K \frac{e^{2\pi ilx}}{l^b} \sum_{m=1}^K \frac{e^{-2\pi im(x-\gamma)}}{m^{s_3}} \sum_{n=1}^K \frac{e^{2\pi in(x+\delta)}}{n^{s_4}} dx \\
& = \zeta_{MT,3}(s_4, a, b, s_3; \delta, 1, 1, \gamma); \\
& \lim_{K \rightarrow \infty} \int_0^1 \sum_{k=-K}^{-1} \frac{e^{2\pi ikx}}{k^a} \sum_{l=1}^K \frac{e^{2\pi ilx}}{l^b} \sum_{m=1}^K \frac{e^{-2\pi im(x-\gamma)}}{m^{s_3}} \sum_{n=1}^K \frac{e^{2\pi in(x+\delta)}}{n^{s_4}} dx \\
& = (-1)^a S_3(a, b, s_3, s_4; 1, 1, \gamma, \delta); \\
& \lim_{K \rightarrow \infty} \int_0^1 \sum_{k=1}^K \frac{e^{2\pi ikx}}{k^a} \sum_{l=-K}^{-1} \frac{e^{2\pi ilx}}{l^b} \sum_{m=1}^K \frac{e^{-2\pi im(x-\gamma)}}{m^{s_3}} \sum_{n=1}^K \frac{e^{2\pi in(x+\delta)}}{n^{s_4}} dx \\
& = (-1)^b S_4(a, b, s_3, s_4; 1, 1, \gamma, \delta); \\
& \lim_{K \rightarrow \infty} \int_0^1 \sum_{k=-K}^{-1} \frac{e^{2\pi ikx}}{k^a} \sum_{l=-K}^{-1} \frac{e^{2\pi ilx}}{l^b} \sum_{m=1}^K \frac{e^{-2\pi im(x-\gamma)}}{m^{s_3}} \sum_{n=1}^K \frac{e^{2\pi in(x+\delta)}}{n^{s_4}} dx \\
& = (-1)^{a+b} \zeta_{MT,3}(a, b, s_3, s_4; 1, 1, \gamma, \delta).
\end{aligned}$$

Hence we see that, for example, for $\Re s_3 > 1$ and $\Re s_4 > 1$,

$$\begin{aligned}
& (-1)^a \zeta_{MT,3}(b, s_3, s_4, a; 1, \gamma, \delta, 1) + (-1)^b \zeta_{MT,3}(s_3, s_4, a, b; \gamma, \delta, 1, 1) \\
& + (-1)^{a+b} S_2(a, b, s_3, s_4; 1, 1, \gamma, \delta) \\
& = \int_0^1 \lim_{K \rightarrow \infty} \sum_{\substack{k=-K \\ k \neq 0}}^K \frac{e^{2\pi ikx}}{k^a} \sum_{\substack{l=-K \\ l \neq 0}}^L \frac{e^{2\pi ilx}}{l^b} \sum_{m=1}^K \frac{e^{2\pi im(x+\gamma)}}{m^{s_3}} \sum_{n=1}^K \frac{e^{2\pi in(x+\delta)}}{n^{s_4}} dx.
\end{aligned}$$

Changing the order of limitation and integration is justified by bounded convergency. Multiplying by $(-1)^{a+b}$ on both sides, and using (3.1)-(3.4), we obtain (3.7), and similarly obtain (3.8). Note that these hold for $\Re s_3 \geq 1$ and $\Re s_4 \geq 1$ by absolute convergency. \square

Next, in the proof of Lemma 3.2, we replace $\sum_{m \geq 1} e^{2\pi im(x+\gamma)} m^{-s}$ and $\sum_{m \geq 1} e^{2\pi im(x+\delta)} m^{-s}$ by $\sum_{m \geq 1} \psi(m) e^{2\pi imx} m^{-s}$ and $\sum_{m \geq 1} \omega(m) e^{2\pi imx} m^{-s}$, respectively. Then we obtain the following.

Lemma 3.3. *For $a, b \in \mathbb{N}$ and $s_3, s_4 \in \mathbb{C}$ with $\Re(s_3) \geq 1$, $\Re(s_4) \geq 1$, we have*

$$\begin{aligned}
(3.9) \quad & (-1)^b L_{MT,3}(b, s_3, s_4, a; \mathbf{1}, \psi, \omega, \mathbf{1}) + (-1)^a L_{MT,3}(s_3, s_4, a, b; \psi, \omega, \mathbf{1}, \mathbf{1}) \\
& + T_2(a, b, s_3, s_4; \mathbf{1}, \mathbf{1}, \psi, \omega) = N_1(a, b, s_3, s_4; \psi, \omega)
\end{aligned}$$

and

$$\begin{aligned}
(3.10) \quad & L_{MT,3}(s_4, a, b, s_3; \omega, \mathbf{1}, \mathbf{1}, \psi) + (-1)^a T_3(a, b, s_3, s_4; \mathbf{1}, \mathbf{1}, \psi, \omega) \\
& + (-1)^b T_4(a, b, s_3, s_4; \mathbf{1}, \mathbf{1}, \psi, \omega) + (-1)^{a+b} L_{MT,3}(a, b, s_3, s_4; \mathbf{1}, \mathbf{1}, \psi, \omega) \\
& = N_2(a, b, s_3, s_4; \psi, \omega).
\end{aligned}$$

Using these results, we can obtain the main theorems.

Theorem 3.4. For $a, b, c \in \mathbb{N}$ and $\delta \in \mathbb{R}$,

$$\begin{aligned}
& \zeta_{MT,3}(a, b, c, s; 1, 1, 1, \delta) - (-1)^{b+c} \zeta_{MT,3}(b, c, s, a; 1, 1, \delta, 1) \\
(3.11) \quad & - (-1)^{c+a} \zeta_{MT,3}(c, s, a, b; 1, \delta, 1, 1) - (-1)^{a+b} \zeta_{MT,3}(s, a, b, c; \delta, 1, 1, 1) \\
& = (-1)^{a+b} M_2(a, b, c, s; 1, \delta) - (-1)^b M_1(a, c, b, s; 1, \delta) \\
& - (-1)^a M_1(c, b, a, s; 1, \delta)
\end{aligned}$$

holds for $s \in \mathbb{C}$ except for singular points of functions on both sides.

Proof. By the definition, we have

$$\begin{aligned}
S_3(a, b, c, s; 1, 1, 1, \delta) &= \sum_{\substack{k, l, m, n=1 \\ k+m=l+n}}^{\infty} \frac{e^{2\pi i n \delta}}{k^a m^c l^b n^s} = S_2(a, c, b, s; 1, 1, 1, \delta), \\
S_4(a, b, c, s; 1, 1, 1, \delta) &= \sum_{\substack{k, l, m, n=1 \\ m+l=k+n}}^{\infty} \frac{e^{2\pi i n \delta}}{m^c l^b k^a n^s} = S_2(c, b, a, s; 1, 1, 1, \delta).
\end{aligned}$$

Therefore we have

$$\begin{aligned}
S_2(a, c, b, s; 1, 1, 1, \delta) &= M_1(a, c, b, s; 1, 1, 1, \delta) \\
&- (-1)^c \zeta_{MT,3}(c, b, s, a; 1, 1, \delta, 1) - (-1)^a \zeta_{MT,3}(b, s, a, c; 1, \delta, 1, 1), \\
S_2(c, b, a, s; 1, 1, 1, \delta) &= M_1(c, b, a, s; 1, 1, 1, \delta) \\
&- (-1)^b \zeta_{MT,3}(b, a, s, c; 1, 1, \delta, 1) - (-1)^c \zeta_{MT,3}(a, s, c, b; 1, \delta, 1, 1).
\end{aligned}$$

Substituting these relations into (3.8), and noting the existence of meromorphic continuation, we obtain the assertion of this theorem. \square

Note that the case $\delta = 1$ coincides with the result in [5, Theorem 4.5]. By the same consideration as in the proof of Theorem 3.4 and using Lemma 3.3, we can obtain the following result which can be regarded as a χ -analogue of [5, Theorem 4.5] and as a triple analogue of the recent results about double zeta and L -functions (see [7, 8, 17, 18]).

Theorem 3.5. For $a, b, c \in \mathbb{N}$,

$$\begin{aligned}
& L_{MT,3}(a, b, c, s; \mathbf{1}, \mathbf{1}, \mathbf{1}, \chi) - (-1)^{b+c} L_{MT,3}(b, c, s, a; \mathbf{1}, \mathbf{1}, \chi, \mathbf{1}) \\
& - (-1)^{c+a} L_{MT,3}(c, s, a, b; \mathbf{1}, \chi, \mathbf{1}, \mathbf{1}) - (-1)^{a+b} L_{MT,3}(s, a, b, c; \chi, \mathbf{1}, \mathbf{1}, \mathbf{1}) \\
& = (-1)^{a+b} N_2(a, b, c, s; \mathbf{1}, \chi) - (-1)^b N_1(a, c, b, s; \mathbf{1}, \chi) \\
& - (-1)^a N_1(c, b, a, s; \mathbf{1}, \chi)
\end{aligned}$$

holds for $s \in \mathbb{C}$ except for singular points of functions on both sides.

Example 3.6. Putting $(a, b, c, s) = (1, 1, 1, 2)$ in Theorem 3.5, we have

$$\begin{aligned}
(3.12) \quad & L_{MT,3}(1, 1, 1, 2; \mathbf{1}, \mathbf{1}, \mathbf{1}, \chi) - 3L_{MT,3}(1, 1, 2, 1; \mathbf{1}, \mathbf{1}, \chi, \mathbf{1}) \\
& = -2L_{MT,2}(2, 2, 1; \chi, \mathbf{1}) - 2L_{MT,2}(2, 1, 2; \mathbf{1}, \mathbf{1}, \chi) \\
& - 4L_{MT,2}(1, 2, 2; \mathbf{1}, \chi, \mathbf{1}) - 2\zeta(2)L(3, \chi).
\end{aligned}$$

4. ALTERNATING TRIPLE ZETA VALUES AND TRIPLE L -VALUES

In this section, we prove new evaluation formulas for alternating triple zeta-functions and for triple L -functions. We first prepare the following lemma which comes from [12, Lemma 8] and is equivalent to [5, Lemma 2.1].

Lemma 4.1. *Let $\phi(s) := \phi(s, 1/2) = \sum_{n \geq 1} (-1)^n n^{-s} = (2^{1-s} - 1) \zeta(s)$. For any function $g : \mathbb{N}_0 \rightarrow \mathbb{C}$, we put*

$$h(j) := \sum_{\mu=0}^{[(j-1)/2]} g(j-2\mu) \frac{(-1)^\mu \pi^{2\mu}}{(2\mu+1)!} \quad (j \in \mathbb{N}).$$

Then we have

$$g(a) = -2 \sum_{\substack{j=1 \\ j \equiv a \pmod{2}}}^a \phi(a-j) h(j) \quad (a \in \mathbb{N}).$$

Proof. We can easily obtain the assertion by putting $P_m = g(2m+1)$ and $Q_m = h(2m+1)$ (resp. $P_m = g(2m+2)$ and $Q_m = h(2m+2)$) for odd (resp. even) a in [12, Lemma 8]. \square

From [5, (3.6) and (3.10)], we have

$$\begin{aligned} (4.1) \quad & \sum_{l,m,n=1}^{\infty} \frac{(-1)^{l+m+n} \cos((l+m+n)\theta)}{l^{2p} m^{2p} n^{2p} (l+m+n)^{2d}} \\ & + \sum_{j=0}^p \phi(2p-2j) \sum_{\nu=0}^{2j} \binom{2d-1+2j-\nu}{2j-\nu} \frac{(-1)^\nu \theta^\nu}{\nu!} \\ & \times \left\{ \sum_{\substack{l,m=1 \\ l \neq m}}^{\infty} \frac{(-1)^{l+m} \cos^{(\nu)}((l-m)\theta)}{l^{2p} m^{2p} (l-m)^{2d+2j-\nu}} - 2 \sum_{l,m=1}^{\infty} \frac{(-1)^{l+m} \cos^{(\nu)}((l+m)\theta)}{l^{2p} m^{2p} (l+m)^{2d+2j-\nu}} \right\} \\ & + \zeta(4p) \sum_{l=1}^{\infty} \frac{(-1)^l \cos(l\theta)}{l^{2p+2d}} = \sum_{\rho=0}^d C_{2d-2\rho}(2p) \frac{(-1)^\rho \theta^{2\rho}}{(2\rho)!} \quad (\theta \in (-\pi, \pi)) \end{aligned}$$

and

$$\begin{aligned} (4.2) \quad & \binom{2e+2p-1}{2p-1} \left\{ \zeta_{MT,2}(2e+2p, 2p, 2p) - \zeta_{MT,2}(2p, 2p, 2e+2p) \right\} \\ & = \sum_{\mu=0}^e C_{2e-2\mu}(2p) \frac{(-1)^\mu \pi^{2\mu}}{(2\mu+1)!} \end{aligned}$$

for $p \in \mathbb{N}$ and $d, e \in \mathbb{N}_0$, where $\{C_{2\nu}(2p) \mid \nu \in \mathbb{N}_0\}$ are constants which can be determined inductively. In fact, applying Lemma 4.1 to (4.2), we have

$$\begin{aligned} (4.3) \quad & C_{2e}(2p) = -2 \sum_{\rho=0}^e \phi(2e-2\rho) \binom{2\rho+2p-1}{2p-1} \\ & \times \{ \zeta_{MT,2}(2\rho+2p, 2p, 2p) - \zeta_{MT,2}(2p, 2p, 2\rho+2p) \}. \end{aligned}$$

On the other hand, letting $\theta = 0$ in (4.1), we have

$$(4.4) \quad C_{2d}(2p) = \zeta_{MT,3}(2p, 2p, 2p, 2d; 1, 1, 1, 1/2) + 2 \sum_{j=0}^p \phi(2p-2j) \binom{2d-1+2j}{2j}$$

$$\begin{aligned} & \times \{\zeta_{MT,2}(2j+2d, 2p, 2p; 1/2, 1, 1) - \zeta_{MT,2}(2p, 2p, 2j+2d; 1, 1, 1/2)\} \\ & + \zeta(4p)\phi(2p+2d) \end{aligned}$$

for $d \in \mathbb{N}_0$. Hence, combining (4.3) and (4.4) in the case $d = e = p$, we obtain the following result which can be regarded as an alternating analogue of [5, Theorem 3.1], and a triple analogue of the previous results about double series [13, 14].

Theorem 4.2. *For $p \in \mathbb{N}$,*

$$\begin{aligned} (4.5) \quad & \zeta_{MT,3}(2p, 2p, 2p, 2p; 1, 1, 1, 1/2) = 2 \sum_{\nu=0}^p \binom{2\nu+2p-1}{2p-1} (2^{1-2p+2\nu} - 1) \\ & \times \zeta(2p-2\nu) \left\{ \zeta_{MT,2}(2p, 2p, 2p+2\nu) - \zeta_{MT,2}(2p+2\nu, 2p, 2p) \right. \\ & \left. + \zeta_{MT,2}(2p, 2p, 2p+2\nu; 1, 1, 1/2) - \zeta_{MT,2}(2p+2\nu, 2p, 2p; 1/2, 1, 1) \right\} \\ & + (1 - 2^{1-4p}) \zeta(4p)^2. \end{aligned}$$

Next we discuss special values of triple L-functions. As well as [5, (3.5)], we can prove that

$$\begin{aligned} (4.6) \quad & \sum_{l,m,n=1}^{\infty} \frac{(-1)^{l+m+n} \sin((l+m+n)\theta)}{lmn(l+m+n)^2} + \theta \sum_{l,m=1}^{\infty} \frac{(-1)^{l+m} \cos((l+m)\theta)}{lm(l+m)^2} \\ & - 2 \sum_{l,m=1}^{\infty} \frac{(-1)^{l+m} \sin((l+m)\theta)}{lm(l+m)^3} + \theta \sum_{l,m=1}^{\infty} \frac{(-1)^m \cos(m\theta)}{lm^2(l+m)} \\ & - 2 \sum_{l,m=1}^{\infty} \frac{(-1)^m \sin(m\theta)}{lm^3(l+m)} = \frac{\zeta(2)^2}{5}\theta + \frac{\zeta(2)}{12}\theta^3 \end{aligned}$$

for $\theta \in (-\pi, \pi)$. For example, we consider the case of $\theta = \pi/2$. For the odd Dirichlet character χ_4 of conductor 4 we know that $\chi_4(k) = \sin(k\pi/2) = -(-1)^k \sin(k\pi/2)$ ($k \in \mathbb{Z}$). Hence, by using $(-1)^k \cos(k\pi/2) = (i^k + i^{-k})/2$ ($k \in \mathbb{Z}$), we can write (4.6) in the case of $\theta = \pi/2$ as

$$\begin{aligned} & -L_{MT,3}(1, 1, 1, 2; \mathbf{1}, \mathbf{1}, \mathbf{1}, \chi_4) + \frac{\pi}{4} \left(\zeta_{MT,2}(1, 1, 2; 1, 1, 1/4) + \zeta_{MT,2}(1, 1, 2; 1, 1, -1/4) \right) \\ & + 2L_{MT,2}(1, 1, 3; \mathbf{1}, \mathbf{1}, \chi_4) + \frac{\pi}{4} \left(\zeta_{MT,2}(1, 2, 1; 1, 1/4, 1) + \zeta_{MT,2}(1, 2, 1; 1, -1/4, 1) \right) \\ & + 2L_{MT,2}(1, 3, 1; \mathbf{1}, \chi_4, \mathbf{1}) = \frac{\pi}{10}\zeta(2)^2 + \frac{1}{96}\zeta(2)\pi^3 = \frac{13}{2880}\pi^5. \end{aligned}$$

Thus we obtain a new evaluation formula

$$\begin{aligned} (4.7) \quad & L_{MT,3}(1, 1, 1, 2; \mathbf{1}, \mathbf{1}, \mathbf{1}, \chi_4) = 2L_{MT,2}(1, 1, 3; \mathbf{1}, \mathbf{1}, \chi_4) + 2L_{MT,2}(1, 3, 1; \mathbf{1}, \chi_4, \mathbf{1}) \\ & + L(1, \chi_4) \left\{ \zeta_{MT,2}(1, 1, 2; 1, 1, 1/4) + \zeta_{MT,2}(1, 1, 2; 1, 1, -1/4) \right. \\ & \left. + \zeta_{MT,2}(1, 2, 1; 1, 1/4, 1) + \zeta_{MT,2}(1, 2, 1; 1, -1/4, 1) \right\} - \frac{104}{75}L(5, \chi_4). \end{aligned}$$

By using the partial fraction decomposition, we can see that $L_{MT,3}(1, 1, 1, 2; \mathbf{1}, \mathbf{1}, \mathbf{1}, \chi_4)$ coincides with $6 \sum_{1 \leq l < m < n} \chi_4(n)/lmn^3$ which is an ordinary triple L -value (see [2]). This

formula (4.7) can be regarded as a χ -analogue of that of $\zeta_{MT,3}$ (see, for example, [5, Example 3.2]), and as a triple analogue of that of double L -values (see [9, 10, 15]). Furthermore, by combining (3.12) and (4.7), we can also give an evaluation formula for $L_{MT,3}(1, 1, 2, 1; \mathbf{1}, \mathbf{1}, \chi_4, \mathbf{1})$ in terms of Dirichlet L -values, double zeta and L -values.

Acknowledgements. The authors wish to express their sincere gratitude to the referee for his (or her) careful reading of the manuscript and helpful advice.

REFERENCES

- [1] T. M. Apostol, *Introduction to Analytic Number Theory*, Springer, 1976.
- [2] T. Arakawa and M. Kaneko: *On multiple L -values*, J. Math. Soc. Japan **56** (2004), 967-992.
- [3] K. Matsumoto, *On the analytic continuation of various multiple zeta-functions*, Number Theory for the Millennium II, Proc. of the Millennial Conference on Number Theory, M. A. Bennett et. al. (eds.), A. K. Peters, (2002), 417-440.
- [4] K. Matsumoto, *On Mordell-Tornheim and other multiple zeta-functions*, Proceedings of the Session in analytic number theory and Diophantine equations (Bonn, January-June 2002), D. R. Heath-Brown and B. Z. Moroz (eds.), Bonner Mathematische Schriften Nr. 360, Bonn 2003, n.25, 17pp.
- [5] K. Matsumoto, T. Nakamura, H. Ochiai and H. Tsumura, *On value-relations, functional relations and singularities of Mordell-Tornheim and related triple zeta-functions*, preprint, submitted for publication.
- [6] L. J. Mordell, *On the evaluation of some multiple series*, J. London Math. Soc. **33** (1958), 368-371.
- [7] T. Nakamura, *A functional relation for the Tornheim double zeta function*, Acta Arith. **125** (2006), 257-263.
- [8] T. Nakamura, *Double Lerch series and their functional relations*, preprint, submitted for publication.
- [9] D. Terhune, *Evaluation of double L -values*, J. Number Theory **105** (2004), 275-301.
- [10] D. Terhune: *Evaluation of a class of double L -values*, Proc. Amer. Math. Soc. **134** (2006), 1881-1889.
- [11] L. Tornheim, *Harmonic double series*, Amer. J. Math. **72** (1950), 303-314.
- [12] H. Tsumura, *On some combinatorial relations for Tornheim's double series*, Acta Arith. **105** (2002), 239-252.
- [13] H. Tsumura, *On alternating analogues of Tornheim's double series*, Proc. Amer. Math. Soc. **131** (2003), 3633-3641.
- [14] H. Tsumura, *Evaluation formulas for Tornheim's type of alternating double series*, Math. Comp. **73** (2004), 251-258.
- [15] H. Tsumura, *On evaluation formulas for double L -values*, Bull. Austral. Math. Soc. **70** (2004), 213-221.
- [16] H. Tsumura, *On Mordell-Tornheim zeta values*, Proc. Amer. Math. Soc. **133** (2005), 2387-2393.
- [17] H. Tsumura, *On some functional relations between the Mordell-Tornheim double L -functions and Dirichlet L -function*, J. Number Theory **120** (2006), 161-178.
- [18] H. Tsumura, *On functional relations between the Mordell-Tornheim double zeta functions and the Riemann zeta function*, to appear in Math. Proc. Cambridge Philos. Soc.
- [19] L. C. Washington, *Introduction to the Cyclotomic Fields*, 2nd ed., Springer-Verlag: New York Berlin Heidelberg, 1997.
- [20] D. Zagier, *Values of zeta functions and their applications*, ECM volume, Progress in Math. **120** (1994), 497-512.

GRADUATE SCHOOL OF MATHEMATICS NAGOYA UNIVERSITY, CHIKUSA-KU, NAGOYA, 464-8602, JAPAN
E-mail address: kohjimat@math.nagoya-u.ac.jp

GRADUATE SCHOOL OF MATHEMATICS NAGOYA UNIVERSITY, CHIKUSA-KU, NAGOYA, 464-8602, JAPAN
E-mail address: m03024z@math.nagoya-u.ac.jp

DEPARTMENT OF MATHEMATICS AND INFORMATION SCIENCES, TOKYO METROPOLITAN UNIVERSITY,
 1-1, MINAMI-OHSAWA, HACHIOJI, TOKYO 192-0397 JAPAN
E-mail address: tsumura@tmu.ac.jp