

Lipopolysaccharide-Deficient Mutants of *Salmonella enterica* Have Increased Sensitivity to Catechins

Miho Yoshii^{1,2}, Akira Okamoto^{2,3*}, Michio Ohta⁴

¹Department of Fundamental Nursing, Graduate School of Medicine and Pharmaceutical Sciences for Research, University of Toyama, Toyama, Japan

²Department of Bacteriology, Nagoya University Graduate School of Medicine, Nagoya, Japan

³Department of School Health Sciences, Aichi University of Education, Kariya, Japan

⁴School of Nursing, Sugiyama Jogakuen University, Nagoya, Japan

Email: *okamoto@auecc.aichi-edu.ac.jp

Received June 20, 2013; revised July 20, 2013; accepted July 31, 2013

Copyright © 2013 Miho Yoshii *et al.* This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

ABSTRACT

Antimicrobial activity is one of the well-known biological characteristics of catechins, the main extract of green tea leaves. It is thought that catechins intercalate into the bacterial cell membrane and damage the lipid bilayer. However, the association between catechins and lipopolysaccharides, which consist of an O side chain, core oligosaccharide, and lipid A, has not been previously investigated. In this study, we evaluated the catechin sensitivity of *Salmonella enterica* mutants that lack the O side chain and have core oligosaccharides of different lengths. These rough mutants were more sensitive to catechins than a bacterial strain with intact lipopolysaccharide. We conclude that the O side chain and core oligosaccharide play an important role in protecting Gram-negative bacteria against the antimicrobial activity of catechins.

Keywords: Antimicrobial Activity; Catechins; Lipopolysaccharide; Gram-Negative Bacteria

1. Introduction

The health benefits of tea have been known for a long time. Over the last quarter-century, several molecules in green tea extract with biological activity have been identified. The main component of green tea extract is catechins, which comprise epicatechin, epigallocatechin, epicatechin gallate, epigallocatechin gallate, and other related compounds. One of the characteristic properties of catechins is antimicrobial activity [1,2]. Although there are several hypotheses about the antimicrobial mechanisms of catechins [3-5], studies with artificial liposomes support the idea that catechins exert their antimicrobial effect mainly by causing damage to the lipid bilayer [6-8].

Generally, Gram-positive bacteria are more sensitive to catechins than Gram-negative bacteria. Ikigai *et al.* proposes that the differences in susceptibility are due to differences in the bacterial cell surface; *i.e.*, the outer membrane of Gram-negative bacteria constitutes a tight barrier to catechins [3]. However, another report indicates that *Vibrio* species are more susceptible to cate-

chins than other Gram-negative species such as *Escherichia coli* or *Salmonella* spp., both of which belong to the family Enterobacteriaceae [2]. While *Vibrio* species also possess an outer membrane with lipopolysaccharide (LPS), the LPS structure is slightly different from that in Enterobacteriaceae [9-13]. These results suggest that the presence or absence of an outer membrane is not the only factor that may affect susceptibility to catechins; the cell surface structures of Gram-negative bacteria should also be considered.

In this study, we investigated the relationship between catechin susceptibility and LPS structure using mutant strains of *Salmonella enterica* that lack O antigen and have core oligosaccharides of different lengths [14,15]. These rough mutant strains had increased susceptibility to catechins, indicating that the O antigen and core oligosaccharide provided some protection against the antimicrobial activity of catechins.

2. Materials and Methods

2.1. Bacterial Strains and Culture Conditions

Rough mutant strains of *Salmonella enterica* serovar

*Corresponding author.

Minnesota (R50, R5, R7, R4, and R595) were used in this study. These strains have been described previously [14-16]. The structure of the O-antigen and R-core oligosaccharide of these strains is shown in **Figure 1**. The wild-type strain LT2 (*S. enterica* serovar Typhimurium), which possesses full-length LPS, was used as a reference.

Bacteria were routinely grown on nutrient agar (Eiken, Tokyo, Japan) and cultured overnight in Heart Infusion Broth (Nissui Pharmaceutical Co., Ltd., Tokyo, Japan) at 37°C.

2.2. Tolerance for Antimicrobial Activity of (-)-Epigallocatechin and (-)-Epigallocatechin Gallate

(-)-Epigallocatechin and (-)-epigallocatechin gallate were purchased from Sigma-Aldrich (St. Louis, MO). Both agents were prepared to give final concentrations of 0.25, 0.5, and 1 mg·mL⁻¹ in PBS with an inoculum density of 1 × 10⁶ to 1 × 10⁷ CFU·mL⁻¹ in a 1 mL volume. Antimicrobial activity of each agent was assessed by time-kill assay. Suspensions were incubated in 96-well plates at 37°C without shaking for 0 to 6 h, and aliquots were removed at 1-h intervals. To obtain colony forming unit counts, dilution series were prepared in PBS (1 × 10⁻¹ to 1 × 10⁻⁹), plated on agar plates (three 100 μL drops per dilution), and incubated at 37°C overnight. For a negative control, PBS was used instead of catechin.

3. Results

The antimicrobial activity of catechins was evaluated by time-kill assay (**Figure 2**). The reference strain LT2 survived 6 h with 0.25 mg·mL⁻¹ of (-)-epigallocatechin or (-)-epigallocatechin gallate (**Figures 2(C) and (F)**). The mutant strains R595, R4, and R7, which possess relatively short core oligosaccharides, were killed within 6 h at all tested catechin concentrations (**Figure 2**). The mutant strain R5, which possesses Rc LPS (*i.e.*, galactosyl-glucose in the core of Rd1 LPS (strain R7)), survived 6 h with 0.25 mg·mL⁻¹ of (-)-epigallocatechin or (-)-epigallocatechin gallate (**Figures 2(C) and (F)**). The mutant strain R50, which possesses Ra LPS (*i.e.*, full-length core oligosaccharide but no O-antigen), survived 6 h with 0.25 mg·mL⁻¹ of (-)-epigallocatechin gallate (**Figure 2(C)**).

4. Discussion

Although the antimicrobial mechanism of catechins remains unclear, there is evidence that their surface activity damages the lipid bilayer [3,6,8]. Moreover, sensitivity to catechins differs among bacterial species and strains; for example, Gram-positive bacteria are more susceptible than Gram-negative bacteria. Ikigai *et al.* proposes that




Figure 1. Structure of the LPS of the *Salmonella enterica* strains used in this study. The LPS phenotypes (Ra, Rc, Rd1, Rd2, and Re) and each mutant strain (R50, R5, R7, R4, and R595) are shown (modified from Osborn (1969)). Glc: glucose; GlcNAc: N-acetyl-glucosamine; Gal: galactose; Hep: heptose; P: phosphate; ETN: ethanolamine; KDO: 2-keto-3-deoxyoctonic acid.

this difference is caused by the repulsion between catechins and the surface of Gram-negative cells, which are coated by LPS [3]. However, among Gram-negative bacteria, *Vibrio* species show higher sensitivity to catechins than Enterobacteriaceae [2]. These reports indicate that sensitivity to catechins is associated with several molecular mechanisms and is not determined only by the presence or absence of LPS on the cell surface. The components of LPS may also affect sensitivity to catechins.

LPS contributes to the mechanical stability of the cell and acts as a barrier to antimicrobial molecules. LPS is a complex glycolipid that consists of three domains: lipid A, core oligosaccharide, and O-polysaccharide. Lipid A is a highly conserved hydrophobic domain that is responsible for the endotoxic properties of many types of LPS. The O side chain, which provides a hydrophilic surface layer, is attached to lipid A through the core oligosaccharide. The O side chain is a polymer of glycosyl residues, and its size, or the number of repeating units, varies among species, strains, or the chemical and physical conditions in which the bacteria are grown. Generally, this size heterogeneity manifests as a ladder pattern in SDS-PAGE.

In this study, the antimicrobial activities of catechins for LPS deficient mutant strains were evaluated by time-kill assay (**Figure 2**). The absence of the O side chain and a shorter core oligosaccharide increased sensitivity to the antimicrobial activity of catechins. Deficiencies in LPS may affect the physical properties of the cell surface, such as rigidity. Another possibility is that the reduction in cell surface hydrophilicity resulting from the absence of the O side chain may allow catechins to more easily access the lipid bilayer. Although other factors, such as multi-drug efflux pumps, may also be involved, the structure of LPS on the cell surface appears to be important for tolerance of the antimicrobial activity of catechins.




Figure 2. Bacterial killing assay for different concentrations of catechins. Antimicrobial activity of (a) $1 \text{ mg}\cdot\text{mL}^{-1}$ (b) $0.5 \text{ mg}\cdot\text{mL}^{-1}$, and (c) $0.25 \text{ mg}\cdot\text{mL}^{-1}$ of epigallocatechin gallate, and (d) $1 \text{ mg}\cdot\text{mL}^{-1}$, (e) $0.5 \text{ mg}\cdot\text{mL}^{-1}$, and (f) $0.25 \text{ mg}\cdot\text{mL}^{-1}$ of epigallocatechin, (g) was compared with control. Symbols were as follows: LT2: cross with dashed line, R50: open circle, R5: filled circle, R7: open triangle, R4: filled triangle, and R595: open square.

5. Acknowledgements

This work was supported by Grant-in-Aid for Young Scientists (KAKENHI) Grant Number 18791632 from Japan Society for the Promotion of Science.

REFERENCES

- [1] N. Caturla, E. Vera-Samper, J. Villalain, C. R. Mateo and V. Micol, "The Relationship between the Antioxidant and the Antibacterial Properties of Galloylated Catechins and the Structure of Phospholipid Model Membranes," *Free*

- Radical Biology & Medicine*, Vol. 34, No. 6, 2003, pp. 648-662. doi:10.1016/S0891-5849(02)01366-7
- [2] T. Taguri, T. Tanaka and I. Kouno, "Antimicrobial Activity of 10 Different Plant Polyphenols against Bacteria Causing Food-Borne Disease," *Biological & Pharmaceutical Bulletin*, Vol. 27, No. 12, 2004, pp. 1965-1969. doi:10.1248/bpb.27.1965
- [3] H. Ikigai, T. Nakae, Y. Hara and T. Shimamura, "Bactericidal Catechins Damage the Lipid Bilayer," *Biochimica et Biophysica Acta*, Vol. 1147, No. 1, 1993, pp. 132-136. doi:10.1016/0005-2736(93)90323-R
- [4] H. Arakawa, M. Maeda, S. Okubo and T. Shimamura, "Role of Hydrogen Peroxide in Bactericidal Action of Catechin," *Biological & Pharmaceutical Bulletin*, Vol. 27, No. 3, 2004, pp. 277-281. doi:10.1248/bpb.27.277
- [5] Y. M. Zhang and C. O. Rock, "Evaluation of Epigallocatechin Gallate and Related Plant Polyphenols as Inhibitors of the FabG and FabI Reductases of Bacterial Type II Fatty-Acid Synthase," *Journal of Biological Chemistry*, Vol. 279, No. 30, 2004, pp. 30994-31001. doi:10.1074/jbc.M403697200
- [6] T. Hashimoto, S. Kumazawa, F. Nanjo, Y. Hara and T. Nakayama, "Interaction of Tea Catechins with Lipid Bilayers Investigated with Liposome Systems," *Bioscience, Biotechnology, and Biochemistry*, Vol. 63, No. 12, 1999, pp. 2252-2255. doi:10.1271/bbb.63.2252
- [7] S. Kumazawa, K. Kajiya, A. Naito, H. Saito, S. Tuzi, M. Tanio, M. Suzuki, F. Nanjo, E. Suzuki and T. Nakayama, "Direct Evidence of Interaction of a Green Tea Polyphenol, Epigallocatechin Gallate, with Lipid Bilayers by Solid-State Nuclear Magnetic Resonance," *Bioscience, Biotechnology, and Biochemistry*, Vol. 68, No. 8, 2004, pp. 1743-1747. doi:10.1271/bbb.68.1743
- [8] Y. Sun, W. C. Hung, F. Y. Chen, C. C. Lee and H. W. Huang, "Interaction of Tea Catechin (-)-Epigallocatechin Gallate with Lipid Bilayers," *Biophysical Journal*, Vol. 96, No. 3, 2009, pp. 1026-1035. doi:10.1016/j.bpj.2008.11.007
- [9] S. Kondo, T. Iguchi and K. Hisatsune, "A Comparative Study of the Sugar Composition of Lipopolysaccharides Isolated from *Vibrio cholerae*, '*Vibrio albensis*' and *Vibrio metschnikovii*," *Journal of General Microbiology*, Vol. 134, No. 6, 1988, pp. 1699-1705.
- [10] C. Whitfield, "Biosynthesis of Lipopolysaccharide O Antigens," *Trends in Microbiology*, Vol. 3, No. 5, 1995, pp. 178-185. doi:10.1016/S0966-842X(00)88917-9
- [11] S. N. Chatterjee and K. Chaudhuri, "Lipopolysaccharides of *Vibrio cholerae*. I. Physical and Chemical Characterization," *Biochimica et Biophysica Acta*, Vol. 1639, No. 2, 2003, pp. 65-79. doi:10.1016/j.bbadis.2003.08.004
- [12] E. Vinogradov, C. Wilde, E. M. Anderson, A. Nakhamchik, J. S. Lam and D. A. Rowe-Magnus, "Structure of the Lipopolysaccharide Core of *Vibrio vulnificus* Type Strain 27562," *Carbohydrate Research*, Vol. 344, No. 4, 2009, pp. 484-490. doi:10.1016/j.carres.2008.12.017
- [13] N. J. Phillips, D. M. Adin, E. V. Stabb, M. J. McFall-Ngai, M. A. Apicella and B. W. Gibson, "The Lipid A from *Vibrio fischeri* Lipopolysaccharide: A Unique Structure Bearing a Phosphoglycerol Moiety," *Journal of Biological Chemistry*, Vol. 286, No. 24, 2011, pp. 21203-21219. doi:10.1074/jbc.M111.239475
- [14] O. Westphal, K. Jann and K. Himmelspach, "Chemistry and Immunochemistry of Bacterial Lipopolysaccharides as Cell Wall Antigens and Endotoxins," *Progress in Allergy*, Vol. 33, 1983, pp. 9-39.
- [15] N. Kato, M. Ohta, N. Kido, H. Ito, S. Naito, Y. Arakawa and T. Komatsu, "Relationship between Mg²⁺-Induced Hexagonal Assembly of R-Form Lipopolysaccharides and Chemical Structure of Their R-Cores," *Microbiology and Immunology*, Vol. 34, No. 3, 1990, pp. 245-257.
- [16] H. Mayer, S. Schlecht and W. Gromska, "Reactivity of Lipopolysaccharides from Various Salmonella SR and R Chemotypes Ra-Re Mutants with Concanavalin A," *Zentralblatt für Bakteriologie, Parasitenkunde, Infektionskrankheiten und Hygiene. Erste Abteilung Originale. Reihe A*, Vol. 233, No. 3, 1975, pp. 327-334.