

Lesson 4


Derivatives of Trigonometric Functions

4A

- Derivative of Sine Function
- Limit of $\frac{\sin x}{x}$
- Derivatives of Basic Trigonometric Function

Derivative of Sine Function

Variation of slopes



Derivative by definition

$$\begin{aligned} f'(x) &= \lim_{h \rightarrow 0} \frac{f(x+h) - f(x)}{h} = \lim_{h \rightarrow 0} \frac{\sin(x+h) - \sin x}{h} \\ &= \lim_{h \rightarrow 0} \frac{\sin x \cos h + \cos x \sin h - \sin x}{h} \\ &= \lim_{h \rightarrow 0} \left(\cos x \frac{\sin h}{h} - \sin x \frac{1 - \cos h}{h} \right) \end{aligned}$$

Limit of $\sin x / x$

Consider a sector with central angle x

Compare the areas of $\triangle OAB$, sector OAB, and $\triangle OAT$

$$\frac{1}{2} \cdot 1 \cdot \sin x < (\pi \cdot 1^2) \cdot \frac{x}{2\pi} < \frac{1}{2} \cdot 1 \cdot \tan x$$

$$\therefore \sin x < x < \tan x$$



Divide by $\sin x (> 0)$

$$\therefore 1 < \frac{x}{\sin x} < \frac{1}{\cos x}$$

$$\therefore 1 > \frac{\sin x}{x} > \cos x$$

As $x \rightarrow 0$

$$\lim_{h \rightarrow 0} \frac{\sin x}{x} = 1$$



Derivative of Sine Function—Cont.

$$f'(x) = \lim_{h \rightarrow 0} \left(\cos x \frac{\sin h}{h} - \sin x \frac{1 - \cos h}{h} \right)$$
$$\frac{(1 - \cos h)(1 + \cos h)}{h(1 + \cos h)} = \frac{1 - \cos^2 h}{h(1 + \cos h)} = \frac{\sin h}{h} \frac{\sin h}{(1 + \cos h)}$$

Therefore

$$(\sin x)' = \cos x$$



That makes sense!

Example

[Example 4-1] Derive the derivative of $\cos x$ and $\tan x$.


Ans.

(1) From the triangle in the right side

$$\cos x = \sin\left(\frac{\pi}{2} - x\right)$$

Therefore

$$\begin{aligned}(\cos x)' &= \left(\sin\left(\frac{\pi}{2} - x\right)\right)' = \cos\left(\frac{\pi}{2} - x\right) \cdot \left(\frac{\pi}{2} - x\right)' \\ &= \sin x \times (-1) = -\sin x\end{aligned}$$



(2) From the quotient rule

$$\begin{aligned}(\tan x)' &= \left(\frac{\sin x}{\cos x}\right)' = \frac{\cos x \cdot \cos x - \sin x(-\sin x)}{\cos^2 x} = \frac{\cos^2 x + \sin^2 x}{\cos^2 x} \\ &= \frac{1}{\cos^2 x} = \sec^2 x\end{aligned}$$

Summary of Derivatives of Tri. Functions

(1) The basic trigonometric derivatives (**Memorize!**)

$$\frac{d}{dx} \sin x = \cos x, \quad \frac{d}{dx} \cos x = -\sin x$$

(2) Other standard relationships

(Derive from (1) if necessary)

$$\frac{d}{dx} \tan x = \sec^2 x, \quad \frac{d}{dx} \sec x = \sec x \tan x$$

$$\frac{d}{dx} \cot x = -\csc^2 x, \quad \frac{d}{dx} \csc x = -\csc x \cot x$$

[note]

These formula are valid only when the angle x is **measured in radians**.

Example

[Example 4.2] Find the derivatives of the following functions:

$$(1) \quad y = \cos^2 x$$

$$(2) \quad y = x \sin x + \cos x$$

Ans.

(1) Chain rule

$$\begin{aligned} y' &= 2 \cos x \cdot (\cos x)' = 2 \cos x \cdot (-\sin x) \\ &= -2 \sin x \cos x = -\sin 2x \end{aligned}$$

(2) Product rule

$$\begin{aligned} y' &= (x \cdot \sin x)' + (\cos x)' \\ &= (1 \cdot \sin x + x \cos x) - \sin x = x \cos x \end{aligned}$$

Exercise

[Ex.4.1] Find the derivatives of the following functions:

(1) $y = \sin ax^2$

(2) $y = \frac{1}{\tan x}$

(3) $y = \cos\left(\frac{x}{2} + \frac{\pi}{6}\right)$

Pause the video and solve the problem by yourself.

Answer to Exercise

[Ex.4.1] Find the derivatives of the following functions:

$$(1) \ y = \sin ax^2 \quad (2) \ y = \frac{1}{\tan x} \quad (3) \ y = \cos\left(\frac{x}{2} + \frac{\pi}{6}\right)$$

$$(1) \ \frac{d}{dx} \sin(ax^2) = \frac{d}{du} \sin u \frac{d}{dx} (ax^2) = \cos u \cdot (2ax) = 2ax \cos(ax^2)$$

$$(2) \ y' = \left(\frac{\cos x}{\sin x}\right)' = \frac{(-\sin x) \sin x - \cos x(\cos x)}{\sin^2 x} = -\frac{(\sin^2 x + \cos^2 x)}{\sin^2 x} = -\frac{1}{\sin^2 x}$$

$$(3) \ y' = \frac{d}{du} \cos u \frac{d}{dx} \left(\frac{x}{2} + \frac{\pi}{6}\right) = -\frac{1}{2} \sin\left(\frac{x}{2} + \frac{\pi}{6}\right)$$

Lesson 4


Derivatives of Trigonometric Functions

4B

- Derivatives and Motions
- Position, Velocity and Acceleration
- Simple Harmonic Motion

Velocity and Acceleration

Point **P** is moving on the straight line :



Its position is given by
 $x = f(t)$

The **average velocity** between t_1 and t_2


$$\bar{v} = \frac{\Delta x}{\Delta t} = \frac{x_2 - x_1}{t_2 - t_1}$$

The **instantaneous velocity** at $t = t_1$

$$v(t_1) = \lim_{\Delta t \rightarrow 0} \frac{\Delta x}{\Delta t} \quad v = \frac{ds}{dt} = f'(t)$$

The **instantaneous acceleration** at $t = t_1$

$$\alpha = \frac{dv}{dt} = f''(t)$$



Example

[**Example 4-3**] The position of the mass moving on the x -axis is given by $s(t) = t^3 - 3t^2 - 9t + 10$


- (1) Find the velocity and the acceleration at $t = 2$.
- (2) Investigate the motion during $-2 \leq t \leq 4$

Ans. (1) Velocity : $v = \frac{ds}{dt} = 3t^2 - 6t - 9 = 3(t+1)(t-3) \quad \therefore v(2) = -9$


Acceleration : $a = \frac{dv}{dt} = 6(t-1) \quad \therefore a(2) = 6$

(2)

t	-2	-1	3	4
v		+	0	-	0	+	
s	8	\nearrow	15	\searrow	-17	\nearrow	-10



Simple Harmonic Motion



Vertical position : $y = A \sin(\omega t + \alpha)$

Velocity : $v = \frac{dy}{dt} = A \omega \cos(\omega t + \alpha)$

Acceleration : $a = \frac{dv}{dt} = \frac{d^2y}{dt^2} = -A \omega^2 \sin(\omega t + \alpha)$

Exercise

[Exercise.4.2] Point P is moving on the x-axis. Its position is given by

$x = 2t + \cos t$. Find the time when the point has the maximum velocity and its maximum velocity.

Pause the video and solve the problem by yourself.

Answer to the Exercise

[Exercise.4.2] Point P is moving on the x-axis. Its position is given by

$$x = 2t + \cos t \quad . \text{ Find the time when the point has the maximum velocity}$$

and its maximum velocity.

Ans.

$$\text{Velocity} \quad v = \frac{dx}{dt} = 2 - \sin t$$

Maximum velocity occurs at

$$\sin t = -1$$

Therefore

$$t = \frac{3}{2}\pi + 2n\pi$$

Maximum velocity is 3.

