


Synthesis of Free-base 10-Azacorroles

Hiroto Omori, Satoru Hiroto,* and Hiroshi Shinokubo*

Department of Applied Chemistry, Graduate School of Engineering, Nagoya University, Aichi, 464-8603, Japan

Supporting Information Placeholder



ABSTRACT: A novel synthetic protocol has been developed for free-base 10-azacorroles. Ni(0)-mediated homo-coupling of nitrogen-bridged bisdipyrrin Zn(II) complexes afforded a series of free-base 5,15-diaryl-10-azacorroles in good yields. Pd(II) and Cu(II) complexes have been prepared through metallation of free-base 10-azacorroles. Optical and electrochemical properties of 10-azacorrole metal complexes can be tuned by *meso*-aryl substituents and central metals. Cyclic voltammetry and theoretical calculations elucidated that the central metals of 10-azacorroles significantly affected their electronic properties.

Introduction of metals into the central cavity of porphyrins has been useful method for fine-tuning of their optical and electrochemical properties. In addition, porphyrin metal complexes have been applied to wide-range of chemical researches in such as supramolecular chemistry, photosynthesis, catalysts, and so forth.^{1,2} Such metal complexes have been usually prepared from their corresponding free-base porphyrins by treatment with various metal sources. Thus, the effective synthesis of free-base porphyrin derivatives is a key issue to expand their chemistry by metal coordination.


Corroles are a porphyrin analogue with one direct pyrrole-pyrrole linkage.^{3,4} Recently, several research groups have developed the synthesis of heterocorroles, namely, corroles containing heteroatoms on their aromatic conjugation circuit.⁵ In particular, installation of heteroatoms at *meso*-positions dramatically changes its electronic structure. For example, D. P. Goldberg and coworkers have developed the chemistry of corrolazine, which has three nitrogen atoms at *meso*-positions.^{6,7} They accomplished the synthesis of free-base corrolazine through removal of phosphorus(V) from a corrolazine cavity under reductive conditions.⁸ Corrolazine stabilizes high oxidation states of the central metals, enabling isolation of reactive intermediate in mimicking of biological oxidation system. Bröring and coworkers reported the synthesis of free-base β -alkyl type 10-oxa-, 10-thia-, and 10-selenacorroles by reduction of their corresponding copper(II) complexes.^{9,10}

Our group independently developed the synthetic protocols for *meso*-aryl-substituted 10-aza-, 10-oxa-, and 10-thiacorroles.¹¹ However, the central metals of these *meso*-aryl-10-heterocorroles have been limited to Ni(II) and Al(III) complexes and preparation of free-base 10-heterocorroles has remained unsuccessful. Removal of Ni(II) and Al(III) proved to

be difficult by known demetallation procedures such as treatment with H₂SO₄ or Grignard reagents.¹²

Here, we disclose a novel protocol to prepare free-base 10-azacorroles from nitrogen bridged bisdipyrrin Zn(II) complexes. This protocol efficiently provides a series of *meso*-aryl substituted 10-azacorroles. In addition, divalent metal complexes such as Zn(II), Cu(II), and Pd(II) have been successfully synthesized.

Scheme 1. Preparation of Zn(II) 10-azacorroles and their demetallation.



We prepared nitrogen-bridged bisdipyrrin Zn(II) complex **1a** according to the previous report. Cyclization of **1a** was accomplished under Yamamoto homo-coupling conditions with Ni(cod)₂, 1,5-cyclooctadiene, and 2,2'-bipyridyl to produce Zn(II) 10-azacorrole **2a** in 83% yield (Scheme 1).¹³ The formation of 10-azacorrole skeleton was confirmed by the ¹H NMR spectrum of **2a**, which exhibited four doublet peaks of β-protons in the aromatic region. Finally, the structure of **2a** was unambiguously elucidated by X-ray diffraction analysis. Figure 1a and 1b show crystal structures of **2a**, in which one of pyrrole rings is tilted. In the crystal, the central Zn(II) metal was ligated by methanol, forming pentacoordinate structure. The mean plane deviation was 0.088 Å, which is larger than the corresponding Ni(II) complex (0.020 Å). Such displacement is due to pentacoordinated conformation and the larger size of Zn(II) metal, which is located out of the mean plane.

Zn(II) metal in **2a** was easily removed by treatment with trifluoroacetic acid (TFA), producing the corresponding free-base 10-azacorrole **3a** in almost quantitative yield. The ¹H NMR spectrum of **3a** exhibited symmetric feature and inner NH protons of **3a** appeared at 5.23 ppm. The larger downfield shift of the NH proton as compared with free-base porphyrins would reflect the lower aromaticity of **3a**. This trend was also observed in other 10-heterocorroles.^{10a} The highly planar structure of **3a** was revealed by X-ray diffraction analysis (Figure 1c and 1d). The position of inner NH protons could not be determined. However, the planar structure of **3a** indicates the existence of two NH protons in the central cavity because inner three NH protons should induce distortion as is observed in corroles. This was also supported by the ¹H NMR spectrum of **3a**, showing two integrated peak at 5.2 ppm, which could be assigned as inner NH protons.




Figure 1. X-ray crystal structures of Zn(II)- and free-base 10-azacorroles. (a) Top view and (b) side view of **2a**. (c) Top view and (d) side view of **3a**. Mesityl substituents are omitted for clarity. The thermal ellipsoids are scaled at 50% probability level.





Figure 2. X-ray crystal structure of **5a**. (a) Top view and (b) side view. *meso*-Mesityl groups are omitted for clarity. The thermal ellipsoids are scaled at 50% probability level.

On the basis of the present cyclization procedure, we prepared 10-azacorroles with various *meso*-aryl substituents. In the case of *meso*-phenyl and 3,5-dimethoxyphenyl groups, demetallation of Zn(II) metal complexes **2b** and **2c** proceeded during silica-gel purification, to furnish free-base azacorroles **3b** and **3c** directly in 89% and 81% yield, respectively.

Scheme 2. Metallation of bisdipyrrins **2a** and **2b** with Cu(II), and Pd(II) ions.



With free-base **3a** in hand, we introduced various metals into its central cavity. Treatment of **3a** with Cu(OAc)₂•H₂O gave Cu(II) complex **4a** in 92% yield. Pd(II) complex **5a** was obtained in 63% yield in the reaction with Pd(OAc)₂ (Scheme 2). The insertion of these metals was confirmed by high-resolution mass spectral analysis. The parent mass ion peaks of **4a** and **5a** was observed at *m/z* = 687.2403 (calcd for (C₄₃H₃₈N₅Cu)⁺ = 687.2418) and *m/z* = 730.2160 (calcd for (C₄₃H₃₈N₅Pd)⁺ = 730.2172). The structure of **5a** was elucidated by X-ray diffraction analysis, showing its highly planar conformation. The mean plane deviation of **5a** was 0.024 Å. The N–Pd bond lengths were 1.92 and 1.93 Å. These values are shorter than those of typical Pd(II) porphyrins (2.03 Å), reflecting smaller cavity of 10-azacorroles than that of porphyrins.¹⁴

Figure 3a shows UV/vis absorption and emission spectra of **3a**, **3b**, and **3c** in CH₂Cl₂. Compared with **3a**, **3b** and **3c** exhibited red-shifted Soret and Q bands, indicating the presence of electronic interaction between the azacorrole core and the *meso*-aryl group except in the case of **3a**. This is due to more flexible conformation of other substituents than the mesityl group in **3a**. This tendency was also confirmed by electrochemical analysis. Cyclic voltammetry of **3a**, **3b**, and **3c** were performed in THF (Figure S21). The result was summarized in Table 1. Both the first oxidation potential and the first reduction potential of **3b** were decreased in comparison to **3a**, resulting in the small HOMO–LUMO gap. This result is in good agreement with the result of optical analysis. The fluorescent quantum yields of these azacorroles were almost identical (0.12 for **3a**, 0.088 for **3b**, and 0.10 for **3c**), indicating that the *meso*-aryl group does not significantly affect the excited state dynamics.




Figure 3. UV/vis absorption (solid line) and emission (dotted line) spectra of (a) **3a** (black), **3b** (red) and **3c** (blue) and (b) **2a** (blue), **3a** (black), **4a** (red), and **5a** (green) in CH_2Cl_2 .

Figure 3b illustrates UV/vis absorption and emission spectra of **2a**, **3a**, and **5a** in CH_2Cl_2 . Zinc(II) complex **2a** exhibited a red-shifted but similar shaped spectrum as compared with **3a**. Zinc(II) complex **2a** and free-base **3a** showed red emission. The fluorescence quantum yields of **2a** and **3a** were 0.11 and 0.026, respectively. These features are similar to those of porphyrins. On the other hand, **5a** exhibited broad and split Soret-band as well as bathochromically shifted Q bands. Table 1 shows the result of electrochemical measurement of azacorroles **2a**, **3a** and **5a** (Figure S22). As is observed in the case of porphyrins, Zn(II) metallation lowered the first oxidation potential in comparison to the free-base.

Table 1. Summary of electrochemical data.^{a,b}

compound	E_{red}^2	E_{red}^1	E_{ox}^1	E_{ox}^2	ΔE
3a	-2.40	-2.01	0.357	0.697	2.37
3b	—	-1.87	0.327	0.689	2.20
3c	-2.23	-1.87	0.347	0.687	2.22
2a	—	-2.19	0.077	0.412	2.27
5a	-2.39	-1.94	0.360	0.716	2.30

a: unit: V, *b*: potentials are shown versus ferrocene/ferrocenium couple, *c*: determined by differential pulse voltammetry, *d*: $\Delta E = E_{\text{ox}}^1 - E_{\text{red}}^1$.

The different absorption feature between **2a** and **5a** was further investigated by theoretical calculations. The absorption

spectra of **2a** and **5a** were simulated by the time-dependent (TD) DFT method at the CAM-B3LYP/6-31G+LanL2DZ level (Figure S24). According to the calculations, two intense bands of **5a** at 393 and 447 nm could be assigned as transitions from HOMO-1 to LUMO and HOMO-3 to LUMO. On the other hand, the sharp Soret band of **2a** at 431 nm could be attributed to transitions from HOMO-1 to LUMO and from HOMO to LUMO+1. As reported previously, the low symmetry of 10-azacorroles resulted in the contribution of the HOMO-3 to LUMO transition into Soret and Q-bands. In fact, HOMO-3 of **5a** contains significant contribution of d orbitals on the central Pd metal to π -orbitals of azacorrole skeleton. (Figure S23). Consequently, the transitions from these orbitals to LUMO become allowed to broaden the spectrum. This feature is not observed for normal porphyrins. Although the divalent nature and emission properties of 10-azacorroles are almost identical with those of porphyrin analogues, the different effect of central metals to the absorption properties highlights the specific feature of 10-azacorroles.

In summary, we have achieved efficient synthesis of free-base 10-azacorroles with various *meso*-aryl substituents. This is the first synthesis of free-base *meso*-aryl substituted 10-heterocorroles. Pd(II) and Cu(II) metal ions were inserted into the central cavity of 10-azacorroles. The structures of free-base, Zn(II), and Pd(II) 10-azacorroles were clearly elucidated by X-ray diffraction analysis. Cyclic voltammetry and theoretical calculations revealed the specific effect of central metals on optical properties of 10-azacorroles. These results would expand the utility of 10-azacorroles in various applications. Further investigations on optical properties dependent on substituents are under way in our group.

ASSOCIATED CONTENT

Supporting Information

Experimental details, compound data, CIF files, and copies of NMR spectra. This material is available free of charge via the Internet at <http://pubs.acs.org>.

AUTHOR INFORMATION

Corresponding Author

*hshino@apchem.nagoya-u.ac.jp; hiroto@apchem.nagoya-u.ac.jp

ACKNOWLEDGMENT

This work was supported by a Grant-in-Aid for Scientific Research on Innovative Areas “New Polymeric Materials based on Element Block” (15H00731) and “ π -System Figuration” (26102003), and Program for Leading Graduate Schools “Integrative Graduate Education and Research in Green Natural Sciences” from MEXT Japan. H.S. is grateful to The Asahi Glass Foundation for financial support.

REFERENCES

- (1) (a) *The Porphyrin Handbook*; Kadish, K.; Smith, K. M.; Guillard, R. Eds.; Academic Press, Boston, **2003**, vol. 11–20. (b) *Handbook of Porphyrin Science*; Kadish, K.; Smith, K. M.; Guillard, R. Eds.; World Scientific Publishing Co. Pte. Ltd. Singapore, **2010**, vol. 1–30.
- (2) (a) Durot, S.; Taesch, J.; Heitz, V. *Chem. Rev.* **2014**, *114*, 8542. (b) Li, L.-L.; Diau, E. W.-G. *Chem. Soc. Rev.* **2013**, *42*, 291. (c) Beletskaya, I.; Tyurin, V. S.; Tsivadze, A. Y.; Guillard, R.; Stern, C. *Chem. Rev.* **2009**, *109*, 1659. (d) Guldi, D. M.; Rahman, G. M. A.; Sgobba, V. Ehli, C. *Chem. Soc. Rev.* **2006**, *35*, 471. (e) Meunier, B. *Chem. Rev.* **1992**, *92*, 1411. (f) Kondratuk, D. V.; Perdigo, L. M. A.; Esmail, A. M. S.; O’Shea, J. N.; Beton,

- P. H.; Anderson, H. L. *Nat. Chem.* **2015**, *7*, 317. (g) Costentin, C.; Drouet, S.; Robert, M.; Savéant, J.-M. *Science*, **2012**, *338*, 90. (h) O'Sullivan, M. C.; Sprafke, J. K.; Kondratuk, D. V.; Rinfrey, C.; Claridge, T. D. W.; Saywell, A.; Blunt, M. O.; O'Shea, J. N.; Beton, P. H.; Malfois, M.; Anderson, H. L. *Nature* **2011**, *469*, 72. (b) Kosal, M. E.; Chou, J. H.; Wilson, S. R.; Suslick, K. S. *Nat. Mater.* **2002**, *1*, 118.
- (3) (a) Paolesse, R. in *The Porphyrin Handbook*; Kadish, K. M., Smith, K. M., Guillard, R., Eds.; Academic Press, San Diego, **2000**, vol. 2, pp201-232. (b) Guillard, R. in *The Porphyrin Handbook*; Kadish, K. M., Smith, K. M., Guillard, R., Eds.; Elsevier, Amsterdam, **2003**, vol 18, pp. 303-351.
- (4) (a) Thomas, K. E.; Alemayehu, A. B.; Conradie, J.; Beavers, C. M.; Gosh, A. *Acc. Chem. Res.* **2012**, *45*, 1203. (b) Lemon, C. M.; Brothers, P. J. *J. Porphyrins Phthalocyanines*, **2011**, *15*, 809. (c) Aviv-Harel, I.; Gross, Z. *Chem. Eur. J.* **2009**, *15*, 8382. (d) Flamigni, L.; Gryko, D. T. *Chem. Soc. Rev.* **2009**, *38*, 1635. (e) Paolesse, R. *Synlett*, **2008**, *15*, 2215.
- (5) (a) Ghosh, A.; Chatterjee, T.; Lee, W.-Z.; Ravikanth, M. *Org. Lett.* **2013**, *15*, 1040. (b) Shetti, V. S.; Prabhu, U. R.; Ravilanth, M. *J. Org. Chem.* **2010**, *75*, 4172. (c) Gryko, D. P.; Fox, J. P.; Goldberg, D. P. *J. Porphyrins Phthalocyanines*, **2004**, *8*, 1091. (d) Sankar, J.; Rath, H.; PrabhuRaja, V.; Chandrashekar, T. K.; Vittal, J. J. *J. Org. Chem.* **2004**, *69*, 5135. (e) Pawlicki, M.; Latos-Grazynski, L.; Szterenber, L. *J. Org. Chem.* **2002**, *67*, 5644. (f) Narayanan, S. J.; Sridevi, B.; Chandrashekar, T. K.; Englich, U.; Ruhlandt-Senge, K. *Org. Lett.* **1999**, *1*, 587.
- (6) McGown, A. J.; Badiei, Y. M.; Leeladee, P.; Prokop, K. A.; De-Beer, S.; Goldberg, D. P. In *Handbook of Porphyrin Science*; Kadish, K. M., Smith, K. M., Guillard, R., Eds.; World Scientific Press: Singapore, **2011**; vol. 14, pp 525-599.
- (7) (a) Neu, H. M.; Baglia, R. A.; Goldberg, D. P. *Acc. Chem. Res.* **2015**, *48*, 2754. (b) Goldberg, D. P. *Acc. Chem. Res.* **2007**, *40*, 626.
- (8) Ramdhanie, B.; Stern, C. L.; Goldberg, D. P. *J. Am. Chem. Soc.* **2001**, *123*, 9447.
- (9) (a) Broadhurst, M. J.; Grigg, R.; Johnson, A. W., *J. Chem. Soc. Perkin Trans. 1* **1972**, 1124. (b) Johnson, W.; Kay, I. T.; Rodrigo, R., *J. Chem. Soc.* **1963**, 2336.
- (10) (a) Sakow, D.; Baabe, D.; Böker, B.; Burghaus, O.; Funk, M.; Kleeberg, C.; Menzel, D.; Pietzonka, C.; Bröring, M. *Chem. Eur. J.* **2014**, *20*, 2913. (b) Sakow, D.; Böker, B.; Brandhorst, K.; Burghaus, O.; Bröring, M., *Angew. Chem. Int. Ed.* **2013**, *52*, 4912. (c) Bröring, M.; Köhler, S.; Pietzonka, C., *J. Porphyrins Phthalocyanines* **2012**, *16*, 641. (d) Bröring, M.; Brégier, F.; Tejero, E. C.; Hell, C.; Holthausen, M. C., *Angew. Chem. Int. Ed.* **2007**, *46*, 445.
- (11) (a) Omori, H.; Hiroto, S.; Shinokubo, H. *Chem. Commun.* **2016**, *52*, 3540. (b) Horie, M.; Hayashi, Y.; Yamaguchi, S.; Shinokubo, H., *Chem. Eur. J.* **2012**, *18*, 5919. (c) Kamiya, H.; Kondo, T.; Sakida, T.; Yamaguchi, S.; Shinokubo, H., *Chem. Eur. J.* **2012**, *18*, 16129. (d) Ito, T.; Hayashi, Y.; Shimizu, S.; Shin, J.-Y.; Kobayashi, N.; Shinokubo, H., *Angew. Chem. Int. Ed.* **2012**, *51*, 8542.
- (12) Murakami, K.; Yamamoto, Y.; Yorimitsu, H.; Osuka, A. *Chem. Eur. J.* **2013**, *19*, 9123.
- (13) Kende, A. S.; Liebeskind, L. S.; Braitsch, D. M. *Tetrahedron Lett.* **1975**, *16*, 3375.
- (14) Sun, D.; Tham, F. S.; Reed, C. A.; Chaker, L.; Burgess, M.; Boyd, P. D. W. *J. Am. Chem. Soc.* **2000**, *122*, 10704.