

Cleaving Dihydrogen with Tetra(*o*-tolyl)diborane(4)

Nana Tsukahara,[†] Hiroki Asakawa,[†] Ka-Ho Lee,[‡] Zhenyang Lin,^{*‡} Makoto Yamashita^{*§¶}

[†]Department of Applied Chemistry, Faculty of Science and Engineering, Chuo University, 1-13-27, Kasuga, Bunkyo-ku, 112-8551, Tokyo, Japan.

[‡]Department of Chemistry, The Hong Kong University of Science and Technology, Clear Water Bay, Kowloon, Hong Kong

[§]Department of Applied Chemistry, Graduate School of Engineering, Nagoya University, Furo-cho, Chikusa-ku, Nagoya, Aichi 464-8603, Japan

[¶]Research Development Initiative, Chuo University, 1-13-27 Kasuga, Bunkyo-ku, Tokyo 112-8551, Japan

Supporting Information Placeholder

ABSTRACT: Tetra(*o*-tolyl)diborane(4) **1** was synthesized and characterized experimentally as well as theoretically by DFT calculations. Exposure of **1** to H₂ (1 bar) at room temperature afforded the corresponding di(*o*-tolyl)hydroborane through cleavage of the H–H and B–B bonds. DFT calculations suggested a diarylborylanion character for the transition state.

Since the discovery of heterogeneously catalyzed hydrogenations by Sabatier,¹ catalytic and stoichiometric hydrogenations with H₂ have represented major research fields in molecular transition-metal chemistry.² However, the limited availability of precious metals, which are active as hydrogenation catalysts, has fueled the quest for more sustainable options. Two such recently developed alternatives are: i) the replacement of the precious metals with base metals, and ii) the use of catalysts based on main-group elements.³ The latter can be subdivided into two classes: i) frustrated Lewis pairs (FLPs),⁴ which can cleave H₂ heterolytically, and ii) main-group element compounds in low oxidation states. The latter class involves group 13 or 14 elements such as C(II), Si(II), Ge(I), Sn(I), P(II), B(o), B(II), Al(I), and Ga(I).⁵ Our recent report regarding the deprotonation of H₂ with boryllithium⁶ is also relevant in this context, as boryl anions can be considered as a B(I) species. As far as reactive trivalent group 13 compounds are concerned, the reaction of an antiaromatic borole with H₂,⁷ the reversible cleavage of H₂ with dianionic diboranthracenes,⁸ and FLP-like reactivity of diaminogallium with cooperation of ligand⁹ have been recently reported.

Recently, we reported the synthesis of the unsymmetrical diborane(4) pinB–BMes₂¹⁰ and its reactivity toward CO and *tert*-butylisocyanide.^{10a} This diborane(4) reacts also with alkynes to form diborylalkenes^{10b} and the regioselectivity of the diboration to afford *syn*- or *anti*-diborylalkenes can be controlled by addition of a catalytic amount of base and 1,2-dimethoxyethane. This diborane(4) furthermore reacts with 2,6-dimethylphenylisocyanide to furnish an 1,2-oxaboretane ring through a ring-contraction of Bpin.^{10c} This characteristic reactivity of pinB–BMes₂ was attributed to the high electron affinity arising from the two parallelly arranged vacant p-orbitals on the two boron atoms.^{10d} Similarly high electron affinity was reported for Mes₂B–B(Mes)Ph, which was synthesized from (MeO)₂B–B(OMe)₂ and is capable to form the corresponding radical anion and dianion.¹¹ Moreover, Mes₂B–B(Mes)Ph represents the only reported example of a tetraaryldiborane(4). Herein, we report the synthesis of tetra(*o*-tolyl)diborane(4) **1** and its reactivity toward H₂ at room temperature, as well as a mechanistic study based on DFT calculations.

By careful choice of the precursor, **1** could be obtained in 30% isolated yield from the reaction between B₂cat₂ and *o*-tolMgBr (Scheme 1). The molecular structure of **1** was determined by single-crystal X-ray diffraction analysis (Figure 1). In the crystal, **1** adopts an almost orthogonally twisted structure [C8–B1–B2–C15 = 101.9(7)°], similar to that of Mes₂B–B(Mes)Ph.^{11a} The ¹H NMR spectrum of **1** exhibited four magnetically equivalent *o*-tol groups at room temperature, indicating free rotation around the B–B and B–C bonds on the NMR timescale. The ¹¹B NMR spectrum of **1** in C₆D₆ showed a broad signal at 89 ppm. Dissolution of **1** in THF did not alter the ¹¹B

NMR chemical shift, indicating that THF does not coordinate to the boron atom in **1**.

Scheme 1. Synthesis of tetra(*o*-tolyl)diborane(**4**) **1**.




Figure 1. Molecular structure of **1** (thermal ellipsoids set at 50% probability; one of two independent molecules and hydrogen atoms were omitted for clarity).

To estimate the electron affinity of **1**, electrochemical measurements and DFT calculations were conducted. The cyclic voltammogram of a THF solution of **1** showed a reversible reduction wave at -2.1 V (vs. $\text{Cp}_2\text{Fe}/\text{Cp}_2\text{Fe}^+$). The first reduction potential of **1** is thus less negative than that of previously reported pinB-BMes₂ (-2.5 V vs. $\text{Cp}_2\text{Fe}/\text{Cp}_2\text{Fe}^+$). Similar to our previous report,^{10d} we also calculated the dependency of the LUMO level and the free energy of **1** on the torsion angle C(*ipso*)-B-B-C(*ipso*). The optimized structure of **1** contains a torsion angle of -96.6° , which is comparable to the crystal structure and leads to a calculated LUMO level of -1.29 eV. Increasing the torsion angle of **1** from -90° to 0° in increments of 10° lowers the LUMO energy (as well as the stability) as shown in Figure 2. When the torsion angle approaches 0° , the LUMO energy drops to -2.20 eV and the corresponding structure is destabilized by 8.51 kcal/mol. This relatively mild destabilization suggests that rotation around the B-B bond should occur at room temperature. The lowering of the LUMO of **1** upon rotation around the B-B bond should be attributed to the overlap of the two vacant p-orbitals on the two boron centers (Figure 3). The LUMO of **1** in the ground state (-96.6°) consists of π^* -orbitals of two *o*-tol rings and one vacant p-orbital of the boron atom. The LUMO+1 is of similar energy and mainly localized on the other B(*o*-tol)₂ moiety, while the LUMO of **1** (0.0°) exhibits two completely merged vacant p-orbitals on the two boron atoms. It should be noted that the HOMOs of

both rotational isomers exhibit a B-B σ -bond character.




Figure 2. Dependency of the LUMO energy level (in eV) and the relative stability (in kcal/mol) of **1** on the torsion angle of the C(*ipso*)-B-B-C(*ipso*) moiety.




Figure 3. Frontier orbitals of the rotational isomers of **1** at C(*ipso*)-B-B-C(*ipso*) = 0.0° (top) and C(*ipso*)-B-B-C(*ipso*) = -96.6° (bottom).

Tetra(*o*-tolyl)diborane(**4**) **1** reacts directly with H₂. Exposing a hexane solution of **1** for 2 h to H₂ (1 bar) at room temperature afforded di(*o*-tolyl)hydroborane **2** in 52% yield (Scheme 2). The crystal structure of **2** revealed a dimeric structure with hydride bridges (Figure 4, left), which is similar to those of previously reported dimeric diarylhydroboranes.¹² The solid state IR spectrum of **2** exhibited a characteristic μ -H vibration at 1520 cm⁻¹, which is comparable to that of [HB(C₆F₅)₂]₂.^{12b} Dissolution of **2** in C₆D₆ or CDCl₃ led to five B-bonded ¹H nuclei and three ¹¹B nuclei in the ¹H{¹¹B} and ¹¹B NMR spectra, respectively. This behavior should be attributed to a monomer/dimer equilibrium¹³ and a potential coordination of solvent molecules to the highly Lewis acidic boron center of the monomer.^{12b,14} In contrast, dissolution of **2** in THF led to a simple NMR spectra indicating a formation of **2**·THF. Addition of pyridine to isolated **2** without solvent at room temperature afforded single crystals of the corresponding pyridine-hydroborane complex **3**, which was completely characterized by

NMR spectroscopy, X-ray analysis, and high-resolution FAB mass spectroscopy.

Scheme 2. Reaction of **1** with H₂





Figure 4. Molecular structures of **2** (left) and **3** (right) (thermal ellipsoids at 50% probability; hydrogen atoms except for B–H were omitted for clarity).

To get a better understanding of the detailed pathway for the reaction of **1** with H₂, DFT calculations were performed (Scheme 3). The results suggested that the overall reaction should consist of four steps. The hydrogenation of the B–B bond in **1** should be initiated by coordination of the H–H bond to one of the vacant p-orbitals of one of the boron atom to afford intermediate **4**, in which the C(*ipso*)–B–B–C(*ipso*) torsion angle changed to -55.1° , indicating that the strong electron affinity of **1** enables the binding to H₂. Upon coordination of H₂, both the H–H and B–B bonds should elongate and the coordination geometry of the H₂-coordinated boron atom should become pyramidalized. Subsequently, cleavage of the B–B σ -bond should be accompanied by a proton-migrating H–H bond cleavage via TS₄₋₅, leading to the formation of the dimeric hydroborane intermediate **5**, which contains a hydride and an *o*-tolyl group as bridging ligands. Here, it should be noted that in TS₄₋₅, the *ipso* carbon atom of one *o*-tolyl group starts to interact with the boron atom it is not directly attached to [C(*ipso*)–B 2.347 Å]. TS₄₋₅ represents the rate-determining step of the reaction sequence and requires an overall activation free energy of 19.8 kcal/mol. The resulting intermediate (**5**) is thus located 8.3 kcal/mol below the reference point (**1** + H₂). Intermediate **5** involves a characteristic B–C–B three-center-two-electron bond, which has recently been confirmed by Wagner.¹⁵ Dissociation of **5** to afford monomeric hydroborane **6**, and the subsequent dimerization of **6** to afford hydride-bridged dimer **2** through the two low-energy transition states TS₅₋₆ and TS₆₋₂ include relatively low activation bar-

riers. The overall reaction from **1** + H₂ to give **5** is exergonic by 20.1 kcal/mol.

Scheme 3. Proposed reaction mechanism and energy profile for the hydrogenolysis of **1** [Ar = *o*-tolyl]; calculated at Mo6/6-31G(d) level of theory with solvent effect of benzene(SMD); dashed lines correspond to the formation/cleavage of bonds or 3-center-2-electron bonds].^a



^aRelative free energies and electronic energies (in parentheses) are given in kcal/mol.




Figure 5. Selected natural bond orbitals: (a) NBO₉₁ and (b) NBO₉₂ of TS₄₋₅, as well as (c) schematic illustration of the diarylborylanion character of B₃ in TS₄₋₅ (numbering of atoms is based on the calculations)

To characterize the transition state for the H–H bond cleavage, an NBO analysis was performed. The calculated natural charge on the migrating hydrogen atom (H₁: +0.306) is more positive than that on the other hydrogen atom (H₂: +0.200), indicating that H₁ should be transferred as a proton. The NBO analysis on TS₄₋₅ showed that the vacant orbital of the migrating proton H₁ interacts with both the bonding orbitals of the H–B and B–B bonds (NBO 9₁ and NBO₉₂ shown in Figure 5a,b), and exhibits significant second-order perturbation energies (*E*₂) of 352.52 and 200.49 kcal/mol, respectively. Thus, the transition states cleaving H–H bond (TS₄₋₅) may be described as an intramolecular deprotonation by an "sp²-sp³ diborane" that exhibits a diarylborylanion

character (Figure 5c). The proton-migrating character of TS_{4-5} is similar to that of the deprotonation of H_2 with boryllithium,⁶ and to that for the cleavage of H–H by an antiaromatic borole involving a B–C bond cleavage.⁷

In conclusion, tetra(o-tolyl)diborane(4) **1** was synthesized, and characterized experimentally as well as theoretically by DFT calculations. Exposure of **1** to H_2 at room temperature afforded the corresponding diarylhydroborane via cleavage of the H–H and B–B bonds. DFT calculations suggested a diarylborylanion character for the transition state.

ASSOCIATED CONTENT

Supporting Information

The Supporting Information is available free of charge on the ACS Publications website at DOI: 10.1021/jacs.x.

- Experimental and computational details (PDF)
- Crystallographic data (CIF)
- DFT Coordinates (XYZ)

AUTHOR INFORMATION

Corresponding Authors

*makoto@oec.apchem.nagoya-u.ac.jp, *chzlin@ust.hk

Notes

The authors declare no competing financial interests.

ACKNOWLEDGMENT

This research was supported by JSPS KAKENHI Grant Number JP24109012 and 26288019, the Research Grants Council of Hong Kong (HKUST16303614 and C5023-14G), CREST from the JST (14529307), and The Asahi Glass Foundation. The authors thank Profs. T. Hiyama (Chuo U) and Y. Nishibayashi (U-Tokyo) for providing us with access to an X-ray diffractometer and to a mass spectrometer. This paper is dedicated to Professor Takayuki Kawashima on the occasion of his 70th birthday.

REFERENCES

- (1) Sabatier, P., *Ind. Eng. Chem.* **1926**, *18*, 1005-1008.
- (2) (a) *The handbook of homogeneous hydrogenation*. Wiley-VCH: 2007; (b) Hartwig, J. F., Chapter 7, Oxidative addition. In *Organotransition Metal Chemistry: From Bonding to Catalysis*, University Science Books: Sausalito, CA, 2010.
- (3) (a) Power, P. P., *Nature* **2010**, *463*, 171-177; (b) Power, P. P., *Acc. Chem. Res.* **2011**, *44*, 627-637; (c) Martin, D.; Soleilhavoup, M.; Bertrand, G., *Chem. Sci.* **2011**, *2*, 389-399.
- (4) (a) Stephan, D. W., *Chem. Commun.* **2010**, *46*, 8526-8533; (b) Welch, G. C.; Juan, R. R. S.; Masuda, J. D.; Stephan, D. W., *Science* **2006**, *314*, 1124-1126; (c) Chase, P. A.; Welch, G. C.; Jurca, T.; Stephan, D. W., *Angew. Chem. Int. Ed.* **2007**, *46*, 8050-8053; (d) Stephan, D. W.; Erker, G., *Angew. Chem. Int. Ed.* **2010**, *49*, 46-76; (e) Stephan, D. W., *Org. Biomol. Chem.* **2012**, *10*, 5740-5746; (f) Hounjet, L. J.; Stephan, D. W., *Org. Process Res. Dev.* **2014**, *18*, 385-391; (g) Stephan, D. W., *Acc. Chem. Res.* **2015**, *48*, 306-316; (h) Stephan, D. W., *J. Am. Chem. Soc.* **2015**, *137*, 10018-10032; (i) Stephan, D. W.; Erker, G., *Angew. Chem. Int. Ed.* **2015**, *54*, 6400-

6441; (j) Weicker, S. A.; Stephan, D. W., *Bull. Chem. Soc. Jpn.* **2015**, *88*, 1003-1016.

(5) (a) Spikes, G. H.; Fettinger, J. C.; Power, P. P., *J. Am. Chem. Soc.* **2005**, *127*, 12232-12233; (b) Frey, G. D.; Lavallo, V.; Donnadiou, B.; Schoeller, W. W.; Bertrand, G., *Science* **2007**, *316*, 439-441; (c) Peng, Y.; Brynda, M.; Ellis, B. D.; Fettinger, J. C.; Rivard, E.; Power, P. P., *Chem. Commun.* **2008**, 6042-6044; (d) Zhu, Z.; Wang, X.; Peng, Y.; Lei, H.; Fettinger, J. C.; Rivard, E.; Power, P. P., *Angew. Chem. Int. Ed.* **2009**, *48*, 2031-2034; (e) Seifert, A.; Scheid, D.; Linti, G.; Zessin, T., *Chem. Eur. J.* **2009**, *15*, 12114-12120; (f) Shoji, Y.; Matsuo, T.; Hashizume, D.; Fueno, H.; Tanaka, K.; Tamao, K., *J. Am. Chem. Soc.* **2010**, *132*, 8258-8260; (g) Li, J.; Schenk, C.; Goedecke, C.; Frenking, G.; Jones, C., *J. Am. Chem. Soc.* **2011**, *133*, 18622-18625; (h) Protchenko, A. V.; Birj Kumar, K. H.; Dange, D.; Schwarz, A. D.; Vidovic, D.; Jones, C.; Kaltsoyannis, N.; Mountford, P.; Aldridge, S., *J. Am. Chem. Soc.* **2012**, *134*, 6500-6503; (i) Schäfer, A.; Reißmann, M.; Schäfer, A.; Schmidtman, M.; Müller, T., *Chem. Eur. J.* **2014**, *20*, 9381-9386; (j) Hadlington, T. J.; Jones, C., *Chem. Commun.* **2014**, 50, 2321-2323; (k) Vasko, P.; Wang, S.; Tuononen, H. M.; Power, P. P., *Angew. Chem. Int. Ed.* **2015**, *54*, 3802-3805; (l) Arrowsmith, M.; Böhnke, J.; Braunschweig, H.; Celik, M. A.; Dellermann, T.; Hammond, K., *Chem. Eur. J.* **2016**, *22*, 17169-17172; (m) Nagata, K.; Murotsuki, T.; Agou, T.; Sasamori, T.; Matsuo, T.; Tokitoh, N., *Angew. Chem. Int. Ed.* **2016**, *55*, 12877-12880; (n) Hinz, A.; Schulz, A.; Villinger, A., *Angew. Chem. Int. Ed.* **2016**, *55*, 12214-12218.

(6) Dettenrieder, N.; Aramaki, Y.; Wolf, B. M.; Maichle-Mössner, C.; Zhao, X.; Yamashita, M.; Nozaki, K.; Anwander, R., *Angew. Chem. Int. Ed.* **2014**, *53*, 6259-6262.

(7) (a) Fan, C.; Mercier, L. G.; Piers, W. E.; Tuononen, H. M.; Parvez, M., *J. Am. Chem. Soc.* **2010**, *132*, 9604-9606; (b) Houghton, A. Y.; Karttunen, V. A.; Fan, C.; Piers, W. E.; Tuononen, H. M., *J. Am. Chem. Soc.* **2013**, *135*, 941-947.

(8) von Grotthuss, E.; Diefenbach, M.; Bolte, M.; Lerner, H.-W.; Holthausen, M. C.; Wagner, M., *Angew. Chem. Int. Ed.* **2016**, *55*, 14067-14071.

(9) Abdalla, J. A. B.; Riddlestone, I. M.; Tirfoin, R.; Aldridge, S., *Angew. Chem. Int. Ed.* **2015**, *54*, 5098-5102.

(10) (a) Asakawa, H.; Lee, K.-H.; Lin, Z.; Yamashita, M., *Nat. Commun.* **2014**, *5*, 4245; (b) Kojima, C.; Lee, K.-H.; Lin, Z.; Yamashita, M., *J. Am. Chem. Soc.* **2016**, *138*, 6662-6669; (c) Katsuma, Y.; Asakawa, H.; Lee, K.-H.; Lin, Z.; Yamashita, M., *Organometallics* **2016**, *35*, 2563-2566; (d) Asakawa, H.; Lee, K.-H.; Furukawa, K.; Lin, Z.; Yamashita, M., *Chem. Eur. J.* **2015**, *21*, 4267-4271.

(11) (a) Moezzi, A.; Olmstead, M. M.; Bartlett, R. A.; Power, P. P., *Organometallics* **1992**, *11*, 2383-2388; (b) Moezzi, A.; Olmstead, M. M.; Power, P. P., *J. Am. Chem. Soc.* **1992**, *114*, 2715-2717; (c) Power, P. P., *Inorg. Chim. Acta* **1992**, *198-200*, 443-447.

(12) (a) Entwistle, C. D.; Marder, T. B.; Smith, P. S.; Howard, J. A. K.; Fox, M. A.; Mason, S. A., *J. Organomet. Chem.* **2003**, *680*, 165-172; (b) Parks, D. J.; Piers, W. E.; Yap, G. P. A., *Organometallics* **1998**, *17*, 5492-5503; (c) Hübner, A.; Qu, Z.-W.; Englert, U.; Bolte, M.; Lerner, H.-W.; Holthausen, M. C.; Wagner, M., *J. Am. Chem. Soc.* **2011**, *133*, 4596-4609; (d) Seven, Ö.; Qu, Z.-W.; Zhu, H.; Bolte, M.; Lerner, H.-W.; Holthausen, M. C.; Wagner, M., *Chem. Eur. J.* **2012**, *18*, 11284-11295.

(13) The possible involvement of trimer and tetramer in the equilibrium was also examined by DFT calculations. However, stable structures could not be obtained for these compounds. For details, see Supporting Information.

(14) The coordination of $\text{B}(\text{C}_6\text{F}_5)_3$ to toluene in the gas phase was calculated to be thermoneutral; see: Timoshkin, A. Y.; Frenking, G., *Organometallics* **2008**, *27*, 371-380.

(15) Hübner, A.; Diefenbach, M.; Bolte, M.; Lerner, H.-W.; Holthausen, M. C.; Wagner, M., *Angew. Chem. Int. Ed.* **2012**, *51*, 12514-12518.

1 Authors are required to submit a graphic entry for the Table of Contents (TOC) that, in conjunction with the man-
2 uscript title, should give the reader a representative idea of one of the following: A key structure, reaction, equation,
3 concept, or theorem, etc., that is discussed in the manuscript. Consult the journal's Instructions for Authors for
4 TOC graphic specifications.
5

