

Anti-DFS70 antibodies in Japanese children with dermatomyositis, localized scleroderma and idiopathic arthritis with iridocyclitis

Journal:	<i>The Journal of Rheumatology</i>
Manuscript ID	2016-1409.R1
Manuscript Type:	Correspondence
Date Submitted by the Author:	n/a
Complete List of Authors:	Muro, Yoshinao; Nagoya University Graduate School of Medicine, Dermatology Iwata, Naomi; Aichi Children's Health and Medical Center, Infection and Immunology Tanaka, Yoshihito; Japan Community Health Care Organization Chukyo Hospital, Dermatology Kodera, Masanari; Japan Community Health Care Organization Chukyo Hospital, Dermatology Kono, Michihiro; Nagoya University Graduate School of Medicine, Dermatology Akiyama, Masashi; Nagoya University Graduate School of Medicine, Dermatology
Keywords:	anti-DFS70 antibody, juvenile dermatomyositis, localized scleroderma, anti-nuclear antibody

Correspondence

**Anti-DFS70 antibodies in Japanese children with dermatomyositis,
localized scleroderma and idiopathic arthritis with iridocyclitis**

YOSHINAO MURO, NAOMI IWATA, YOSHIHITO TANAKA,
MASANARI KODERA, MICHIIHIRO KONO, MASASHI AKIYAMA

Key indexing terms: anti-DFS70 antibody, anti-nuclear antibody, juvenile
dermatomyositis, juvenile idiopathic arthritis, localized scleroderma

Yoshinao Muro, MD, PhD: Department of Dermatology, Nagoya University Graduate
School of Medicine, Showa-ku, Nagoya, Japan

Naomi Iwata, MD: Department of Infection and Immunology, Aichi Children's Health
and Medical Center, Obu, Aichi, Japan

Yoshihito Tanaka, MD: Department of Dermatology, Japan Community Health Care
Organization Chukyo Hospital, Nagoya, Japan

Masanari Kodera, MD, PhD: Department of Dermatology, Japan Community Health
Care Organization Chukyo Hospital, Nagoya, Japan

Michihiro Kono, MD, PhD: Department of Dermatology, Nagoya University Graduate
School of Medicine, Showa-ku, Nagoya, Japan

Masashi Akiyama, MD, PhD: Department of Dermatology, Nagoya University
Graduate School of Medicine, Showa-ku, Nagoya, Japan

Correspondence and reprint requests to: Yoshinao Muro, MD, PhD

Division of Connective Tissue Disease and Autoimmunity, Department of Dermatology,
Nagoya University Graduate School of Medicine, 65 Tsurumai-cho, Showa-ku,
Nagoya 466-8550, JAPAN

TEL: 81-52-744-2314

FAX: 81-52-744-2318

E-mail: ymuro@med.nagoya-u.ac.jp

Short running title: Anti-DFS70 in JDM and LS

To the Editor:

We read with interest the article by Schmeling, *et al.*¹ They screened a large number of sera from children with antinuclear antibody (ANA)-associated rheumatic disease (AARD) and related conditions, as well as from a reference cohort, for anti-dense fine speckled 70 autoantibodies (anti-DFS70ab). They showed that the frequency of anti-DFS70ab in healthy children (3/145, 2.1%) was within the lower range of that reported in adults, but that anti-DFS70ab were found at significantly higher frequencies in children with localized scleroderma (LS; 4/29, 13.8%), juvenile dermatomyositis (JDM; 2/11, 18.2%), and uveitis (3/26, 11.5%). For 331 patients with systemic lupus erythematosus (SLE) and 41 patients with systemic scleroderma, they investigated concomitant AARD-associated autoantibodies and found that only 6 patients with SLE had isolated anti-DFS70ab. This result fits the theory that anti-DFS70ab positivity in AARD is usually associated with additional AARD-related antibodies². However, since their disease cohort was small, except for SLE and juvenile idiopathic arthritis (JIA) (202 patients), the higher frequency of the antibodies in LS (4/29), JDM (2/11) or uveitis (3/26) is inconclusive. In addition, neither several recently identified DM-specific autoantibodies (anti-MDA5, anti-TIF, anti-NXP-2, etc.),³ nor anti-ssDNA antibodies that are associated with some patients with LS⁴ were

examined in their study. We examined anti-DFS70ab and myopathy-associated autoantibodies in Japanese children with JDM and investigated anti-DFS70ab and anti-ssDNA antibodies in Japanese children with LS. Moreover, we investigated anti-DFS70ab in JIA patients complicated with iridocyclitis.

Our child cohort consisted of 29 patients with JDM, 14 with LS and 10 with JIA. The age range for children was set at 18 years and younger, in line with previous publications. This study was approved by the Ethics Committee of each hospital, and it complies with the Declaration of Helsinki guidelines. ANA testing was performed by indirect immunofluorescence (IIF) on HEp-2 cell substrates (MBL, Nagoya, Japan)⁵, and anti-DFS70ab levels were measured by ELISA (MBL). JDM-specific autoantibodies including anti-Mi-2, anti-TIF1 γ , anti-MDA5, anti-NXP2, anti-SAE, anti-MLH1, and anti-PMS1 antibodies were measured for JDM patients by the ELISA we developed, which uses *in vitro* transcription and translation biotinylated recombinant protein⁶⁻⁸. The anti-aminoacyl-tRNA synthetase antibodies anti-Jo-1, anti-PL-7, anti-PL-12, anti-EJ, and anti-KS were also investigated by ELISA (MBL) for JDM patients. Anti-ssDNA antibody was measured by ELISA (MBL) for LS patients.

The anti-DFS70ab ELISA found 7 (24.1%) positive patients in our JDM cohort

(Table 1). Two patients with JDM showed the DFS pattern in IIF studies but were anti-DFS70ab negative in the ELISA. Immunoblotting analysis with bacterially expressed DFS70 recombinant protein and HeLa cell extract⁵ (data not shown) confirmed that one of the two was positive for anti-DFS70ab (Table 2, case #8) and the other was negative. One patient with anti-DFS70ab did not show the characteristic DFS pattern in IIF. This patient (Table 2, case #5) was confirmed to be positive for the antibody by the same immunoblotting analysis (discussed below).

The frequency of anti-DFS70ab in JDM was similar to that of the previous study by Schmeling *et al.*¹ (27.6% vs. 18.2%). The most important result is that there was only 1 (3.4%) isolated (monospecific) anti-DFS70ab-positive JDM patient in the present study (Table 2). This datum fits our previous data indicating that the monospecificity of anti-DFS70ab differs between AARD and non-AARD in adults.² Although the Schmeling study showed a relatively high frequency of anti-DFS70ab in patients with LS (13.8%), none of our patients with LS had that antibody. However, there was no statistical difference ($p < 0.287$ by Fisher's exact test). Interestingly, our LS patients often had anti-ssDNA antibodies (9/14, 64.3%).

Schmeling *et al.*¹ studied anti-DFS70ab also in patients with uveitis that was associated with JIA or that was idiopathic. Two of 19 (10.5%) JIA patients with uveitis

and 1 of 7 (14.3%) idiopathic uveitis patients had anti-DFS70ab. According to the study by Ravelli *et al.*,⁹ JIA patients with iridocyclitis very frequently had ANA (58/60, 96.7%). Surprisingly, in our present study, 5 of 10 JIA patients with iridocyclitis had anti-DFS70ab. The association of anti-DFS70ab with JIA with iridocyclitis in Japanese children should be confirmed by future studies. Although 1 of these 5 anti-DFS70ab-positive sera did not show a DFS pattern in IIF, the antibody-positive result was confirmed by immunoblotting analysis (Table 2, case #12). In a total of 53 examined sera, 4 sera had discrepant results between the DFS pattern in IIF and the anti-DFS70ab in ELISA. Schmeling *et al.*¹ also showed that, although the concordance rate of IIF patterns and chemiluminescence immunoassay results for anti-DFS70ab was >90%, IIF patterns do not always correspond to the other immunoassays results for anti-DFS70ab. Since recognizing the DFS pattern in IIF is not an easy task,¹⁰ a specific immunoassay for anti-DFS70ab should be used to confirm their presence. Moreover, clinicians should not over-interpret isolated ANA results in anti-DFS70ab-positive patients, but should pay attention to detecting concomitant disease-specific autoantibodies.

REFERENCES

1. Schmeling H, Mahler M, Levy DM, Moore K, Stevens AM, Wick J, et al. Autoantibodies to dense fine speckles in pediatric diseases and controls. *J Rheumatol* 2015;42:2419-26.
2. Muro Y, Sugiura K, Morita Y, Tomita Y. High concomitance of disease marker autoantibodies in anti-DFS70/LEDGF autoantibody-positive patients with autoimmune rheumatic disease. *Lupus*. 2008;17:171-6.
3. Deakin CT, Yasin SA, Simou S, Arnold KA, Tansley SL, Betteridge ZE, et al. Muscle biopsy findings in combination with myositis-specific autoantibodies aid prediction of outcomes in juvenile dermatomyositis. *Arthritis Rheumatol* 2016;68:2806-16
4. Arkachaisri T, Fertig N, Pino S, Medsger TA Jr. Serum autoantibodies and their clinical associations in patients with childhood- and adult-onset linear scleroderma. A single-center study. *J Rheumatol* 2008;35:2439-44.
5. Watanabe A, Kodera M, Sugiura K, Ususa T, Tan EM, Takasaki Y, et al. Anti-DFS70 antibodies in 597 healthy hospital workers. *Arthritis Rheum* 2005;32:2144-9.
6. Muro Y, Sugiura K, Nara M, Sakamoto I, Suzuki N, Akiyama M. High incidence

- of cancer in anti-small ubiquitin-like modifier activating enzyme antibody-positive dermatomyositis. *Rheumatology (Oxford)* 2015;54:1745-7.
7. Muro Y, Sugiura K, Akiyama M. A new ELISA for dermatomyositis autoantibodies: rapid introduction of autoantigen cDNA to recombinant assays for autoantibody measurement. *Clin Dev Immunol* 2013;2013:856815.
 8. Muro Y, Nakashima R, Hosono Y, Sugiura K, Mimori T, Akiyama M. Autoantibodies to DNA mismatch repair enzymes in polymyositis/dermatomyositis and other autoimmune diseases: a possible marker of favorable prognosis. *Arthritis Rheumatol* 2014;66:3457-62.
 9. Ravelli A, Felici E, Magni-Manzoni S, Pistorio A, Novarini C, Bozzola E, et al. Patients with antinuclear antibody-positive juvenile idiopathic arthritis constitute a homogeneous subgroup irrespective of the course of joint disease. *Arthritis Rheum* 2005;52:826-32.
 10. Bentow C, Fritzler MJ, Mummert E, Mahler M. Recognition of the dense fine speckled (DFS) pattern remains challenging: results from an international internet-based survey. *Autoimmun Highlights* 2016;7:8

Table 1. Clinical profiles and frequency of anti-DFS70 antibodies in the present cohorts.

Cohort	n	Sex M:F	Age (yrs) Range (median)	Anti-nuclear antibody*			Anti-DFS70-positive, n (%)	Other autoantibody -positive, n (%)
				Positive, n (%)	Range (median)	DFS-pattern positive, n (%)		
Juvenile DM	29	11:18	1 – 16 (9)	23 (79.3)	1/40 – 1/320 (x80)	8 (27.6)	8 (27.6)	anti-TIF1 γ , 9 (31.0) anti-MDA5, 8 (27.6) anti-PMS1, 6 (20.7) anti-NXP2, 5 (17.2) anti-MLH1, 3 (10.3)
Localized scleroderma	14	2:12	3 – 18 (12)	13 (92.9)	1/40 – 1/640 (x80)	0 (0)	0 (0)	anti-ssDNA, 9 (64.3)
JIA with iridocyclitis	10	4:6	1 – 11 (6)	10 (100)	1/40 – 1/640 (x160)	4 (40)	5 (50)	NT

*"Anti-nuclear antibody" also includes anti-cytoplasmic antibody.

DFS: dense fine speckled; DM: dermatomyositis; JIA: juvenile idiopathic arthritis; M:F: male:female; NT: not tested

Table 2. Profiles of patients with anti-DFS70 antibodies.

Patient No.	Diagnosis	Age, yrs	Sex	DFS70 reactivity by ELISA, unit/ml*	ANA pattern/ titer by IIF	Other autoantibodies
1	JDM	6	female	22.5	DFS 1/80	Negative
2	JDM	10	female	111.9	DFS 1/160	anti-TIF1 γ
3	JDM	1	male	51.3	DFS 1/80	anti-TIF1 γ
4	JDM	9	male	121.7	DFS 1/320	anti-NXP2, anti-MLH1, anti-PMS1
5	JDM	6	male	85.4	Speckled 1/160	anti-NXP2, anti-MLH1, anti-PMS1
6	JDM	9	female	40.5	DFS 1/80	anti-MDA5
7	JDM	11	female	33.3	DFS 1/80	anti-MDA5
8	JDM	11	female	12.8	DFS 1/80	anti-MDA5
9	JIA iridocyclitis	2	female	252.5	DFS 1/320	NT
10	JIA iridocyclitis	7	female	68.8	DFS 1/160	NT
11	JIA iridocyclitis	6	female	62.0	DFS 1/160	NT
12	JIA iridocyclitis	5	male	52.9	Nucleolar 1/80	NT
13	JIA iridocyclitis	5	male	23.2	DFS 1/80	NT

*positive >15 unit/ml. Sera from all patients were confirmed to have anti-DFS70 antibodies by immunoblotting analysis with DFS70 recombinant protein and with HeLa cell extract.

DFS: dense fine speckled; JDM; juvenile dermatomyositis; JIA: juvenile idiopathic arthritis; NT: not tested.

Correspondence

**Anti-DFS70 antibodies in Japanese children with dermatomyositis,
localized scleroderma and idiopathic arthritis with iridocyclitis**

YOSHINAO MURO, NAOMI IWATA, YOSHIHITO TANAKA,
MASANARI KODERA, MICHIIHIRO KONO, MASASHI AKIYAMA

Key indexing terms: anti-DFS70 antibody, anti-nuclear antibody, juvenile
dermatomyositis, juvenile idiopathic arthritis, localized scleroderma

Yoshinao Muro, MD, PhD: Department of Dermatology, Nagoya University Graduate
School of Medicine, Showa-ku, Nagoya, Japan

Naomi Iwata, MD: Department of Infection and Immunology, Aichi Children's Health
and Medical Center, Obu, Aichi, Japan

Yoshihito Tanaka, MD: Department of Dermatology, Japan Community Health Care
Organization Chukyo Hospital, Nagoya, Japan

Masanari Kodera, MD, PhD: Department of Dermatology, Japan Community Health
Care Organization Chukyo Hospital, Nagoya, Japan

Michihiro Kono, MD, PhD: Department of Dermatology, Nagoya University Graduate
School of Medicine, Showa-ku, Nagoya, Japan

Masashi Akiyama, MD, PhD: Department of Dermatology, Nagoya University
Graduate School of Medicine, Showa-ku, Nagoya, Japan

Correspondence and reprint requests to: Yoshinao Muro, MD, PhD

Division of Connective Tissue Disease and Autoimmunity, Department of Dermatology,
Nagoya University Graduate School of Medicine, 65 Tsurumai-cho, Showa-ku,
Nagoya 466-8550, JAPAN

TEL: 81-52-744-2314

FAX: 81-52-744-2318

E-mail: ymuro@med.nagoya-u.ac.jp

Short running title: Anti-DFS70 in JDM and LS

To the Editor:

We read with interest the article by Schmeling, *et al.*¹ They screened a large number of sera from children with antinuclear antibody (ANA)-associated rheumatic disease (AARD) and related conditions, as well as from a reference cohort, for anti-dense fine speckled 70 autoantibodies (anti-DFS70ab). They showed that the frequency of anti-DFS70ab in healthy children (3/145, 2.1%) was within the lower range of that reported in adults, but that anti-DFS70ab were found at significantly higher frequencies in children with localized scleroderma (LS; 4/29, 13.8%), juvenile dermatomyositis (JDM; 2/11, 18.2%), and uveitis (3/26, 11.5%). For 331 patients with systemic lupus erythematosus (SLE) and 41 patients with systemic scleroderma, they investigated concomitant AARD-associated autoantibodies and found that only 6 patients with SLE had isolated anti-DFS70ab. This result fits the theory that anti-DFS70ab positivity in AARD is usually associated with additional AARD-related antibodies². However, since their disease cohort was small, except for SLE and juvenile idiopathic arthritis (JIA) (202 patients), the higher frequency of the antibodies in LS (4/29), JDM (2/11) or uveitis (3/26) is inconclusive. In addition, neither several recently identified DM-specific autoantibodies (anti-MDA5, anti-TIF, anti-NXP-2, etc.),³ nor anti-ssDNA antibodies that are associated with some patients with LS⁴ were

examined in their study. We examined anti-DFS70ab and myopathy-associated autoantibodies in Japanese children with JDM and investigated anti-DFS70ab and anti-ssDNA antibodies in Japanese children with LS. Moreover, we investigated anti-DFS70ab in JIA patients complicated with iridocyclitis.

Our child cohort consisted of 29 patients with JDM, 14 with LS and 10 with JIA. The age range for children was set at 18 years and younger, in line with previous publications. This study was approved by the Ethics Committee of each hospital, and it complies with the Declaration of Helsinki guidelines. ANA testing was performed by indirect immunofluorescence (IIF) on HEp-2 cell substrates (MBL, Nagoya, Japan)⁵, and anti-DFS70ab levels were measured by ELISA (MBL). JDM-specific autoantibodies including anti-Mi-2, anti-TIF1 γ , anti-MDA5, anti-NXP2, anti-SAE, anti-MLH1, and anti-PMS1 antibodies were measured for JDM patients by the ELISA we developed, which uses *in vitro* transcription and translation biotinylated recombinant protein⁶⁻⁸. The anti-aminoacyl-tRNA synthetase antibodies anti-Jo-1, anti-PL-7, anti-PL-12, anti-EJ, and anti-KS were also investigated by ELISA (MBL) for JDM patients. Anti-ssDNA antibody was measured by ELISA (MBL) for LS patients.

The anti-DFS70ab ELISA found 7 (24.1%) positive patients in our JDM cohort

(Table 1). Two patients with JDM showed the DFS pattern in IIF studies but were anti-DFS70ab negative in the ELISA. Immunoblotting analysis with bacterially expressed DFS70 recombinant protein and HeLa cell extract⁵ (data not shown) confirmed that one of the two was positive for anti-DFS70ab (Table 2, case #8) and the other was negative. One patient with anti-DFS70ab did not show the characteristic DFS pattern in IIF. This patient (Table 2, case #5) was confirmed to be positive for the antibody by the same immunoblotting analysis (discussed below).

The frequency of anti-DFS70ab in JDM was similar to that of the previous study by Schmeling *et al.*¹ (27.6% vs. 18.2%). The most important result is that there was only 1 (3.4%) isolated (monospecific) anti-DFS70ab-positive JDM patient in the present study (Table 2). This datum fits our previous data indicating that the monospecificity of anti-DFS70ab differs between AARD and non-AARD in adults.² Although the Schmeling study showed a relatively high frequency of anti-DFS70ab in patients with LS (13.8%), none of our patients with LS had that antibody. However, there was no statistical difference ($p < 0.287$ by Fisher's exact test). Interestingly, our LS patients often had anti-ssDNA antibodies (9/14, 64.3%).

Schmeling *et al.*¹ studied anti-DFS70ab also in patients with uveitis that was associated with JIA or that was idiopathic. Two of 19 (10.5%) JIA patients with uveitis

and 1 of 7 (14.3%) idiopathic uveitis patients had anti-DFS70ab. According to the study by Ravelli *et al.*,⁹ JIA patients with iridocyclitis very frequently had ANA (58/60, 96.7%). Surprisingly, in our present study, 5 of 10 JIA patients with iridocyclitis had anti-DFS70ab. The association of anti-DFS70ab with JIA with iridocyclitis in Japanese children should be confirmed by future studies. Although 1 of these 5 anti-DFS70ab-positive sera did not show a DFS pattern in IIF, the antibody-positive result was confirmed by immunoblotting analysis (Table 2, case #12). In a total of 53 examined sera, 4 sera had discrepant results between the DFS pattern in IIF and the anti-DFS70ab in ELISA. Schmeling *et al.*¹ also showed that, although the concordance rate of IIF patterns and chemiluminescence immunoassay results for anti-DFS70ab was >90%, IIF patterns do not always correspond to the other immunoassays results for anti-DFS70ab. Since recognizing the DFS pattern in IIF is not an easy task,¹⁰ a specific immunoassay for anti-DFS70ab should be used to confirm their presence. Moreover, clinicians should not over-interpret isolated ANA results in anti-DFS70ab-positive patients, but should pay attention to detecting concomitant disease-specific autoantibodies.

REFERENCES

1. Schmeling H, Mahler M, Levy DM, Moore K, Stevens AM, Wick J, et al. Autoantibodies to dense fine speckles in pediatric diseases and controls. *J Rheumatol* 2015;42:2419-26.
2. Muro Y, Sugiura K, Morita Y, Tomita Y. High concomitance of disease marker autoantibodies in anti-DFS70/LEDGF autoantibody-positive patients with autoimmune rheumatic disease. *Lupus*. 2008;17:171-6.
3. Deakin CT, Yasin SA, Simou S, Arnold KA, Tansley SL, Betteridge ZE, et al. Muscle biopsy findings in combination with myositis-specific autoantibodies aid prediction of outcomes in juvenile dermatomyositis. *Arthritis Rheumatol* 2016;68:2806-16
4. Arkachaisri T, Fertig N, Pino S, Medsger TA Jr. Serum autoantibodies and their clinical associations in patients with childhood- and adult-onset linear scleroderma. A single-center study. *J Rheumatol* 2008;35:2439-44.
5. Watanabe A, Kodera M, Sugiura K, Ususa T, Tan EM, Takasaki Y, et al. Anti-DFS70 antibodies in 597 healthy hospital workers. *Arthritis Rheum* 2005;32:2144-9.
6. Muro Y, Sugiura K, Nara M, Sakamoto I, Suzuki N, Akiyama M. High incidence

- of cancer in anti-small ubiquitin-like modifier activating enzyme antibody-positive dermatomyositis. *Rheumatology (Oxford)* 2015;54:1745-7.
7. Muro Y, Sugiura K, Akiyama M. A new ELISA for dermatomyositis autoantibodies: rapid introduction of autoantigen cDNA to recombinant assays for autoantibody measurement. *Clin Dev Immunol* 2013;2013:856815.
8. Muro Y, Nakashima R, Hosono Y, Sugiura K, Mimori T, Akiyama M. Autoantibodies to DNA mismatch repair enzymes in polymyositis/dermatomyositis and other autoimmune diseases: a possible marker of favorable prognosis. *Arthritis Rheumatol* 2014;66:3457-62.
9. Ravelli A, Felici E, Magni-Manzoni S, Pistorio A, Novarini C, Bozzola E, et al. Patients with antinuclear antibody-positive juvenile idiopathic arthritis constitute a homogeneous subgroup irrespective of the course of joint disease. *Arthritis Rheum* 2005;52:826-32.
10. Bentow C, Fritzler MJ, Mummert E, Mahler M. Recognition of the dense fine speckled (DFS) pattern remains challenging: results from an international internet-based survey. *Autoimmun Highlights* 2016;7:8