

## ***Replica-Exchange* Monte Carlo Method for Ar Fluid**

Takeshi NISHIKAWA,<sup>\*)</sup> Hiroshi OHTSUKA, Yuji SUGITA,  
Masuhiro MIKAMI\* and Yuko OKAMOTO

*Institute for Molecular Science, Okazaki, Aichi 444-8585, Japan*

<sup>\*)</sup> *National Institute of Materials and Chemical Research, Tsukuba 305-8565, Japan*

(Received October 11, 1999)

Lennard-Jones fluid systems such as Ar fluid have a huge number of states of energy local minima. Hence, a simulation by conventional algorithms based on the NVT canonical ensemble tends to get trapped in these local-minimum states. *Generalized-ensemble algorithms* alleviate this difficulty by performing a random walk in energy space. Multicanonical algorithm<sup>1)</sup> and simulated tempering<sup>2)</sup> are two of the most well-known such methods. However, in these methods we have to make iterations of preliminary runs in order to determine the (non-Boltzmann) weight factors. This process can be tedious and time-consuming. The *Replica-Exchange method*<sup>3)-7)</sup> (the method is also referred to as *replica Monte Carlo method*,<sup>4)</sup> *multiple Markov chain method*,<sup>6)</sup> and *parallel tempering*<sup>7)</sup>), is a new promising algorithm in which the above complicated procedure of weight determination is not necessary. We have applied the *Replica-Exchange Monte Carlo method*, *REMC*, to an Ar fluid system. We show that the new algorithm is indeed very effective and that accurate low-temperature canonical distributions can be obtained much more efficiently than conventional methods.

The generalized ensemble for replica-exchange method consists of  $M$  *non-interacting* replicas of the original system in the canonical ensemble at  $M$  different temperatures  $T_m$  ( $m = 1, \dots, M$ ). Because the replicas are non-interacting, the weight factor for the state  $X$  in this generalized ensemble is given by the product of Boltzmann factors for each replica (or at each temperature). We now consider exchanging a pair of replica  $i$  and replica  $j$  which are at temperatures  $T_m$  ( $= 1/k_B\beta_m$ ) and  $T_n$  ( $= 1/k_B\beta_n$ ), respectively. In order for this exchange process to converge towards an equilibrium distribution, it is sufficient to impose the detailed balance condition on the transition probability  $w(X \rightarrow X')$ :

$$W_{REMC}(X) w(X \rightarrow X') = W_{REMC}(X') w(X' \rightarrow X) . \quad (1)$$

The transition probability,  $w(X \rightarrow X')$  which satisfies the detailed balance condition is then given by:<sup>3)-8)</sup>

$$w(X \rightarrow X') \equiv w(x_m^{[i]} | x_n^{[j]}) = \begin{cases} 1 , & \text{for } \Delta \leq 0 , \\ \exp(-\Delta) , & \text{for } \Delta > 0 , \end{cases} \quad (2)$$

where

$$\Delta \equiv (\beta_n - \beta_m) \left( U(q^{[i]}) - U(q^{[j]}) \right) , \quad (3)$$

---

<sup>\*)</sup> E-mail address: nisikawa@ims.ac.jp

and  $U(q^{[i]})$  and  $U(q^{[j]})$  are the potential energies of replicas  $i$  and  $j$ , respectively. The effectiveness of the method was tested with Ar fluid. We remark that the exchanged quantity does not have to be temperature as long as it is in one-to-one correspondence with the replica (for instance, we can simulate the same system of  $M$  replicas with  $M$  different volumes). This is why we prefer the name replica-exchange method<sup>3)</sup> to parallel tempering<sup>7)</sup> (likewise, we prefer the term replica-exchange method to multiple Markov chain method,<sup>6)</sup> because molecular dynamics algorithm is also possible<sup>8)</sup>).




Fig. 1. Energy of *REMC* method.




Fig. 2. Energy of canonical MC.

We now present the results of our simulations based on the *REMC* algorithm and conventional canonical MC algorithm. We used the sixteen temperatures between  $T_H^* = 2.0$  and  $T_L^* = 0.72$ . Each replica consists of  $n^* = 0.5$ ,  $N = 256$ , Ar fluid. All the simulations were started from random configurations. Compared with conventional method, *REMC* method seems to achieve fast thermalization (compare Fig. 1 with Fig. 2) and explores energy-surface without getting trapped in states of energy local minima at low temperature (see Fig. 3).




Fig. 3. Average energy as a function of temperature.

The *Replica-Exchange* method is indeed very effective and that accurate low-temperature canonical distributions can be obtained much more efficiently than conventional methods.

## References

- 1) B. A. Berg and T. Neuhaus, Phys. Lett. **B267** (1991), 249.
- 2) E. Marinari and G. Parisi, Europhys. Lett. **19** (1992), 451.
- 3) a) K. Hukushima and K. Nemoto, J. Phys. Soc. Jpn. **65** (1996), 1604.  
b) K. Hukushima, H. Takayama and K. Nemoto, Int. J. Mod. Phys. **C7** (1996), 337.
- 4) R. H. Swendsen and J.-S. Wang, Phys. Rev. Lett. **57** (1986), 2607.
- 5) C. J. Geyer, in *Computing Science and Statistics: Proc. 23rd Symp. on the Interface*, ed. E. M. Keramidas (Interface Foundation, Fairfax Station, 1991), p. 156.
- 6) M. C. Tesi, E. J. J. van Rensburg, E. Orlandini and S. G. Whittington, J. Stat. Phys. **82** (1996), 155.
- 7) E. Marinari, G. Parisi and J. J. Ruiz-Lorenzo, in *Spin Glasses and Random Fields*, ed. A. P. Young (World Scientific, Singapore, 1998), p. 59.
- 8) Y. Sugita and Y. Okamoto, Chem. Phys. Lett., in press.