

A cross-sectional study on tobacco advertising, promotion and sponsorship (TAPS) and violations of tobacco sale regulations in Myanmar: Do these factors affect current tobacco use among Myanmar high school students?

Su Myat Cho¹, Yu Mon Saw^{1,2}, NyiNyi Latt³, Thu Nandar Saw⁴, Hein Htet⁵, Moe Khaing^{1,6}, Thet Mon Than⁶, Ei Mon Win⁷, ZawZaw Aung⁷, Tetsuyoshi Kariya^{1,2}, Eiko Yamamoto¹, Nobuyuki Hamajima¹

¹*Department of Healthcare Administration, Nagoya University Graduate School of Medicine, Nagoya, Japan*

²*Nagoya University Asian Satellite Campuses Institute, Nagoya, Japan*

³*AungMyinMyint Mo Hospital, Gyobingauk, Myanmar*

⁴*Department of Community and Global Health, Graduate School of Medicine, The University of Tokyo, Tokyo, Japan*

⁵*Department of Preventive and Social Medicine, University of Medicine, Mandalay, Myanmar*

⁶*Department of Medical Services, Ministry of Health and Sports, NayPyi Taw, Myanmar*

⁷*Department of Public Health, Ministry of Health and Sports, NayPyi Taw, Myanmar*

Corresponding author:

Yu Mon Saw

Department of Healthcare Administration, Nagoya University Graduate School of Medicine,
65 Tsurumai-cho, Showa-ku, Nagoya 466-8550, Japan

E-mail address: sawyumon@med.nagoya-u.ac.jp

Word count- 4111

ABSTRACT

Objectives: To examine the associations of current tobacco use with tobacco advertising, promotion and sponsorship (TAPS), and illicit tobacco sales exposures among Myanmar high school students.

Design: A quantitative, cross-sectional study.

Setting: Seven high schools from both urban and rural areas of four states and regions in Myanmar.

Participants: In total, 1,174 high school students (482 males and 692 females) were interviewed using a self-administered questionnaire.

Main outcome measure: Current tobacco use of participants, defined as using any kind of smoked or smokeless tobacco product at least one occasion within the past 30 days.

Results: The prevalence of TAPS exposure was 90.9% among high school students in Myanmar. Current tobacco use was positively associated with being over 14 years old (AOR 9.81; 95%CI 4.54–21.19), being male (AOR 28.06; 95%CI 13.29–59.25), exposure to any kind of TAPS (AOR 6.59; 95%CI 2.33–18.64), having seen any smoked tobacco product for sale inside or within 100 feet of the school premises (AOR 4.17; 95%CI 1.65–10.58), having seen the sale or gifting of any smoked tobacco product to minors (AOR 6.40; 95%CI 2.18–19.12), and having seen the sale or distribution of any smoked tobacco product by minors (AOR 2.42; 95% CI 1.42–4.10). Having ever received health education about tobacco use (AOR 0.45; 95% CI 0.27–0.78), or having a higher perception score of tobacco use (AOR 0.17; 95% CI 0.10–0.30) were negatively associated with current tobacco use.

Conclusions: There was an alarming prevalence of TAPS exposure among Myanmar high school students. TAPS exposure and violations of tobacco sale regulations were strong risk factors for current tobacco use among Myanmar high school students, while health education about tobacco products was reported as an effective protective factor. Specific smokeless tobacco sale regulations for minors are needed immediately in Myanmar.

Key words: Tobacco advertising, tobacco sale, current tobacco use, high school students, Myanmar

STRENGTHS AND LIMITATIONS OF THIS STUDY

- This study is the first in Myanmar to report the associations of current tobacco use with TAPS, and illicit tobacco sales exposures among high school students.
- The findings of this study cannot be generalized for the whole nation. It was conducted among 1,174 students from two states and two regions out of a total of seven states, seven regions, and one territory in Myanmar.
- This study could not confirm the causal relationships of current tobacco use among high school students because of the cross-sectional nature of our study.
- Due to the limitations of the cross-sectional study design, this study calls for further TAPS-related interventional and longitudinal studies to explore Myanmar adolescents' tobacco use behaviours.

INTRODUCTION

“Tobacco use,” defined as the use of any type of smoked or smokeless tobacco,¹⁾ is regarded as an important public health concern worldwide. Globally, it was estimated that 24 million (7.0%) adolescents aged between 13 and 15 years had smoked cigarettes in the past 30 days and 13.4 million (3.6%) had used smokeless tobacco products in the past 30 days during 2017.¹⁾ In the United States, it was reported that 4 million high school students were current tobacco users in 2018.²⁾ In Southeast Asia, the prevalence of adolescents' tobacco use in the past 30 days was 5.7% and that of smokeless tobacco use was 7.2% over the period of 2007 to 2017.¹⁾ A recent multi-national study reported that, between 2009 and 2013, adolescent smokeless tobacco use in the World Health Organization (WHO) South-East Asia Region (SEAR) was at its highest in Bhutan (23.2%), followed by Nepal (16.2%), Timor-Leste (14.2%), Myanmar (9.8%), India (9.0%), Sri Lanka (8.5%), the Maldives (6.2%), Bangladesh (5.9%), and Thailand (5.7%).³⁾ Thus, Myanmar ranked the fourth-highest for adolescent smokeless tobacco use among these nine countries. Myanmar is one of the countries with high prevalence of tobacco use among young population in the WHO SEAR countries.⁴⁾ Cigarette smoking among schoolchildren is much higher in Myanmar compared to other SEAR countries, i.e., Bangladesh, India, Maldives, Nepal, and Sri Lanka.⁵⁾ The use of smokeless tobacco product is also highly prevalent in Myanmar as compared to other countries.⁶⁾

Tobacco use is responsible for five million deaths every year globally, a figure that is expected to rise to 10 million per year by 2030.⁷⁾ In 2004, it was estimated that 600,000 people had died from the effects of second-hand smoke, accounting for roughly 1% of global mortality.⁸⁾ Adolescents are highly vulnerable to addiction to the nicotine in tobacco. A recent study has shown that the earlier people become dependent on nicotine, the more likely they are to become addicted to smoking in adulthood.⁹⁾ Moreover, nicotine consumption may negatively impact brain development during adolescence.¹⁰⁾ Studies have shown that adolescent smoking is associated with poor academic performance¹¹⁾ and attention and cognitive deficits.¹²⁾ Smokeless tobacco use is as dangerous as smoked forms of tobacco because it contains nicotine, carcinogens and other toxic chemicals.¹³⁾ Smokeless tobacco use has deleterious effects to oral health including the staining and discolouration of teeth, leukoplakia, erythroplakia and oral cancer.¹⁴⁾ A recent systematic review from India reported a positive association between smokeless tobacco use and various cancers (oral, oesophageal, pancreatic) in the South-East Asian Region and Eastern Mediterranean Region.¹⁵⁾ Moreover, another recent systematic review in the United States found an increased risk of heart disease and stroke among smokeless tobacco users.¹⁶⁾

Studies have shown that tobacco and smokeless tobacco use is highly prevalent among Myanmar high school students. Since 2001, the Global Youth Tobacco Survey

(GYTS) has been conducted every 3 to 5 years in Myanmar to monitor tobacco use among high school students. The findings from the 2016 GYTS conducted among high school students pointed out that the prevalence of the current use of smoked tobacco products and the prevalence of the current use of smokeless tobacco products was 10.6% and 5.7% respectively.¹⁷⁾ In parallel with the GYTS, Myanmar has been conducting the nationwide Global School-based Student Health Survey (GSHS) to monitor the understanding of health risk behaviours among high school students. The 2016 survey also reported that the prevalence of current tobacco smoking and current smokeless tobacco use among high school students was 7.2% and 8.5%, respectively.¹⁸⁾ Another study conducted among high school students in 2015 in Nay Pyi Taw, Myanmar, reported that 34.7% were smokers and 28.3% were smokeless tobacco users.¹⁹⁾ Therefore, the use of smoked and smokeless tobacco among high school students in Myanmar is an important public health issue as well as a social one. Moreover, all these studies also pointed out that most high school students began using tobacco before the age of 14. However, in Myanmar, parents tend to show less concern about their children becoming smokeless tobacco users (especially chewing betel quid with tobacco) because there is a widespread misconception that the use of smokeless tobacco is not as harmful as the use of cigarettes.¹³⁾ It is important to monitor the initiation and pattern of tobacco use among adolescents and youths, especially among high school students.

121 More than a decade ago, Myanmar signed the WHO Framework Convention on
122 Tobacco Control (FCTC) and enacted the first Tobacco Control Law in 2006, regulating
123 tobacco advertising, promotion, and sponsorship (TAPS) and tobacco sales to minors.¹⁷⁾
124 Subsequently, the restrictions on smoking in all indoor public places, the introduction of
125 graphic health warnings on tobacco product packaging, and the raising of tobacco product
126 taxes were all promulgated. Despite this, the prevalence of current tobacco use among
127 Myanmar high school students has not changed significantly over the past 15 years.¹⁷⁾

128 A comparison of the 2007 and the 2016 GSHS also revealed that the prevalence of
129 cigarette smoking among high school students had increased significantly, from 2.0% to
130 6.7%.¹⁸⁾ Although the Tobacco Control Law has banned tobacco sales to minors, adolescent
131 smokers can still buy cigarettes from large stores, retail shops, or street vendors very easily.¹⁷⁾
132 These alarming findings indicate the failure of efforts to control tobacco consumption among
133 young people in Myanmar.

134 Although the Tobacco Control Law in Myanmar prohibits TAPS activities by the
135 tobacco industry, TAPS activities are still common. Most tobacco companies distribute
136 tobacco products and personal goods with tobacco product labels either free of charge or as
137 gifts. According to the 2016 Myanmar GYTS conducted among high school students, 8.7% of
138 boys and 3.6% of girls reported that tobacco companies had offered free tobacco products,

and 7.3% of boys and 4.2% of girls reported owning something with tobacco branding or a tobacco logo.¹⁷⁾ Furthermore, 83.4% of the students reported noticing someone using tobacco products on television or in videos and movies.¹⁷⁾

Researchers from other parts of the world have reported that TAPS exposure can affect adolescent smoking behaviours.²⁰⁻²³⁾ A positive association between exposure to cigarette advertisements and initiating smoking has been reported among Indonesian students.²⁰⁾ A longitudinal study conducted in Germany also pointed out that, with every additional 10 tobacco advertisements, the adjusted relative risk for established smoking and daily smoking was raised by 38% and 30%, respectively.²⁴⁾ Adolescent students, in a phase of life where curiosity is at its peak, are vulnerable to adopting smoked or smokeless tobacco use.^{13, 25)}

At present, research in Myanmar is limited where concerns the patterns of high school students' tobacco use and their connection to TAPS exposure. Our previous study,²⁶⁾ conducted among the same participants, revealed the low awareness of the Tobacco Control Law among Myanmar high school students, but we did not examine how TAPS exposure and violations of tobacco sales regulations might affect Myanmar high school students' tobacco use. The present study aims to investigate (1) the prevalence of TAPS and illicit tobacco sale

exposures and (2) their associations with current tobacco use among Myanmar high school students.

MATERIALS AND METHODS

Study population

A cross-sectional study was conducted among grade 10 and 11 high school students from seven high schools in Shan State, Mon State, Bago region, and Magway region in Myanmar. A simple random sampling technique using a drawing method was applied to select the study areas and schools. The details of the sampling procedure have been described elsewhere.²⁶⁾ In total, 1,339 high school students answered a self-administrated questionnaire. Of these, 165 were excluded due to missing or incomplete responses to TAPS exposure questions. In sum, the total number of participants was 1,174 (482 males and 692 females) and the response rate was 87.7%.

Data collection

Data were collected by using a pre-tested, anonymous, paper and pencil self-administered questionnaire. The questionnaire in the Myanmar language contained 40 questions, covering nine components: 1) background information, 2) experience with tobacco products, 3) exposure to second-hand smoking, 4) perception of smoked and smokeless tobacco products,

173 5) sale of tobacco, 6) health warnings and information, 7) tobacco advertisement, promotion,
174 and sponsorship, 8) smoke-free areas, and 9) the Tobacco Control Law and its enforcement.

175 ***Study measures***

176 ***Dependent variable***

177 The outcome variable was “current tobacco use” among high school students. It was defined
178 as the use of any kind of smoked or smokeless tobacco product on at least one occasion
179 within the 30 days preceding the survey.

180 ***Independent variables***

181 After controlling socio-demographic characteristics, smoking exposure at home and school,
182 receiving health education about tobacco use, and the perception of tobacco use as covariates,
183 the independent variables in this study were participants’ exposure to any kind of TAPS and
184 illicit tobacco sale exposures. “Exposure to any kind of TAPS” was defined using the
185 following variables: 1) having seen or heard tobacco advertisements or sponsorship in any
186 form, 2) having seen any goods displaying the label of any tobacco product being used in
187 promotion, and 3) having seen any toy, comestible, or wares made in the form of any tobacco
188 product. If a student had had at least one TAPS exposure, the response was counted as a
189 “Yes,” and if they had never experienced exposure, their response was counted as a “No.”

For illicit tobacco sale exposures, we measured four different types of tobacco sales to students contravening several of the tobacco sale regulations prohibited by Myanmar Tobacco Control Law. These four variables were: 1) having seen any smoked tobacco product for sale inside or within 100 feet of the school premises within the last 12 months, 2) having seen the sale or gifting of any smoked tobacco product to minors within the last 12 months, 3) having seen the sale or distribution of any smoked tobacco product by minors within the last 12 months, and 4) having seen the sale of cigarettes singly or in packs containing less than 20 within the last 12 months.

Study analysis

The data were coded, entered, and analysed using the Statistical Package for Social Science (SPSS) software program version 24.0 (IBM SPSS Inc.). Categorical data were analysed by chi-square tests for hypothesis testing. For multivariable logistic regression, all the variables were re-coded on a dichotomous scale. All analyses were two-tailed, setting $p < 0.05$ as the significance value.

RESULTS

Table 1 shows the characteristics of the study participants by gender. Most of the participants (84.7%) were younger than or equal to 15 years of age. More than two-thirds of the students

207 (68.5% of males and 67.1% of females) were grade 10 students. Of those who took part,
208 25.3% of male and 1.3% of female students were current users of smoked or smokeless
209 tobacco at the time of the survey. Male students had more exposure to parental smoking
210 (6.0%), sibling smoking (11.4%), and peer smoking (9.3%) than did female students. Nearly
211 two-thirds of females (66.0%) reported receiving health education about tobacco use,
212 compared to 52.7% of males. Regarding the perception of tobacco use, out of eight items, we
213 set the mean score of seven as the cut-off point. Overall, 68.7% of males and 83.2% of
214 females scored more than seven.

215 Table 2 presents different kinds of TAPS exposures and its associations with current
216 tobacco use among the study participants. In total, 90.9% had TAPS exposure in any form.
217 The findings indicate that 71.7% had seen or heard of tobacco advertisement and sponsorship
218 in any form, and 68.1% had seen goods displaying the label of any tobacco product being
219 used in promotions. One out of three students had seen toys, comestibles or wares made in the
220 form of any tobacco product. More than one-third of the participants (35.3%) had seen or
221 heard tobacco advertisements in sponsorship or support of sports, funfairs, exhibitions, or
222 other social activities.

223 Table 3 describes illicit tobacco sale exposures and their associations with current
224 tobacco use among the study participants. Within the last 12 months, more than 80.0% had

been exposed to the sale of any smoked tobacco product inside or within 100 feet of the school premises, the sale or gifting of any smoked tobacco product to minors, or the sale of cigarettes singly or in packs of less than 20. Nearly 56.0% had seen the sale or distribution of any smoked tobacco product by minors.

Table 4 presents the unadjusted odds ratio (UOR), adjusted odds ratio (AOR), and 95% confidence intervals (CI) of current tobacco use among Myanmar high school students. In the binary logistic regression, being a grade 11 student (UOR 3.24; 95% CI 2.24-4.70) and having seen the sale of cigarettes, either singly or in packs of less than 20 (UOR 7.45; 95%CI 2.34–23.70) were associated with current tobacco use.

From the multiple logistic regression, current tobacco use was found to be positively associated with being over 14 years old (AOR 9.81; 95%CI 4.54–21.19); being male (AOR 28.06; 95%CI 13.29–59.25); being exposed to any kind of TAPS (AOR 6.59; 95%CI 2.33–18.64); having seen any smoked tobacco product for sale inside or within 100 feet of the school premises within the last 12 months (AOR 4.17; 95%CI 1.65–10.58); having seen the sale or gifting of any smoked tobacco product to minors within the last 12 months (AOR 6.40; 95%CI 2.18–19.12); and having seen the sale or distribution of any smoked tobacco product by minors within the last 12 months (AOR 2.42; 95% CI 1.42–4.10). Having ever received health education about tobacco use (AOR 0.45; 95% CI 0.27–0.78) and having a higher

perception score with regards to tobacco use (AOR 0.17; 95% CI 0.10–0.30) were negatively associated with current tobacco use (Table 3).

DISCUSSION

To the best of our knowledge, this is the first study to report the associations of current tobacco use with TAPS and illicit tobacco sale exposures among high school students in Myanmar. The majority of the study participants reported having been exposed to TAPS and illicit tobacco sales. TAPS exposure and illicit tobacco sale exposures increase the odds of current tobacco use among high school students.

In this study, a high prevalence of TAPS exposure (91%) was reported among high school students in Myanmar, a country that has completely banned all forms of direct or indirect TAPS, including Corporate Social Responsibility (CSR) activities by the tobacco industry.²⁷⁾ Myanmar has a well-established Tobacco Control Law named the “Control of Smoking and Consumption of Tobacco Product Law,” which has been enacted since 2006.²⁷⁾ However, the monitoring, reporting and punishment of TAPS activities prohibited by the law are not common in Myanmar. Other studies conducted in Myanmar also pointed out that the awareness of the tobacco control law among high school students was low and that the lack of tobacco control law enforcement was in a critical state.^{19,26)} It is not rare to see sponsored events and CSR activities executed by tobacco companies, violating the TAPS regulations

261 and reframing tobacco products' image among Myanmar youths.^{19,26)} The global tobacco
262 industry has been focusing on expanding its market in developing countries that have low
263 tobacco taxes, partial TAPS bans, and weak law enforcement of TAPS regulations, rather than
264 in developed countries with high tobacco taxes, comprehensive and complete TAPS bans, and
265 the strict implementation of TAPS regulations.²⁸⁻³¹⁾

266 The weakness of tobacco control law enforcement in Myanmar creates opportunities
267 for tobacco companies and retailers to violate it. Nigerian researchers have reported that
268 exposure to events sponsored by the tobacco industry was associated with current cigarette
269 use and demonstrated the importance of the tobacco control law and its enforcement in
270 reducing tobacco use. Our findings highlighted the urgent need to enforce this law in
271 Myanmar to reduce TAPS exposure among adolescents in order to decrease tobacco use. It
272 also encourages Myanmar policymakers to formulate specific TAPS regulations addressing
273 newly developed smoking and smokeless tobacco products and complete comprehensive
274 TAPS bans, including cross-border TAPS. Local authorities need to monitor TAPS among
275 Myanmar youths strictly and to punish the tobacco companies, stores, and retailers violating
276 TAPS regulations.

277 High school students exposed to TAPS in any form were six times more likely to be
278 current tobacco users than those without any TAPS exposure. A recent study from Myanmar

279 has reported that only half of the high school students had heard about the Tobacco Control
280 Law, and none of the study participants had ever heard of any reporting of or punishment for
281 any violation of the Tobacco Control Law that bans TAPS activities and tobacco sales among
282 minors.²⁶⁾ Our study suggests that Myanmar high school students should be properly informed
283 not only about the dangers of TAPS exposure but also about the country's Tobacco Control
284 Law and its punishments.

285 In this study, the prevalence of current tobacco use among Myanmar high school
286 students was 11.2%, which is higher than the national figure of 9.8% reported in the 2016
287 WHO GSHS for students between 13 and 17 years old in Myanmar.¹⁸⁾ Our study also reported
288 that male high school students were more likely to be current tobacco users than their female
289 peers were. A sub-national-level study also reported a considerably high prevalence of
290 smoked (34.7%) and smokeless tobacco use (28.3%) among high school students.¹⁹⁾ The
291 findings suggest Myanmar policymakers to consider both smoked and smokeless tobacco use
292 among adolescents as a public health priority and to put more effort into implementing control
293 measures.

294 In Myanmar, it has been reported that there were only three full-time staff assigned
295 to national tobacco control, meaning they were each responsible for 2,080,000 smokers.³²⁾
296 The country also has an insufficient government budget for tobacco control and there is no

297 health promotion funding mechanism in place for the use of tobacco taxes.³²⁾ To tackle the
298 current situation in Myanmar, the effective implementation of tobacco control measures is
299 needed to reduce current tobacco use, and human and financial resources for national tobacco
300 control should be improved.

301 Increasing tobacco tax, as recommended for implementation of Article 6 of the WHO
302 FCTC, functions as a cost-effective demand-reducing measure in global tobacco control.³²⁾
303 However, cigarettes in Myanmar are relatively cheap, indicating that the country's tobacco
304 tax policies need to be re-evaluated. Among the Association of Southeast Asian Nations
305 (ASEAN), the price of the most popular local cigarette brand in Myanmar is the second
306 lowest (0.6 USD/20-stick pack), and that of the most popular foreign brand in Myanmar is the
307 fourth highest (2.11 USD/20-stick pack).³²⁾ This indicates a gap in tax differences between
308 local and foreign brands of cigarettes. The price at which youths in Myanmar are deterred
309 from smoking is reportedly 11 USD/20-stick pack, the lowest among the ASEAN.³²⁾
310 Therefore, the Myanmar government should reevaluate tobacco tax levels and adjust tax
311 policies to decrease tobacco use and TAPS exposures.

312 In this study, students who had seen any smoked tobacco product for sale inside or
313 within 100 feet of the school premises during the last 12 months were four times more likely
314 to be current tobacco users than their peers were. It is illegal to sell any kind of smoked

315 tobacco products inside or within 100 feet of school premises in Myanmar.²⁷⁾ This finding
316 implies that the lack of monitoring and reporting of illicit tobacco sales inside or near school
317 premises exerts an influence on current tobacco use among Myanmar high school students. In
318 addition, these illicit tobacco sales provide easy access to tobacco for students as well as for
319 school personnel, triggering second-hand smoking exposure and student curiosity about
320 tobacco use. Therefore, school personnel, students, and parents should monitor and report any
321 illicit tobacco sales to local authorities and actions should be taken according to the rules and
322 regulations.

323 However, there is no specific regulation on smokeless tobacco sales and distributions
324 to and by minors in Myanmar, the sale or gifting of any smoked tobacco product to minors
325 and the sale or distribution of any smoked tobacco product by minors are illegal.²⁷⁾
326 Nonetheless, 82.5% of participants in this study reported that they had seen someone selling
327 or gifting of any smoked tobacco product to a minor within the last 12 months, and 55.6% of
328 the participants reported having witnessed the sale or distribution of any smoked tobacco
329 product by minors within the last 12 months. The study also revealed that having seen these
330 two illicit tobacco activities within the last 12 months was significantly associated with
331 current tobacco use among the study participants. A study of adolescents in the United States
332 has also reported an association between smokeless tobacco use and smoking.³³⁾ In order to

reduce current tobacco use among Myanmar high school students, this study recommends the urgent need of specific smokeless tobacco regulations concerning sales made to and by minors in Myanmar, and strict law enforcement on sales and distributions all smoked and smokeless tobacco to and by minors in the country.

Another common violation of the Tobacco Control Law in Myanmar is the sale of cigarettes singly or in packs of less than 20. Despite the significant price difference³²⁾ both foreign and local brands become affordable for smokers when they are sold singly or in packs of less than 20. Such sales are contrary to the demand-reducing tobacco control measures of the national policy and negatively affect the tobacco use of all age groups, especially adolescents.

In addition, other smoked and smokeless tobacco products, such as cheroots, cigars, pipes, betel quids, etc., can be purchased singly or in small quantities in Myanmar. To reduce tobacco use in all age groups, our study recommends that the 2006 Tobacco Control Law be updated to address the sale and purchase of all forms of smoked and smokeless tobacco products in small quantities, along with strict law enforcement, especially among minors.

The findings of this study imply that effective health education and high perception of tobacco use can be effective factors in combating the current tobacco use among Myanmar high school students. In this study, students who had received health education about tobacco

use and those with a higher perception score of tobacco use were less likely to be current tobacco users than their counterparts. Nearly two-thirds of the participants had received health education about tobacco use, and nearly 80% had a higher perception score. Health education programs and tobacco control measures targeting the young should address the newly developed and popular tobacco products like electronic cigarettes, shisha, pipes, menthol and fruit-flavoured cigarettes etc. Youths' awareness of not only the harms of tobacco products but also the country's Tobacco Control Law should be promoted.

The study findings also revealed that 10.7% of current tobacco users were 14 years old or less. Another Myanmar researcher has reported that the average age for first tobacco use is 14 years.¹⁹⁾ Therefore, tobacco control intervention measures should be introduced to Myanmar high school students before the age of 14. Awareness of the Tobacco Control Law remains quite low in Myanmar.^{19,26)} Furthermore, the provision of health education via youth-friendly media, such as the internet, mobile applications, and social networking services, may attract more attention from high school students.

In contrast to other studies,^{19,26)} parental smoking, sibling smoking, and peer smoking were not found to be associated with current tobacco use in this study. This may be because only the smoking status of parents, siblings and peers of the participants was assessed in this study and most of the study participants did not have exposure to parents, siblings, or peers

who smoked. Having high knowledge and perception about tobacco use may also have prevented them from using tobacco.¹⁹⁾

Despite being the very first study in Myanmar to report the associations between current tobacco use, TAPS and illicit tobacco sale exposure among Myanmar high school students, the present study did not explore the associations of first-time or daily tobacco use, and TAPS or illicit tobacco sales among Myanmar high school students. This study was conducted among 1,174 high school students from two states and two regions out of a total of seven states, seven regions and one territory of Myanmar, and its findings cannot be generalized for the whole nation. Due to the limitations of the cross-sectional nature of our study, we recommend further interventional or longitudinal studies of TAPS and sales exposure for a better understanding of adolescent tobacco use in Myanmar.

CONCLUSIONS

This study reported high prevalences of TAPS and illicit tobacco sales exposures among Myanmar high school students. Current tobacco use among Myanmar high school students was statistically associated with overall TAPS exposure. Violations of tobacco sales regulations were reported to be strong risk factors for current smoked and smokeless tobacco use among Myanmar high school students. Our findings highlight that Myanmar's Tobacco Control Law enforcement is in an alarming state and requires urgent improvement. Sales and

387 purchase of not only cigarettes but also all forms of smoked and smokeless tobacco products
388 in small quantities should be regulated. Specific smokeless tobacco sale regulations for
389 minors are urgently needed in the country.

390 **ABBREVIATIONS**

391 TAPS Tobacco Advertising, Promotion, and Sponsorship

392 AOR Adjusted Odds Ratios

393 UOR Unadjusted Odds Ratios

394 CI Confidence Intervals

395 WHO World Health Organization

396 GYTS Global Youth Tobacco Survey

397 GSHS Global School-based Student Health Survey

398 FCTC Framework Convention on Tobacco Control

399 SPSS Statistical Package for Social Science

400 IBM International Business Machines

401 CSR Corporate Social Responsibility

402 ASEAN Association of South-East Asian Nations

403 USD The United States Dollar

404

ACKNOWLEDGEMENTS

The authors would like to thank Ministry of Health and Sports and Ministry of Education in Myanmar for granting a permission to collect the data at high schools. We also would like to thank the local authorities in the survey areas, the staff from participating schools, last but not least, all the high school students those who participated voluntarily in this study.

Funding statement

This research received no specific grant from any funding agency in the public, commercial or not-for-profit sectors.

Data sharing statement

No additional data are available.

Author contributions

YMS and NNL conceptualised the study and designed the study. NNL, and TNS contributed to data collection and data acquisition. YMS, TNS, HH, and SMC performed data analyses and data interpretations. YMS, HH, and SMC wrote the first draft of the manuscript. MK, TMT, EMW, TK, EY, and ZZA contributed to data acquisition. NH contributed to the study design, data interpretation, and revising the manuscript. YMS, TNS, SMC, and NH revised and edited the manuscript. All authors had full access to the data, and take responsibility to the accuracy of data analysis. All authors approved the final manuscript and agreed to submit it for publication and take accountability.

Ethical approval and consent to participate

This study was ethically approved by the Department of Medical Services, Ministry of Health and Sports, Myanmar (Letter No. 617 of Planning/Research issued on August 26, 2015), and the Ministry of Education, Myanmar (Letter No. 12125 of Information/Research issued on October 19, 2015), as well as the ethical review committee of Nagoya University School of Medicine (No. 6518 issued on August 31, 2015). To conduct this school- based survey, permissions from Ministry of Education, Regional Offices of Basic Education, Ministry of Health and Sports, local educational steering committees and authorities, the schools' authorities, the headmasters of participated schools and local Parents-Teacher Associations were obtained. The survey procedure was approved by Ministry of Education and Ministry of Health and Sports. After thoroughly explaining the study's objectives, contents of the survey questionnaire, and rights of the study participants, the written-informed consents from local educational steering committees and authorities, the schools' authorities, the headmasters of participated schools, local Parents-Teacher Associations, and parents were obtained. One week prior to the survey, the information sheet and the written-informed consents that stating the study's objectives, the survey's procedure and the contents of the questionnaires, and the rights of the study participants were sent to parents. Researchers also explained the study's objectives, contents of the survey questionnaire, the voluntary nature and procedure of the survey, and the rights of the participants to collaborators, students and teachers before conducting the survey. All data collection and analytical processes remain anonymous for privacy and confidentiality. The locations, names, and numbers of the eligible participants of the schools involved were not documented.

Competing interests

All authors declared no conflicts of interest for this study.

Additional file

449 There is no additional files.

450

REFERENCES

1. World Health Organization. *WHO global report on trends in prevalence of tobacco smoking 2000–2025, second edition*.
<https://www.who.int/tobacco/publications/surveillance/trends-tobacco-smoking-second-edition/en/>. Accessed May 5, 2019.
2. Gentzke AS, Creamer M, Cullen KA, et al. Vital Signs: Tobacco Product Use among Middle and High School Students - United States, 2011–2018. *MMWR Morb Mortal Wkly Rep*. 2019;68(6):157–164. doi:10.15585/mmwr.mm6806e1.
3. Sinha D N, Palipudi K M, Jones C K, et al. Levels and trends of smokeless tobacco use among youth in countries of the World Health Organization South-East Asia Region. *Indian J Cancer*. 2014;51(5):50-53. doi:10.4103/0019-509X.147472.
4. World Health Organization, Regional Office for South-East Asia. (2015). Monitoring tobacco control among youth in countries of the South-East Asia Region: 2014. World Health Organization. <https://apps.who.int/iris/handle/10665/155159>. Accessed Nov 24, 2019.
5. Tun NA, Chittin T, Agarwal N, et al. Tobacco use among young adolescents in Myanmar: Findings from global youth tobacco survey. *Indian J Public Health* 2017; 61(Suppl 1):S54-S59. doi: 10.4103/ijph.IJPH_236_17.
6. World Health Organization, Regional Office for South-East Asia. (2012). Expert group meeting on smokeless tobacco control and cessation: New Delhi, India, 16-17 August 2011. WHO Regional Office for South-East Asia.
<https://apps.who.int/iris/handle/10665/205054>. Accessed Nov 24, 2019.
7. World Health Organization. *WHO global report: mortality attributable to tobacco*.
https://www.who.int/tobacco/publications/surveillance/rep_mortality_attributable/en/. Accessed April 27, 2019.
8. Oberg M, Jaakkola MS, Woodward A, Peruga A, Prüss-Ustün A. Worldwide burden of disease from exposure to second-hand smoke: a retrospective analysis of data from 192 countries. *Lancet*. 2011;377(9760):139-146. doi:10.1016/S01406736(10)61388-8.
9. Selya AS, Dierker L, Rose JS, Hedeker D, Mermelstein RJ. Early-emerging nicotine dependence has lasting and time-varying effects on adolescent smoking behavior. *Prev Sci*. 2016;17(6):743-750. doi:10.1007/s11121-016-0673-0.
10. Centers for Disease Control and Prevention. *The health consequences of smoking- 50 years of progress: A report of the Surgeon General*.
<https://www.surgeongeneral.gov/library/reports/50-years-of-progress/full-report.pdf>. Accessed April 29, 2019.

11. Robert PO, Kuipers MAG, Rathmann K, et al. Academic performance and adolescent smoking in 6 European cities: the role of friendship ties. *Int J Adolesc Youth*. 2019;24(1):125-135. doi:10.1080/02673843.2018.1475288.
12. Treur JL, Willemsen G, Bartels M, et al. Smoking during adolescence as a risk factor for attention problems. *Biol Psychiatry*. 2015;78(9):656-63. doi:10.1016/j.biopsych.2014.06.019.
13. Kyaing NN, Sein T, Sein AA, Htike MMT, Tun A, Shein NN. Smokeless tobacco use in Myanmar. *Indian J Cancer*. 2012; 49(4): 347–351. doi:10.4103/0019-509X.107727.
14. Muthukrishnan A, Warnakulasuriya S. Oral health consequences of smokeless tobacco use. *Indian J Med Res*. 2018;148(1):35–40. doi:10.4103/ijmr.IJMR_1793_17.
15. Gupta S, Gupta R, Sinha DN, Mehrotra R. Relationship between type of smokeless tobacco & risk of cancer: A systematic review. *Indian J Med Res*. 2018;148(1):56–76. doi:10.4103/ijmr.IJMR_2023_17.
16. Rostron BL, Chang JT, Anic GM, Tanwar M, Chang CM, Corey CG. Smokeless tobacco use and circulatory disease risk: a systematic review and meta-analysis. *Open Heart*. 2018;5(2):e000846. doi:10.1136/openhrt-2018-000846.
17. World Health Organization. *Report of fifth Global Youth Tobacco Survey (GYTS) Myanmar, 2016*. <https://apps.who.int/iris/handle/10665/274351>. Accessed May 2, 2019.
18. World Health Organization. *Report of second Global School-based Student Health Survey in Myanmar 2016*. <http://www.who.int/iris/handle/10665/274350>. Accessed May 2, 2019.
19. Myint MNT, Yamamoto E, Ko MH, Khaing M, Reyer JA, Hamajima N. Knowledge, attitude, and usage pattern of tobacco among high school students in Nay Pyi Taw, Myanmar. *Nagoya J Med Sci*. 2019;81(1):65–79. doi:10.18999/nagjms.81.1.65.
20. Prabandari YS, Dewi A. How do Indonesian youth perceive cigarette advertising? A cross-sectional study among Indonesian high school students. *Glob Health Action*. 2016;9:30914. doi:10.3402/gha.v9.30914.
21. Chido-Amajuoyi OG, Mantey DS, Clendennen SL, Pérez A. Association of tobacco advertising, promotion and sponsorship (TAPS) exposure and cigarette use among Nigerian adolescents: implications for current practices, products and policies. *BMJ Glob Health*. 2017;2(3):e000357. doi:10.1136/bmjgh-2017-000357.
22. Braverman MT, Aaro LE. Adolescent smoking and exposure to tobacco marketing under a tobacco advertising ban: findings from 2 Norwegian national samples. *Am J Public Health*. 2004;94(7):1230–1238.

23. Soneji S, Ambrose BK, Lee W, Sargent J, Tanski S. Direct-to-consumer tobacco marketing and its association with tobacco use among adolescents and young adults. *J Adolesc Health*. 2014;55(2):209–215. doi:10.1016/j.jadohealth.2014.01.019.
24. Morgenstern M, Sargent JD, Isensee B, Hanewinkel R. From never to daily smoking in 30 months: the predictive value of tobacco and nontobacco advertising exposure. *BMJ Open*. 2013;3(6): e002907. doi: <http://dx.doi.org/10.1136/bmjopen-2013-002907>.
25. Akl EA, Jawad M, Lam WY, Co CN, Obeid R, Irani J. Motives, beliefs and attitudes towards waterpipe tobacco smoking: a systematic review. *Harm Reduct J*. 2013;10:12. doi:10.1186/1477-7517-10-12.
26. LattNN, Saw YM, Myat Cho S, Kariya T, Yamamoto E, Hamajima N. Tobacco Control Law awareness, enforcement, and compliance among high school students in Myanmar. *Nagoya J Med Sci*. 2018;80(3):379–389. doi:10.18999/nagjms.80.3.379.
27. Ministry of Health. *The control of smoking and consumption of Tobacco Product Law*. <https://seatca.org/dmdocuments/Myanmar%20-%20TC%20Law.pdf>. Accessed May 17, 2019.
28. Gilmore AB, Fooks G, DropeJ, Bialous SA, Jackson RR. Exposing and addressing tobacco industry conduct in low-income and middle-income countries. *Lancet* 2015;385(9972):1029–1043. doi:10.1016/S0140-6736(15)60312-9.
29. Gostin LO. FDA regulation of tobacco: politics, law, and the public's health. *JAMA*. 2009;302(13):1459-1460. doi:10.1001/jama.2009.1421.
30. Jamison N, Tynan M, MacNeil A, *et al*. Federal and state cigarette excise taxes - United States, 1995-2009. *MMWR Morb Mortal Wkly Rep*. 2009;58(19):524–527.
31. Chaloupka FJ, Yurekli A, Fong GT. Tobacco Taxes as a tobacco control strategy. *Tob Control* 2012;21(2):172–180. doi:10.1136/tobaccocontrol-2011-050417.
32. Southeast Asia Tobacco Control Alliance. The Tobacco Control Atlas: ASEAN Region, Fourth Edition. <https://seatca.org/dmdocuments/Tobacco%20Control%20Atlas%20ASEAN%20Region%204th%20Ed%20Feb%202019.pdf>. Accessed May 19, 2019.
33. Wiener RC. Association of smokeless tobacco use and smoking in adolescents in the United States: an analysis of data from the Youth Risk Behavior Surveillance System survey, 2011. *J Am Dent Assoc*. 2013;144(8):930–938.

553 **Table List**

554 **Table 1** Characteristics of study participants

555 **Table 2** Different kinds of TAPS ^{a)} exposures among study participants

556 **Table 3** Illicit tobacco sale exposures among study participants

557 **Table 4** Odds ratio (OR) and 95% confidence interval (CI) of current tobacco use among
558 study participants.

TABLES

Table 1 Characteristics of study participants (N=1,174)

Characteristics	Male (N=482)		Female (N=692)		Total (N=1,174)	
	N	%	N	%	N	%
Age						
≤14 years	195	40.5	307	44.4	502	42.8
15 years	199	41.3	293	42.3	492	41.9
16 years	61	12.7	91	13.2	152	12.9
17 years	21	4.4	1	0.1	22	1.9
≥18 years	6	1.2	0	0.0	6	0.5
Grade						
10	330	68.5	464	67.1	794	67.6
11	152	31.5	228	32.9	380	32.4
Current tobacco use						
No	360	74.7	683	98.7	1,043	88.8
Yes	122	25.3	9	1.3	131	11.2
Parent smoking						
No	453	94.0	662	95.7	1,115	95.0
Yes	29	6.0	30	4.3	59	5.0
Sibling smoking						
No	427	88.6	652	94.2	1,079	91.9
Yes	55	11.4	40	5.8	95	8.1
Peer smoking						
No	437	90.7	641	92.6	1,078	91.8
Yes	45	9.3	51	7.4	96	8.2
Ever received health education about tobacco use						
No	228	47.3	235	34.0	463	39.4
Yes	254	52.7	457	66.0	711	60.6
Perception of tobacco use						
≤7	151	31.3	116	16.8	267	22.7
>7	331	68.7	576	83.2	907	77.3

559

560

Table 2 Different kinds of TAPS ^{a)} exposures among study participants (N=1,174)

TAPS ^{a)}	Use ^{b)} (N=131)		No use ^{c)} (N=1,043)		Total (N=1,174)	
	N	%	N	%	N	%
Ever seen or heard tobacco advertising and sponsorship in any form **						
No	25	19.1	307	29.4	332	28.3
Yes	106	80.9	736	70.6	842	71.7
Ever seen tobacco advertising signboards (vinyl, LED, stickers, etc.)						
No	128	97.7	994	95.3	1,122	52.0
Yes	3	2.3	49	4.7	52	4.4
Ever seen tobacco advertising drawing or painting (on vehicles, on walls, on boards, etc.)						
No	118	90.1	925	88.7	1,043	88.8
Yes	13	9.9	118	11.3	131	11.2
Ever seen tobacco advertising in journals, magazines, newspapers, and pamphlets						
No	125	95.4	998	95.7	1,123	95.7
Yes	6	4.6	45	4.3	51	4.3
Ever seen broadcasting of tobacco advertisements (TV, radio, internet, social network like Facebook, etc.)						
No	120	91.6	982	94.2	1,102	93.9
Yes	11	8.4	61	5.8	72	6.1
Ever seen or heard about the distribution of tobacco products free of charge or as gifts						
No	124	94.7	953	91.4	1,077	91.7
Yes	7	5.3	90	8.6	97	8.3
Ever seen or heard distributions of personal goods with tobacco product labels free of charge or as gifts ***						
No	64	48.9	702	67.3	766	65.2
Yes	67	51.1	341	32.7	408	34.8
Ever seen or heard about tobacco advertising with lucky draw, exchange of old cigarette pack with new ones, bonus to sellers, car stickers, etc. *						
No	125	95.4	1,025	98.3	1,150	98.0
Yes	6	4.6	18	1.7	24	2.0
Ever seen any goods with the label of any tobacco product being used in promotions (clothes, hats, lighters, key chains, tissue boxes, stationeries, kitchen utensils, etc.)						
No	34	26.0	340	32.6	703	67.4

Yes	374	31.9	97	74.0	800	68.1
Ever seen any toy, comestible or wares made in the form of any tobacco product (toys, chewing gums, sweet sticks, key chains, lighters, balloons, etc.)						
No	88	67.2	702	67.3	790	67.3
Yes	43	32.8	341	32.7	384	32.7
Ever heard about the announcements of tobacco advertisement at fairs and festivals *						
No	128	97.7	1,040	99.7	1,168	99.5
Yes	3	2.3	3	0.3	6	0.5
Ever seen or heard about the tobacco advertising as sponsorship or support to sports, funfairs, exhibitions, or any social activities **						
No	70	53.4	689	66.1	759	64.7
Yes	61	46.6	354	33.9	415	35.3
Exposure to any kind of TAPS ^{a)}						
No	8	6.1	99	9.5	107	9.1
Yes	123	93.9	944	90.5	1,067	90.9

*p<0.05, **p<0.01, ***p<0.001; ^{a)}Tobacco advertisement, promotion, and sponsorship, ^{b)}Current tobacco use,

^{c)}No current tobacco use

561

562

Table 3 Illicit tobacco sale exposures among study participants (N=1,174)

Illicit tobacco sale	Use ^{b)}		No use ^{c)}		(N=1,174)
	(N=131)		(N=1,043)		
	N	%	N	%	
Had seen any smoked tobacco product for sale inside or within 100 feet of the school premises within the last 12 months ***					
No	92	19.1	93	13.4	
Yes	390	80.9	599	86.6	
Had seen the sale or gifting of any smoked tobacco product to minors ^{a)} within the last 12 months ***					
No	86	17.8	120	17.3	
Yes	396	82.2	572	82.7	
Had seen the sale or distribution of any smoked tobacco product by minors ^{a)} within the last 12 months ***					
No	209	43.4	312	45.1	
Yes	273	56.6	380	54.9	
Had seen the sale of cigarettes singly or in packs less than 20 cigarettes within the last 12 months ***					
No	50	10.4	108	15.6	
Yes	432	89.6	584	84.4	100.0

*p<0.05, **p<0.01, ***p<0.001; ^{a)}Under 18 years old, ^{b)}Current tobacco use, ^{c)}No current tobacco use

563

564

Table 4 Odds ratio (OR) and 95% confidence interval (CI) of current tobacco use among study participants (N=1,174)

Characteristics	Use ^{a)}	No use ^{b)}	Unadjusted		Adjusted ^{c)}	
	(N=131)	(N=1043)	OR	95% CI	OR	95% CI
Age						
≤ 14 years	14 (10.7)	488 (46.8)	1	Reference	1	Reference
> 14 years	117 (89.3)	555 (53.2)	7.35	(4.17-12.96)***	9.81	(4.54-21.19)***
Gender						
Female	9 (6.9)	683 (65.5)	1	Reference	1	Reference
Male	122 (93.1)	360 (34.5)	25.72	(12.91-51.22)***	28.06	(13.29-59.25)***
Grade						
10	56 (42.7)	738 (70.8)	1	Reference	1	Reference
11	75 (57.3)	305 (29.2)	3.24	(2.24-4.70)***	1.52	(0.85-2.73)
Parent smoking						
No	128 (97.7)	987 (94.6)	1	Reference	1	Reference
Yes	3 (2.3)	56 (5.4)	0.41	(0.13-1.34)	0.31	(0.08-1.29)
Sibling smoking						
No	118 (90.1)	961 (92.1)	1	Reference	1	Reference
Yes	13 (9.9)	82 (7.9)	1.29	(0.70-2.39)	1.06	(0.47-2.37)
Peer smoking						
No	119 (90.8)	959 (91.9)	1	Reference	1	Reference
Yes	12 (9.2)	84 (8.1)	1.15	(0.61-2.17)	0.41	(0.17-0.95)

Exposure to any kind of TAPS ^{a)}

No	89 (6.1)	99 (9.5)	1	Reference	1	Reference
Yes	123 (93.9)	944 (90.5)	1.61	(0.77-3.40)	6.59	(2.33-18.64)***

Had seen any smoked tobacco product for sale inside or within 100 feet of the school premises within the last 12 months

No	9 (6.9)	176 (16.9)	1	Reference	1	Reference
Yes	122 (93.1)	867 (83.1)	2.75	(1.37-5.52)**	4.17	(1.65-10.58)**

Had seen the sale or gifting of any smoked tobacco product to minors ^{a)} within the last 12 months

No	4 (3.1)	202 (19.4)	1	Reference	1	Reference
Yes	127 (96.9)	841 (80.6)	7.63	(2.79-20.88)***	6.46	(2.18-19.12)***

Had seen the sale or distribution of any smoked tobacco product by minors ^{a)} within the last 12 months

No	44 (33.6)	477 (45.7)	1	Reference	1	Reference
Yes	87 (66.4)	566 (54.3)	1.67	(1.14-2.44)**	2.42	(1.42-4.10)***

Had seen the sale of cigarettes singly or in packs less than 20 cigarettes within the last 12 months

No	3 (2.3)	155 (14.9)	1	Reference	1	Reference
Yes	128 (97.7)	888 (85.1)	7.45	(2.34-23.70)***	3.32	(0.93-11.85)

Ever received health education about tobacco use

No	70 (53.4)	393 (37.7)	1	Reference	1	Reference
Yes	61 (46.6)	650 (62.3)	0.53	(0.37-0.76)***	0.45	(0.27-0.78)**

Perception of tobacco use

≤ 7	80 (61.1)	187 (17.9)	1	Reference	1	Reference
>7	51 (38.9)	856 (82.1)	0.14	(0.10-0.21)***	0.17	(0.10-0.30)***

* $p < 0.05$, ** $p < 0.01$, *** $p < 0.001$; ^{a)}Current tobacco use, ^{b)}No current tobacco use, ^{c)}Adjusted for age, gender, grade, parent smoking, sibling smoking, peer smoking, ever received health education about tobacco use, and perception of tobacco use, ^{d)}Under 18 years old

565

566 **STROBE 2007 (v4) Statement—Checklist of items that should be included in reports of**
567 ***cross-sectional studies***
568

Section/Topic	Item #	Recommendation	Reported on page #
Title and abstract	1	(a) Indicate the study’s design with a commonly used term in the title or the abstract	1-2
		(b) Provide in the abstract an informative and balanced summary of what was done and what was found	2-3
Introduction			
Background/rationale	2	Explain the scientific background and rationale for the investigation being reported	5-9
Objectives	3	State specific objectives, including any prespecified hypotheses	8-9
Methods			
Study design	4	Present key elements of study design early in the paper	9
Setting	5	Describe the setting, locations, and relevant dates, including periods of recruitment, exposure, follow-up, and data collection	9-10
Participants	6	(a) Give the eligibility criteria, and the sources and methods of selection of participants	9
Variables	7	Clearly define all outcomes, exposures, predictors, potential confounders, and effect modifiers. Give diagnostic criteria, if applicable	10-11

Data sources/ measurement	8*	For each variable of interest, give sources of data and details of methods of assessment (measurement). Describe comparability of assessment methods if there is more than one group	9-11
Bias	9	Describe any efforts to address potential sources of bias	21
Study size	10	Explain how the study size was arrived at	9
Quantitative variables	11	Explain how quantitative variables were handled in the analyses. If applicable, describe which groupings were chosen and why	10-11
Statistical methods	12	(a) Describe all statistical methods, including those used to control for confounding	11
		(b) Describe any methods used to examine subgroups and interactions	-
		(c) Explain how missing data were addressed	9
		(d) If applicable, describe analytical methods taking account of sampling strategy	-
		(e) Describe any sensitivity analyses	-
Results			
Participants	13*	(a) Report numbers of individuals at each stage of study—eg numbers potentially eligible, examined for eligibility, confirmed eligible, included in the study, completing follow-up, and analysed	9,11-12

		(b) Give reasons for non-participation at each stage	-
		(c) Consider use of a flow diagram	-
Descriptive data	14*	(a) Give characteristics of study participants (eg demographic, clinical, social) and information on exposures and potential confounders	11-13 Table 1 to 3
		(b) Indicate number of participants with missing data for each variable of interest	9
Outcome data	15*	Report numbers of outcome events or summary measures	11-14 Table 1 Table 2 Table 3
Main results	16	(a) Give unadjusted estimates and, if applicable, confounder-adjusted estimates and their precision (eg, 95% confidence interval). Make clear which confounders were adjusted for and why they were included	13-14 Table 4
		(b) Report category boundaries when continuous variables were categorized	-
		(c) If relevant, consider translating estimates of relative risk into absolute risk for a meaningful time period	-
Other analyses	17	Report other analyses done—eg analyses of subgroups and interactions, and sensitivity analyses	-

Discussion			
Key results	18	Summarise key results with reference to study objectives	14
Limitations	19	Discuss limitations of the study, taking into account sources of potential bias or imprecision. Discuss both direction and magnitude of any potential bias	21
Interpretation	20	Give a cautious overall interpretation of results considering objectives, limitations, multiplicity of analyses, results from similar studies, and other relevant evidence	14-21
Generalisability	21	Discuss the generalisability (external validity) of the study results	21
Other information			
Funding	22	Give the source of funding and the role of the funders for the present study and, if applicable, for the original study on which the present article is based	23

569

570 *Give information separately for cases and controls in case-control studies and, if applicable,
571 for exposed and unexposed groups in cohort and cross-sectional studies.

572

573 **Note:** An Explanation and Elaboration article discusses each checklist item and gives
574 methodological background and published examples of transparent reporting. The STROBE
575 checklist is best used in conjunction with this article (freely available on the Web sites of
576 PLoS Medicine at <http://www.plosmedicine.org/>, Annals of Internal Medicine at
577 <http://www.annals.org/>, and Epidemiology at <http://www.epidem.com/>). Information on the
578 STROBE Initiative is available at www.strobe-statement.org.

579

580