

# CELL SURFACE GLYCOPROTEINS: BIOCHEMICAL, IMMUNOLOGICAL AND MOLECULAR BIOLOGICAL STUDIES

TAKASHI MURAMATSU

*Department of Biochemistry, Nagoya University School of Medicine, Nagoya, Japan*

## ABSTRACT

This article briefly summarizes the author's contribution to the study of cell surface glycoproteins. At first, much effort was devoted to developing analytical methods, especially exoglycosidases and endoglycosidases. An endo- $\beta$ -*N*-acetylglucosaminidase found in *Streptococcus pneumoniae* was the first example of an endoglycosidase acting on glycoproteins. A combination of radioactive labeling, glycosidase digestion and lectin affinity chromatography enabled the characterization of carbohydrate moieties of cell surface glycoproteins derived from cultured cells. Application of the method to teratocarcinoma stem cells and preimplantation mouse embryos led to the discovery of marked changes in the carbohydrate moieties of cell surface glycoproteins during embryogenesis. Combining biochemical and immunohistochemical methods, an overall picture was obtained for carbohydrate changes in early embryogenesis of the mouse, and molecular biological approaches have been adopted to determine their biological significance. Furthermore, some core proteins carrying developmentally regulated carbohydrate markers were found and characterized by molecular cloning. One example of such a protein is embigin, which enhances integrin-mediated cell-substratum adhesion and established the existence of a new group in the immunoglobulin superfamily. Carbohydrate immunochemical markers useful in the analysis of mouse embryogenesis were also found to be effective in the classification of human carcinomas with respect to metastatic potential.

## INTRODUCTION

Molecular interactions at the cell surface play important roles in the regulation of cellular activities such as growth, differentiation, migration and adhesion.<sup>1)</sup> The author has been engaged in studies on cell surface molecules, namely cell surface glycoproteins and a novel heparin binding growth factor, midkine,<sup>2-25)</sup> the latter of which has recently been reviewed.<sup>2,3)</sup> Taking this opportunity, I wish to summarize our contribution to a research area, namely cell surface glycoproteins, with some personal reflections.

## EXOGLYCOSIDASES

As a graduate student, I joined the laboratory of Prof. F. Egami, Faculty of Science, Tokyo University. Prof. Egami identified ribonuclease T<sub>1</sub>, which is useful in structural studies of ribonucleic acid. He proposed that I study carbohydrases of a marine gastropod, *Charonia lampas* (Japanese name: *bōshubora* or *horagai*), considering that studies of complex carbohydrates would become important and that the identification and characterization of enzymes capable of their degradation would be highly desirable.

Among the wide variety of complex carbohydrates, I chose to study glycoproteins, since little was known about these molecules and enzymes acting on the carbohydrate moieties. The apparent specificity of the glycosylation products promised both interesting and important results. I prepared extracts from the hepatopancreas of *C. lampas*, and examined enzymes acting on glycopeptides isolated from both ovalbumin and ovomucoid. Enzymes releasing mannose, N-acetylglucosamine and galactose were found, but no oligosaccharides were detected even for short reaction times, suggesting that the sugars were released by the step-wise activity of exoglycosidases.<sup>26)</sup> The enzyme releasing mannose was found to be novel, and was considered to be an  $\alpha$ -mannosidase. Thus, attempts were made to purify it using p-nitrophenyl  $\alpha$ -mannoside as a substrate. The purified enzyme indeed released mannose from glycopeptides.<sup>27)</sup> However, an  $\alpha$ -mannosidase was also purified from jack bean meal, and was shown to act on glycoprotein carbohydrates. The characterization of the jack bean enzyme was published slightly earlier than that of the *C. lampas* enzyme.

The importance of enzymes acting on glycoprotein carbohydrate moieties became apparent to several investigators at the time. I, together with co-workers, systematically purified various exoglycosidases from *C. lampas*, and also from another marine gastropod, *Turbo cornutus* (Japanese name: *sazae*) (Table 1).<sup>26-33)</sup> Among these,  $\beta$ -mannosidase and  $\beta$ -xylosidase were the first to be purified. These enzymes act on glycoprotein carbohydrates, and have since proved useful in structural and biological studies of glycoproteins as well as bacterial lipopolysaccharides. (Table 1). They became commercially available from Seikagaku Kogyo Co. During my time in Prof. Egami's laboratory, I also studied both glycoproteins from the jelly coat of starfish eggs<sup>34)</sup> and sulfated glycopeptides from bovine brain,<sup>35)</sup> the former of which were later shown to participate in fertilization by Prof. M. Hoshi, Tokyo Institute of Technology.

Table 1. Exoglycosidase Purified from Marine Gastropods

Enzyme	Substrates including applied subjects
$\alpha$ -Mannosidase	Ovalbumin glycopeptide, Stem bromelain glycopeptide, Japanese encephalitis virus receptor, Bacterial lipopolysaccharide
$\beta$ -Mannosidase	Bacterial lipopolysaccharide
$\alpha$ -L-Fucosidase	Blood group substances, Milk oligosaccharides
$\beta$ -Xylosidase	Stem bromelain glycopeptide, Xylosyl serine linkage in proteoglycan
$\beta$ -N-Acetylhexosaminidase	Ovomucoid
$\alpha$ -N-Acetylgalactosaminidase	—

## MAJOR HISTOCOMPATIBILITY ANTIGEN

After receiving my Ph. D., I wished to study the biological roles of cell surface glycoproteins, and joined the laboratory of Dr. Stanley G. Nathenson, Albert Einstein College of Medicine, as a postdoctoral fellow. He published a paper on the purification of H-2 antigen (major histocompatibility antigen of the mouse), and demonstrated that it is a glycoprotein. I felt that the H-2 antigenic determinant may be carbohydrate as is the case for the ABH blood group antigen.

The H-2 antigen could be obtained in only very small amounts, making biochemical studies difficult. However, Einstein College at that time was a major world center of cell biology, and new techniques including SDS polyacrylamide gel electrophoresis were developed there. In

## CELL SURFACE GLYCOPROTEINS

Nathenson's laboratory, radioactively labeled H-2 antigen was prepared by culturing H-2 bearing tumor cells in the presence of radioactive amino acids. Following this method, I isolated carbohydrate-labeled H-2 antigen, which could be used to characterize the carbohydrate portion.<sup>36-39)</sup> A central question was whether carbohydrate is the antigenic determinant. While an article in *Nature* reported that this was indeed the case, all my data strongly suggested otherwise, and the *Nature* paper was subsequently shown to be mistaken. Although results obtained were not dramatic, I learned how powerful the application of radiolabeling methods to glycoprotein research could be. Using the technique, we characterized the thymus leukemia (TL) antigen.<sup>40)</sup>

## ENDOGLYCOSIDASES AND FIBROBLAST OLIGOSACCHARIDES

Until 1970, all enzymes acting on glycoprotein carbohydrates were thought to be exoglycosidases. When I was analyzing radioactively labeled H-2 glycopeptides using an enzyme preparation from *Streptococcus pneumoniae*, I found an unusual product, best explained by cleavage by an endoglycosidase. Changing the substrate to carbohydrate-labeled IgG glycopeptide, I indeed showed the presence of an endoglycosidase acting on the glycoprotein.<sup>41)</sup> The enzyme appeared to be an endo- $\beta$ -*N*-acetylglucosaminidase.

After my appointment at the laboratory of Prof. A. Kobata, Kobe University School of Medicine, I began systematic studies of the endoglycosidase. The ovalbumin glycopeptide (Man)<sub>5</sub>(GlcNAc)<sub>2</sub>An was [<sup>14</sup>C]-*N*-acetylated and used to assay the enzyme; Asn [<sup>14</sup>C]-acetyl GlcNAc released was quantified. The enzyme was purified and its specificity determined.<sup>42-46)</sup> It was shown to cleave di-*N*-acetylchitobiose structure, and have strict specificity for the oligomannosyl cores of glycopeptides. This endoglycosidase, named endo- $\beta$ -*N*-acetylglucosaminidase D (Endo D), acts on complex-type carbohydrate chains after releasing external side chain sugars. After the discovery of Endo D, another endo- $\beta$ -*N*-acetylglucosaminidase was found in culture filtrates of *Streptomyces* by A. Tarentino. We found Endo D and the *Streptomyces* enzyme (endo- $\beta$ -*N*-acetylglucosaminidase H: Endo H) to have complementary specificities (Fig. 1).<sup>46)</sup> The exact specificity of Endo D was elucidated by structural studies of the substrates.<sup>45,47-49)</sup> We also studied endo- $\beta$ -*N*-acetylglucosaminidases from rat liver,<sup>50)</sup> *Clostridium perfringens*<sup>51)</sup> and fig.<sup>52)</sup> Endo- $\beta$ -*N*-acetylglucosaminidases have recently been reviewed.<sup>53)</sup>




Fig. 1. Different specificities of Endo D and Endo H. The upper substrate was hydrolyzed by Endo D, but not by Endo H; the lower substrate was hydrolyzed by Endo H, but not by Endo D.

The endoglycosidases have been used in structural studies of ovalbumin glycopeptides<sup>47)</sup> and IgG glycopeptides.<sup>49)</sup> Furthermore, combination of exoglycosidase and Endo D proved useful in the removal of carbohydrates from IgG.<sup>54)</sup> This enzymatic digestion procedure demonstrated the importance of carbohydrates in Fc region functions such as binding to Fc receptors. The restricted and complementary specificities of Endo D and Endo H and related enzymes were used by three groups in the USA and, consequently, they identified and characterized a processing pathway for asparagine-linked oligosaccharides.<sup>53)</sup>

Radiolabeled glycopeptides were also used to study the specificity of concanavalin A columns. We found that two  $\alpha$ -mannosyl residues, which may be substituted only at C-2, are required for binding to the column.<sup>55)</sup> We also extended studies on exoglycosidases using radiolabeled substrates.<sup>56-58)</sup>

Soon after the discovery of Endo D, I felt that it would be useful for the characterization of radioactively labeled glycopeptides from the cell surface of cultured fibroblast cells.<sup>59)</sup> Using the complementary specificities of Endo D and Endo H, we found that the level of high mannose oligosaccharides is elevated in growing cells as compared with nongrowing cells.<sup>60,61)</sup> Endo D and Endo H digestion was useful in determining the overall structure of asparagine-linked cell surface oligosaccharides.<sup>61-64)</sup> Furthermore, in addition to the enzymatic method, concanavalin A affinity chromatography was employed. Consequently, we identified increased branching in cell surface oligosaccharides from virally-transformed cells.<sup>65)</sup>

## TERATOCARCINOMAS AND EARLY MOUSE EMBRYOS

After studying endo- $\beta$ -*N*-acetylglucosaminidases and fibroblast glycopeptides, I wished to study more directly the function of cell surface carbohydrates, and selected early mouse embryogenesis as the experimental system. As a Franco-Japanese exchange scholar, I joined the laboratory of Prof. F. Jacob, Pasteur Institute. At that time, Prof. Jacob concentrated on the mouse teratocarcinoma system to study the molecular mechanisms of early embryogenesis. Since nothing was known about carbohydrates of cell surface glycoproteins in early embryonic cells, I analyzed [<sup>3</sup>H]-fucose-labeled glycopeptides from plasma membranes and whole cells of teratocarcinoma stem cells (EC cells), which resemble the multipotent cells of early embryos.<sup>66,67)</sup> Unexpectedly, high levels of the label were eluted in the excluded volume of Sephadex G-50 columns. This radioactivity progressively decreased when EC cells differentiated *in vitro*. Preimplantation and early postimplantation mouse embryos also synthesized large fucosyl glycopeptides, while during the midgestational period, the synthetic activity became scarcely detectable.<sup>66,68)</sup> Various biochemical experiments excluded the possibility that the large glycopeptides were mucopolysaccharides, mucin-type glycopeptides or glycolipids.<sup>67)</sup> Thus, we concluded that there exists a class of unusual glycoproteins in early embryonic cells, and that the carbohydrate profile changes markedly during embryogenesis.<sup>66-72)</sup> The large carbohydrates were found to carry certain cell surface markers of EC cells, such as *Lotus tetragonolobus* agglutinin binding sites, and glycosidase digestion showed them to be poly-*N*-acetylglucosamines.<sup>69)</sup>

Two years after returning from France, I was appointed as a professor of biochemistry, Kagoshima University Faculty of Medicine, and opened a new laboratory. We continued studies on the large carbohydrate from EC cells and isolated and biochemically characterized it.<sup>73-81)</sup> In particular, we showed it to be highly branched, and to have a molecular weight of around 10,000 or more. On the basis of its complexity and the markers it carried, this large glycan was different from other poly-*N*-acetylglucosamines so far identified, and we named it embryoglycan.<sup>75)</sup> The protein-carbohydrate linkage of embryoglycan is of asparagine-linked one.<sup>76,80)</sup>

## CELL SURFACE GLYCOPROTEINS

Embryoglycan has Le<sup>x</sup> structure, epitope of SSEA-1, which is the most well-known cell surface marker of EC cells.<sup>80)</sup> Furthermore, the major carrier of SSEA-1 in EC cells was shown to be high molecular weight glycoproteins with embryoglycan.<sup>82)</sup>

In addition to biochemical analysis, we employed immunohistochemical methods to analyze carbohydrate changes during development.<sup>83-95)</sup> For example, an *N*-acetylgalactosamine-recognizing lectin, *Dolichos biflorus* agglutinin (DBA), specifically reacted with extraembryonic endoderm in early postimplantation embryos,<sup>86)</sup> and in the cells derived from pluripotent EC cells.<sup>91)</sup>

Summarizing all these results and citing works performed in other laboratories, I presented an overall picture of carbohydrate changes during early mouse embryogenesis.<sup>96-98)</sup>

The epitope structure of the binding site of DBA was determined as shown in Table 2.<sup>99)</sup> DBA also specifically reacts with thymocytes in early developmental stages,<sup>83)</sup> some leukemia cells<sup>100)</sup> and restricted tissues in adult mice.<sup>84)</sup> EC cells whose direction of differentiation is restricted to extraembryonic endoderm cells are also reactive to DBA.<sup>101)</sup>

Table 2. Carbohydrate Markers Present on Poly-*N*-acetylactosamine (embryoglycan) and Useful for Analysis of Early Mouse Embryogenesis

Name	Determinant
Antigens	
SSEA-1 (Le <sup>x</sup> )	Galβ 1→4(Fucα 1→3)GlcNAc
I	Galβ1→4GlcNAcβ1
	$\searrow$ 6 Galβ1→4GlcNAcβ1→3Gal $\nearrow$ 3 Galβ1→4GlcNAcβ1
i	Galβ1→4GlcNAcβ1→3Galβ1→4GlcNAcβ1→3Gal
Lectin receptors	
LTA (FBP)	Galβ1→4(Fucα1→3)GlcNAc
PNA	Galβ1→3GalNAc
DBA	GalNAcβ1→4(NeuAcα2→3)Galβ1→4GlcNAcβ1→3Gal
GSI-B <sub>4</sub>	Galα1→3Gal

The basis of the carbohydrate change is alteration in glycosyltransferase activity. Indeed, we verified that the reduction in SSEA-1 antigenicity after differentiation of EC cells was accompanied by a concomitant reduction in α1-3 fucosyltransferase activity.<sup>102)</sup> The key fucosyltransferase was purified and characterized.<sup>103)</sup>

Embryoglycan carries numerous markers found in early embryonic cells (Table 2).<sup>82,95-98,104-107)</sup> In addition, sera from patients with germ cell tumors frequently have precipitating antibodies which recognize an epitope on embryoglycan.<sup>108-110)</sup> Using endo-β-galactosidase C, a novel endo-β-galactosidase identified in our laboratory, which acts on Galα1-3Gal β1-4 GlcNAc sequence releasing Gal α1-3 Gal disaccharide,<sup>111-112)</sup> the epitope recognized by the patients' sera was shown to be Gal α1-3Gal.<sup>113)</sup> Endo-β-galactosidase C was also useful in correcting the structure of glycopeptides from Ehrlich carcinoma cells.<sup>114)</sup>

To investigate the function of carbohydrates on EC cells, we prepared anti-carbohydrate monoclonal antibodies by using EC cells as the immunogen. A monoclonal antibody thus

obtained, 4C9, inhibited cell-substratum adhesion of EC cells.<sup>115)</sup> The epitope recognized by the 4C9 antibody is Le<sup>X</sup> structure. The distribution of 4C9 antigen is similar to SSEA-1, while 4C9 is more specific for cells in lineages starting from totipotent embryonic cells leading to primordial germ cells.<sup>116)</sup> After several control experiments, we proposed that Le<sup>X</sup> structure enhances cell-substratum adhesion. This proposal has been supported by a recent finding in our laboratory; transfection with  $\alpha$ 1-3 fucosyltransferase that forms Le<sup>X</sup> structure, resulted in increased cell-substratum adhesion of transfected L cells (Sudou, A. et al., unpublished).

We also searched for lectins in teratocarcinoma cells to find endogeneous sugar-recognizing proteins. A lectin was found in teratocarcinoma OTT6050, but identified as laminin.<sup>117)</sup> The finding of lectin activity in laminin is, however, important in considering the function of laminin.

A candidate for a carbohydrate-recognizing protein on the surface of EC cells is  $\beta$ -galactosyltransferase. We prepared a monoclonal antibody to the enzyme and verified its cell surface location.<sup>118)</sup>

To obtain a more definitive answer concerning the function of carbohydrates, we decided to use transgenic techniques. We produced transgenic mice carrying  $\alpha$  1-3 galactosyltransferase cDNA, and these mice became the first example of transgenic animals expressing a specific carbohydrate structure ectopically.<sup>119)</sup> They tended to show some abnormalities, such as increased protein level in the urine. Undoubtedly, gene targeting experiments will be highly informative, and work along this line is currently in progress.

## CORE PROTEINS

We were also interested in proteins that carry developmentally regulated carbohydrate chains. To identify these, glycoproteins were isolated from teratocarcinoma cells by affinity chromatography through lectin-agarose columns. Rabbits were immunized against the glycoproteins, and affinity-purified antibodies were used to immunochemically characterize glycoproteins.<sup>120,121)</sup> Subsequently, monoclonal antibody technique was also used.<sup>122)</sup> Then, we used the affinity purified antibodies to screen a  $\lambda$ gt 11 expression library. When DBA lectin was used, a core protein named embigin was identified, the protein moiety of which had a molecular weight of 30 kDa, while that of the whole glycoprotein was 70 to 90 kDa.<sup>123)</sup> Embigin has a transmembrane domain, and the extracellular portion contains two immunoglobulin-like domains. During mouse embryogenesis, embigin is preferentially expressed before 10 days of gestation, and *in situ* hybridization analysis has identified the preferential site of this expression.<sup>124)</sup> L cells lack embigin, and those transfected with embigin cDNA exhibited increased cell-substratum adhesion activity.<sup>125)</sup> This increased adhesion was inhibited by anti-integrin antibody. Thus, embigin was concluded to have a role in enhancing integrin-mediated cell-substratum adhesion.

Using antibodies raised against glycoproteins binding to *Lotus tetragonolobus* agglutinin (LTA),<sup>126)</sup> we also cloned a core protein named basigin.<sup>127,128)</sup> Basigin shows strong homology to embigin throughout most of the molecule, indicating that embigin is a prototype of a new class of molecules in the Ig superfamily. The molecular weight of the protein portion of basigin is 27 kDa, while that of the glycosylated molecule is 44 to 66 kDa. The immunoglobulin-like domain close to the transmembrane region is unique, since it has homology to both Ig V domain and MHC class II $\beta$ -chain, which has a C domain (Fig. 2). Basigin may be related to the primordial form of the immunoglobulin superfamily. Basigin was also found independently by other groups and given different names such as gp42 and HT7. Although basigin is broadly distributed in various embryonic and adult tissues, it may still be used as a marker of brain blood vessels; basigin was reported to be absent in neurons and systemic blood vessels. Attempts are currently

## CELL SURFACE GLYCOPROTEINS




Fig. 2. Relationship between members of the Ig superfamily. Cited from Ref. 128.

underway to determine the function of basigin by gene targeting.

Using antibodies to DBA-binding glycoproteins from teratocarcinomas, we also found two new proteins by cDNA cloning. One is heparin-binding protein-44 (HBP-44),<sup>129-130</sup> which forms a complex with brushin<sup>120,131</sup> (also called GP-330 or Heymann nephritis antigen), a calcium sensor common to the kidney and extraembryonic endoderm. The other is reticulocalbin, a calcium-binding protein which is located in the endoplasmic reticulum and has 4 EF-hand.<sup>132</sup> Antibody screening was also applied to identify a new core protein in peanut agglutinin (PNA)-binding glycoproteins from human gastric carcinomas.<sup>133-135</sup> MGC-24 (*multi-glycosylated core protein-24*) thus found, is a hydrophilic, cysteine-rich protein which carries numerous Gal $\beta$ 1-3 GalNAc oligosaccharides.<sup>135</sup>

## CARBOHYDRATES AS METASTATIC AND PROGNOSTIC MARKERS

At Kagoshima University Faculty of Medicine, we embarked upon numerous collaborative studies with clinical departments; the subjects were glycoproteins such as those in photoreceptor cells and human carcinomas.<sup>136-151</sup> The most notable finding produced by these collaborative studies was that the carbohydrate markers we used to analyse mouse embryogenesis were also useful in analyzing the biological properties of human carcinomas. Thus, lung carcinomas with [DBA(+), 4C9(-)] carbohydrate profile are less metastatic, and patient prognoses are better in comparison with carcinomas with different carbohydrate profiles.<sup>146</sup>

In the case of transient cell carcinoma of the urinary bladder, primary tumors that strongly express *Lotus tetragonolobus* agglutinin binding sites (LTA), showed a tendency to be more metastatic than those expressing them only weakly or not at all.<sup>147</sup> Also, expression of LTA-related markers, namely Le<sup>X</sup> and sialyl Le<sup>X</sup>, correlated with metastatic potential of the tumors.<sup>145,148</sup> The latter is interesting from the point of view that selectins, which are carbohydrate receptors involved in adhesion of leukocytes and which recognize sialyl Le<sup>X</sup>, may be involved in metastasis of human carcinomas. These metastasis-associated carbohydrate markers are present on a 60 kDa glycoprotein.<sup>149</sup> The influence of carbohydrate profiles on metastatic potential and prognosis of human carcinomas has been recently reviewed by the author.<sup>152</sup>

## CONCLUDING REMARKS

In the three decades from when I began studying glycoproteins, this research area has undergone many major changes. Functions of cell surface carbohydrates have been mentioned in diverse biological systems,<sup>153)</sup> carbohydrate structures have been elucidated for many glycoproteins, glycosyltransferases forming the chains have been cloned, and carbohydrate receptors called selections have been identified and cloned. Glycobiology is now the key word for this busy area. Since gene targeting has come to the center stage, the biological functions of the carbohydrate chains will be definitively determined. I wish to contribute to glycobiology also at this important stage, from the Department of Biochemistry, Nagoya University School of Medicine, to which I have recently moved.

## REFERENCES

- 1) Muramatsu, T.: Cell Surface and Differentiation. Chapman and Hall, London (1990).
- 2) Muramatsu, T.: Midkine, the product of a retinoic acid responsive gene, and pleiotrophin constitute a new protein family regulating growth and differentiation. *Int. J. Dev. Biol.*, 37, 183–188 (1993).
- 3) Muramatsu, T.: The midkine family of growth differentiation factors. *Dev. growth Differ*, 36, 1–8 (1994).
- 4) Kadomatsu, K., Tomomura, M. and Muramatsu, T.: cDNA cloning and sequencing of a new gene intensely expressed in early differentiation stages of embryonal carcinoma cells and in mid-gestation period of mouse embryogenesis. *Biochem. Biophys. Res. Commun.*, 151, 1312–1318 (1988).
- 5) Kadomatsu, K., Huang, R.-P., Suganuma, T. et al.: A retinoic acid responsive gene MK found in the teratocarcinoma system is expressed in spatially and temporally controlled manner during mouse embryogenesis. *J. Cell Biol.*, 110, 607–616 (1990).
- 6) Matsubara, S., Tomomura, M., Kadomatsu, K. et al.: Structure of a retinoic acid-responsive gene, MK, which is transiently activated during the differentiation of embryonal carcinoma cells and the midgestation period of mouse embryogenesis. *J. Biol. Chem.*, 265, 9441–9443 (1990).
- 7) Tomomura, M., Kadomatsu, K., Matsubara, S. et al.: A retinoic acid-responsive gene, MK, found in the teratocarcinoma system. Heterogeneity of the transcript and the nature of the translation product. *J. Biol. Chem.*, 265, 10765–10770 (1990).
- 8) Tomomura, M., Kadomatsu, K., Nakamoto, M. et al.: A retinoic acid responsive gene, MK, produces a secreted protein with heparin binding activity. *Biochem. Biophys. Res. Commun.*, 171, 603–609 (1990).
- 9) Huang, R.-P., Muramatsu, H. and Muramatsu, T.: Effects of different conditions of retinoic acid treatment on expression of MK gene, which is transiently activated during differentiation of embryonal carcinoma cells. *Dev. Growth Differ.*, 32, 189–195 (1990).
- 10) Muramatsu, H. and Muramatsu, T.: Purification of recombinant midkine and examination of its biological activities: functional comparison of new heparin binding factors. *Biochem. Biophys. Res. Commun.*, 177, 652–658 (1991).
- 11) Tsutsui, J., Uehara, K., Kadomatsu, K. et al.: A new family of heparin-binding factors: strong conservation of midkine (MK) sequences between the human and the mouse. *Biochem. Biophys. Res. Commun.*, 176, 792–797 (1991).
- 12) Simon-Chazottes, D., Matsubara, S., Miyauchi, T. et al.: Chromosomal localization of two cell-surface associated molecules of potential importance in development: midkine (Mdk) and basigin (Bsg). *Mammalian Genome*, 2, 269–271 (1992).
- 13) Uehara, K., Matsubara, S., Kadomatsu, K. et al.: Genomic structure of human midkine (MK), a retinoic acid-responsive growth/differentiation factor. *J. Biochem.*, 111, 563–567 (1992).
- 14) Nakamoto, M., Matsubara, S., Miyauchi, T. et al.: A new family of heparin binding growth/differentiation factors: differential expression of the midkine (MK) and HB-GAM genes during mouse development. *J. Biochem.*, 112, 340–349 (1992).
- 15) Maruta, H., Bartlett, P.F., Nurcombe, V. et al.: Midkine (MK), a retinoic acid (RA)-inducible gene product, produced in *E. coli* acts on neuronal and HL60 leukemia cells. *Growth Factors*, 8, 119–134 (1993).
- 16) Tsutsui, J., Kadomatsu, K., Matsubara, S. et al.: A new family of heparin-binding growth differentiation factors: increased midkine expression in Wilms' tumor and other human carcinomas. *Cancer Res.*, 53, 1281–1285 (1993).



## CELL SURFACE GLYCOPROTEINS

- 17) Fabri, L., Maruta, H., Muramatsu, H. et al.: Structural characterization of native and recombinant forms of the neurotrophic cytokine MK. *J. Chromatog.*, 646, 213–226 (1993).
- 18) Kitamura, M., Shirasawa, T., Mitarai, T. et al.: A retinoid responsive cytokine gene, MK, is preferentially expressed in the proximal tubules of the kidney and human tumor cell lines. *Am J. Pathol.*, 142, 425–431 (1993).
- 19) Kaname, T., Kuwano, A., Murano, I. et al.: Midkine gene (MDK), a gene for prenatal differentiation and neuroregulation, maps to band 11p11.2 by fluorescence in situ hybridization. *Genomics*, 17, 514–515 (1993).
- 20) Michikawa, M., Kikuchi, S., Muramatsu, H. et al.: Retinoic acid responsive gene product, midkine (MK), has neurotrophic functions for mouse spinal cord and dorsal root ganglion neurons in culture. *J. Neurosci. Res.*, 35, 530–539 (1993).
- 21) Yasuhara, O., Muramatsu, H., Kim, S.U. et al.: Midkine, a novel neurotrophic factor, is present in senile plaques of Alzheimer disease. *Biochem. Biophys. Res. Commun.*, 192, 246–251 (1993).
- 22) Michikawa, M., Xu, R.Y., Muramatsu, H. et al.: Midkine is a mediator of retinoic acid induced neuronal differentiation of embryonal carcinoma cells. *Biochem. Biophys. Res. Commun.*, 192, 1312–1318 (1993).
- 23) Satoh, J., Muramatsu, H., Moretto, G. et al.: Midkine that promotes survival of human neurons is produced by fetal human astrocytes in culture. *Dev. Brain Res.*, 75, 201–205 (1993).
- 24) Muramatsu, H., Shirahama, H., Yonezawa, S. et al.: Midkine (MK), a retinoic acid-inducible growth/differentiation factor: immunochemical evidence for the function and distribution. *Dev. Biol.*, 159, 392–402 (1993).
- 25) Kikuchi, S., Muramatsu, H., Muramatsu, T. et al.: Midkine, a novel neurotrophic factor, promotes survival of mesencephalic neurons in culture. *Neurosci. Lett.*, 160, 9–12 (1993).
- 26) Muramatsu, T. and Egami, F.: Hydrolysis of ovalbumin glycopeptide and ovomucoid glycopeptide by an enzyme preparation from the liver of *Charonia lampas*. *Jap. J. Exp. Med.*, 35, 171–179 (1965).
- 27) Muramatsu, T.: Purification of  $\alpha$ -D-mannosidase and  $\beta$ -D-mannosidase from marine gastropods. *Arch. Biochem. Biophys.*, 115, 427–429 (1966).
- 28) Muramatsu, T.: Purification and properties of  $\alpha$ -mannosidase from the liver of *Charonia lampas*. *J. Biochem.*, 62, 487–491 (1967).
- 29) Muramatsu, T. and Egami, F.:  $\alpha$ -mannosidase and  $\beta$ -mannosidase from the liver of *Turbo cornutus*: purification, properties and application in carbohydrate research. *J. Biochem.*, 62, 700–709 (1968).
- 30) Fukuda, M., Muramatsu, T., Egami, F. et al.: Purification of  $\beta$ -xylosidase and its action on O- $\beta$ -xylosyl L-serine and stem bromelain glycopeptide. *Biochim. Biophys. Acta.*, 159, 215–216 (1968).
- 31) Muramatsu, T.: N-Acetylhexosaminidases from a gastropod, *Turbo cornutus*. *J. Biochem.*, 64, 521–531 (1968).
- 32) Fukuda, M., Muramatsu, T. and Egami, F.:  $\beta$ -xylosidase from the liver of *Charonia lampas*. I. Purification, properties and application in carbohydrate research. *J. Biochem.*, 65, 191–199 (1969).
- 33) Iijima, Y., Muramatsu, T. and Egami, F.: Purification of  $\alpha$ -L-fucosidase from the liver of a marine gastropod, *Turbo cornutus*, and its action on blood group substances. *Arch. Biochem. Biophys.*, 145, 50–54 (1971).
- 34) Muramatsu, T.: A sulphated mucopolysaccharide from the jelly coat of starfish eggs. *J. Biochem.*, 57, 223–225 (1965).
- 35) Arima, T., Muramatsu, T., Saigo, K. et al.: Sialomucopolysaccharides from bovine brain. *Jap. J. Exp. Med.*, 39, 301–309 (1969).
- 36) Muramatsu, T. and Nathenson, S.G.: Isolation of a chromatographically unique glycopeptide from murine histocompatibility-2 (H-2) membrane alloantigens labelled with H3-fucose or H3-glucosamine. *Biochem. Biophys. Res. Commun.*, 38, 1–8 (1970).
- 37) Muramatsu, T. and Nathenson, S.G.: Studies on the carbohydrate portion of membrane-located H-2 alloantigens. *Biochemistry*, 9, 4875–4883 (1970).
- 38) Muramatsu, T. and Nathenson, S.G.: Comparison of the carbohydrate portion of membrane H-2 alloantigens isolated from spleen cells and tumor cells. *Biochim. Biophys. Acta*, 241, 195–199 (1971).
- 39) Nathenson, S.G. and Muramatsu, T.: Properties of the carbohydrate portion of mouse H-2 alloantigen glycoproteins, Glycoproteins from Red Cells and Plasma. Jamieson, G.A. Ed., Lippincott, Philadelphia, pp245–262 (1971).
- 40) Muramatsu, T., Nathenson, S.G., Boyse, E.A. et al.: Some biochemical properties of thymus leukemia antigens solubilized from cell membranes by papain digestion. *J. Exp. Med.*, 137, 1256–1262 (1973).
- 41) Muramatsu, T.: Demonstration of an endo-glycosidase acting on a glycoprotein. *J. Biol. Chem.*, 246, 5535–5537 (1971).

- 42) Koide, N. and Muramatsu, T.: Endo- $\beta$ -*N*-acetylglucosaminidase acting on carbohydrate moieties of glycoproteins: purification and properties of the enzyme from *Diplococcus pneumoniae*. J. Biol. Chem., 249, 4897–4904 (1974).
- 43) Koide, N. and Muramatsu, T.: Specific inhibition of endo- $\beta$ -*N*-acetylglucosaminidase D by mannose and simple mannosides. Biochem. Biophys. Res. Commun., 66, 411–416 (1975).
- 44) Muramatsu, T.: Endo- $\beta$ -*N*-acetylglucosaminidase D from *Diplococcus pneumoniae*. Methods Enzymol., 50, 555–559 (1978).
- 45) Muramatsu, T., Koide, N. and Maeyama, K.: Further studies on endo- $\beta$ -*N*-acetylglucosaminidase D. J. Biochem., 83, 363–370 (1978).
- 46) Arakawa, M. and Muramatsu, T.: Endo- $\beta$ -*N*-acetylglucosaminidases acting on the carbohydrate moieties of glycoproteins: the differential specificities of the enzymes from *Streptomyces griseus* and *Diplococcus pneumoniae*. J. Biochem., 76, 307–314 (1974).
- 47) Tai, T., Yamashita, K., Ogata-Arakawa, M. et al.: Structure of two ovalbumin glycopeptides in relation to endo- $\beta$ -*N*-acetylglucosaminidase specificity. J. Biol. Chem., 250, 8569–8575 (1975).
- 48) Ito, S., Muramatsu, T. and Kobata, A.: Release of galactosyl oligosaccharides by endo- $\beta$ -*N*-acetylglucosaminidase D. Biochem. Biophys. Res. Commun., 63, 938–944 (1975).
- 49) Tai, T., Ito, S., Yamashita, K. et al.: Asparagine-linked oligosaccharide chains of IgG: a revised structure. Biochem. Biophys. Res. Commun., 65, 968–974 (1975).
- 50) Nishigaki, M., Muramatsu, T. and Kobata, A.: Endoglycosidases acting on carbohydrate moieties of glycoproteins: demonstration in mammalian tissue. Biochem. Biophys. Res. Commun., 59, 638–645 (1974).
- 51) Ito, S., Muramatsu, T. and Kobata, A.: Endo- $\beta$ -*N*-acetylglucosaminidases acting on carbohydrate moieties of glycoproteins: purification and properties of the two enzymes with different specificities from *Clostridium perfringens*. Arch. Biochem. Biophys., 171, 78–86 (1975).
- 52) Ogata-Arakawa, M., Muramatsu, T. and Kobata, A.: Partial purification and characterization of an endo- $\beta$ -*N*-acetylglucosaminidase from fig extract. J. Biochem., 82, 611–614 (1977).
- 53) Muramatsu, T.: Endo- $\beta$ -*N*-acetylglucosaminidases, CRC Handbook of Endoglycosidases and Glycoamidases. Takahashi, N. and Muramatsu, T. Eds., CRC Press, Boca Raton, Ann Arbor, London, pp13–40 (1992).
- 54) Koide, N., Nose, M. and Muramatsu, T.: Recognition of IgG by Fc receptor and complement: effects of glycosidase digestion. Biochem. Biophys. Res. Commun., 75, 838–843 (1977).
- 55) Ogata, S., Muramatsu, T. and Kobata, A.: Fractionation of glycopeptides by affinity column chromatography on concanavalin A-Sepharose. J. Biochem., 78, 687–696 (1975).
- 56) Nishigaki, M., Muramatsu, T., Kobata, A. et al.: The broad aglycon specificity of  $\alpha$ -L-fucosidase from marine gastropods. J. Biochem., 75, 509–517 (1974).
- 57) Arakawa, M., Ogata, S., Muramatsu, T. et al.:  $\beta$ -Galactosidase from jack bean and almond emulsin: application for the enzymatic distinction of Gal $\beta$ 1-4GlcNAc and Gal $\beta$ 1-3GlcNAc linkages. J. Biochem., 75, 707–714 (1974).
- 58) Ogata-Arakawa, M., Muramatsu, T. and Kobata, A.:  $\alpha$ -L-Fucosidases from almond emulsin: characterization of the two enzymes with different specificities. Arch. Biochem. Biophys., 181, 353–359 (1977).
- 59) Muramatsu, T., Atkinson, P.H., Nathanson, S.G. et al.: Cell surface glycopeptides: growth-dependent changes in the carbohydratepeptide linkage region. J. Mol. Biol., 80, 781–799 (1973).
- 60) Ceccarini, C., Muramatsu, T., Tsang, J. et al.: Growth-dependent alterations in oligomannosyl cores of glycopeptides. Proc. Natl. Acad. Sci., 72, 3139–3143 (1975).
- 61) Muramatsu, T., Koide, N., Ceccarini, C. et al.: Characterization of mannose-labeled glycopeptides from human diploid cells and their growth-dependent alterations. J. Biol. Chem., 251, 4673–4679 (1976).
- 62) Muramatsu, T., Koide, N. and Ogata, M.: Analysis of oligomannosyl cores of cellular glycopeptides by digestion with endo- $\beta$ -*N*-acetylglucosaminidases. Biochem. Biophys. Res. Commun., 66, 881–888 (1975).
- 63) Muramatsu, T., Ogata, M. and Koide, N.: Characterization of fucosyl glycopeptides from cell surface and cellular material of rat fibroblasts. Biochim. Biophys. Acta, 444, 53–68 (1976).
- 64) Koide, N., Muramatsu, T. and Kobata, A.: Studies on mannosyl glycopeptides from normal and transformed cells. J. Biochem., 85, 149–155 (1979).
- 65) Ogata, S., Muramatsu, T. and Kobata, A.: New structural characteristic of the large glycopeptides from transformed cells. Nature, 259, 580–582 (1976).
- 66) Muramatsu, T., Gachelin, G., Nicolas, J.F. et al.: Carbohydrate structure and cell differentiation: unique properties of fucosyl glycopeptides isolated from embryonal carcinoma cells. Proc. Natl. Acad. Sci., 75, 2315–2319 (1978).
- 67) Muramatsu, T., Gachelin, G. and Jacob, F.: Characterization of glycopeptides from membranes of F9 embryonal carcinoma cells. Biochim. Biophys. Acta, 587, 392–406 (1979).

## CELL SURFACE GLYCOPROTEINS

- 68) Muramatsu, T., Condamine, H., Gachelin, G. et al.: Changes in fucosyl glycopeptides during early postimplantation embryogenesis in the mouse. *J. Embryol. Exp. Morphol.*, 57, 25–36 (1980).
- 69) Muramatsu, T., Gachelin, G., Damonville, M. et al.: Cell surface carbohydrates of embryonal carcinoma cells: polysaccharidic side chains of F9 antigens and receptors to two lectins. FBP and PNA, *Cell*, 18, 183–191 (1979).
- 70) Muramatsu, T., Avner, P., Fellous, M. et al.: Distinctive properties of fucosyl glycopeptides on human teratoma cells. *Somatic Cell Genet.*, 5, 753–761 (1979).
- 71) Prujansky-Jacobovits, A., Gachelin, G., Muramatsu, T. et al.: Surface galactosyl glycopeptides of embryonal carcinoma cells. *Biochem. Biophys. Res. Commun.*, 89, 448–455 (1979).
- 72) Muramatsu, T., Gachelin, G. and Jacob, F.: High-mannose glycopeptides from embryonal carcinoma cells. *J. Biochem.*, 88, 685–688 (1980).
- 73) Muramatsu, H. and Muramatsu, T.: Decreased synthesis of large fucosyl glycopeptides during differentiation of embryonal carcinoma cells induced by retinoic acid and dibutyryl cyclic AMP. *Dev. Biol.*, 90, 441–444 (1982).
- 74) Muramatsu, H., Muramatsu, T. and Avner, P.: Biochemical properties of the high-molecular-weight glycopeptides released from the cell surface of human teratocarcinoma cells. *Cancer Res.*, 42, 1749–1752 (1982).
- 75) Muramatsu, H., Ishihara, H., Miyauchi, T. et al.: Glycoprotein-bound large carbohydrates of early embryonic cells: structural characteristic of the glycan isolated from F9 embryonal carcinoma cells. *J. Biochem.*, 94, 799–810 (1983).
- 76) Muramatsu, H. and Muramatsu, T.: Effects of succinyl concanavalin A and tunicamycin on F9 embryonal carcinoma cells: inhibition of cell spreading. *Dev. Growth Differ.*, 26, 303–310 (1984).
- 77) Muramatsu, H., Miura, I., Gachelin, G. et al.: <sup>1</sup>H-NMR spectroscopy of the glycoprotein-bound large carbohydrates from embryonal carcinoma cells. *Biochem. Int.*, 9, 569–575 (1984).
- 78) Arita, Y., Ogata, S., Ozawa, M. et al.: Distinct properties of receptors for Dolichos biflorus agglutinin (DBA) isolated from small intestine of adult mice and endoderm cells of early embryos and teratocarcinomas. *Differentiation*, 28, 254–259 (1985).
- 79) Noguchi, S., Ogata, S. and Muramatsu, T.: Altered synthesis of glucosamine-labeled glycopeptides during postimplantation embryogenesis of the mouse. *Cell Differ.*, 16, 223–228 (1985).
- 80) Kamada, Y., Arita, Y., Ogata, S. et al.: Receptors for fucose-binding proteins of Lotus tetragonolobus isolated from mouse embryonal carcinoma cells: structural characteristics of the poly(*N*-acetyllactosamine)-type glycan. *Eur. J. Biochem.*, 163, 497–502 (1987).
- 81) Muramatsu, H. and Muramatsu, T.: Analysis of glycoprotein-bound carbohydrates from pluripotent embryonal carcinoma cells by pokeweed agglutinin-agarose. *J. Biochem.*, 107, 629–634 (1990).
- 82) Ozawa, M., Muramatsu, T. and Solter, D.: SSEA-1, a stage-specific embryonic antigen of the mouse, is carried by the glycoprotein-bound large carbohydrate in embryonal carcinoma cells. *Cell Differ.*, 16, 169–173 (1985).
- 83) Kasai, M., Ochiai, Y., Habu, S. et al.: A new differentiation marker selectively expressed on mouse fetal thymocytes. *Immunol. Lett.*, 2, 157–158 (1980).
- 84) Watanabe, M., Muramatsu, T., Shirane, H. et al.: Discrete distribution of binding sites for Dolichos biflorus agglutinin (DBA) and for peanut agglutinin (PNA) in mouse organ tissues. *J. Histochem. Cytochem.*, 29, 779–790 (1981).
- 85) Miyauchi, T., Yonezawa, S., Takamura, T. et al.: A new fucosyl antigen expressed on colon adenocarcinoma and embryonal carcinoma cells. *Nature*, 299, 168–169 (1982).
- 86) Noguchi, M., Noguchi, T., Watanabe, M. et al.: Localization of receptors for Dolichos biflorus agglutinin in early postimplantation embryos in mice. *J. Embryol. Exp. Morph.*, 72, 39–52 (1982).
- 87) Murata, F., Tsuyama, S., Suzuki, S. et al.: Distribution of glycoconjugates in the kidney studied by use of labeled lectins. *J. Histochem. Cytochem.*, 31, 139–144 (1983).
- 88) Hamada, H., Sato, M., Murata, F. et al.: Differential expression of lectin receptors in germ layers of the mouse egg cylinder and teratocarcinomas. *Exp. Cell Res.*, 144, 489–495 (1983).
- 89) Sato, M., Ogata, S., Ueda, R. et al.: Reactivity of a monoclonal antibody raised against human leukemic cells to embryonic and adult tissues of the mouse and teratocarcinomas. *Dev. Growth Differ.*, 25, 333–344 (1983).
- 90) Sato, M. and Muramatsu, T.: Reactivity of five *N*-acetylgalactosamine-recognizing lectins with preimplantation embryos, early postimplantation embryos, and teratocarcinoma cells of the mouse. *Differentiation*, 29, 29–38 (1985).
- 91) Muramatsu, H., Hamada, H., Noguchi, S. et al.: Cell-surface changes during in vitro differentiation of pluripotent embryonal carcinoma cells. *Dev. Biol.*, 110, 284–296 (1985).

- 92) Fujimoto, H., Muramatsu, T., Urushihara, H. et al.: Receptors to *Dolichos biflorus* agglutinin: a new cell surface marker common to teratocarcinoma cells and preimplantation mouse embryos. *Differentiation*, 22, 59–61 (1982).
- 93) Sato, M., Yonezawa, S., Uehara, H. et al.: Differential distribution of receptors for two fucose-recognizing lectins in embryos and adult tissues of the mouse. *Differentiation*, 30, 211–219 (1986).
- 94) Sato, M. and Muramatsu, T.: Oncodevelopmental carbohydrate antigens: distribution of ECMA 2 and 3 antigens in embryonic and adult tissues of the mouse and in teratocarcinomas. *J. Reprod. Immunol.*, 9, 123–135 (1986).
- 95) Muramatsu, T., Gachelin G., Moscona, A.A. et al.: *Teratocarcinoma and Embryonic Cell Interactions*, Japan Scientific Societies Press / Academic Press. Tokyo and New York (1982).
- 96) Muramatsu, T.: Developmentally regulated expression of cell surface carbohydrates during mouse embryogenesis. *J. Cell. Biochem.*, 36, 1–14 (1988).
- 97) Muramatsu, T.: Alterations of cell-surface carbohydrates during differentiation and development. *Biochimie*, 70, 1587–1596 (1988).
- 98) Muramatsu, T.: *Early embryogenesis, Cell Surface Carbohydrates and Cell Development*. Fukuda, M. Ed., CRC Press, Boca Raton, Ann Arbor, London, pp239–256 (1991).
- 99) Kamada, Y., Muramatsu, H., Arita, Y. et al.: Structural studies on a binding site for *Dolichos biflorus* agglutinin in the small intestine of the mouse. *J. Biochem.*, 109, 178–183 (1991).
- 100) Muramatsu, T., Muramatsu, H., Kasai, M. et al.: Receptors for *Dolichos biflorus* agglutinin: new cell surface markers on a spontaneous leukemia. *Biochem. Biophys. Res. Commun.*, 96, 1547–1553 (1980).
- 101) Muramatsu, T., Muramatsu, H. and Ozawa, M.: Receptors for *Dolichos biflorus* agglutinin on embryonal carcinoma cells. *J. Biochem.*, 89, 473–481 (1981).
- 102) Muramatsu, H. and Muramatsu, T.: A fucosyltransferase in teratocarcinoma stem cells: decreased activity after differentiation to parietal endoderm cells. *FEBS Lett.*, 163, 181–184 (1983).
- 103) Muramatsu, H., Kamada, Y. and Muramatsu, T.: Purification and properties of *N*-acetylglucosaminide  $\alpha$  1-3 fucosyltransferase from embryonal carcinoma cells. *Eur. J. Biochem.*, 157, 71–75 (1986).
- 104) Ozawa, M., Muramatsu, T. and Kemler, R.: Molecular properties of ECMA 2 and ECMA 3 antigens defined by monoclonal antibodies against embryonal carcinoma cells. *J. Biochem.*, 97, 307–315 (1985).
- 105) Ozawa, M. and Muramatsu, T.: The glycoprotein-bound large carbohydrates from embryonal carcinoma cells carry receptors for several lectins recognizing *N*-acetylgalactosamine and galactose. *J. Biochem.*, 97, 317–324 (1985).
- 106) Muramatsu, T.: Cell surface glycoproteins as markers in monitoring in vitro differentiation of embryonal carcinoma cells. *Cell Differ.*, 15, 101–108 (1984).
- 107) Muramatsu, T., Muramatsu, H., Ozawa, M. et al.: High-molecular-weight carbohydrates in cell-surface glycoproteins from teratocarcinomas and early embryos, *Teratocarcinoma Stem Cells*. Silver, L., Martin, G. and Strickland, S. Eds., Cold Spring Harbor Press, Cold Spring Harbor, pp173–184 (1983).
- 108) Kawata, M., Higaki, K., Sekiya, S. et al.: Antibodies to large glycopeptides in sera from patients with ovarian germ cell tumors. *Clin. Exp. Immunol.*, 51, 401–406 (1983).
- 109) Ozawa, M., Higaki, K., Kawata, M. et al.: An  $\alpha$ -galactosyl residue in the large carbohydrates of teratocarcinoma cells: the antigenic determinant recognized by sera from patients with ovarian germ cell tumors. *Biochem. Biophys. Res. Commun.*, 115, 268–274 (1983).
- 110) Kawata, M., Sekiya, S., Takamizawa, H. et al.: Molecular properties of F9 embryoglycan recognized by a unique antibody in sera from patients with germ cell tumors. *Cancer Res.*, 47, 2288–2294 (1987).
- 111) Fushuku, N., Muramatsu, H., Uezono, M.M. et al.: A new endo- $\beta$ -galactosidase releasing Gal $\alpha$ 1-3Gal from carbohydrate moieties of glycoproteins and from a glycolipid. *J. Biol. Chem.*, 262, 10086–10092 (1987).
- 112) Muramatsu, T.: An endo- $\beta$ -galactosidase releasing Gal $\alpha$ 1-3Gal. *Methods Enzymol.*, 179, 496–504 (1989).
- 113) Kamada, Y., Muramatsu, H., Muramatsu, T. et al.: Glycopeptides from murine teratocarcinoma cells: structure of the determinants recognised by *Griffonia simplicifolia* agglutinin I and by sera from patients with ovarian germ cell tumors. *Carbohydr. Res.*, 176, 237–243 (1988).
- 114) Kamada, Y., Muramatsu, H., Kawata, M. et al.: Poly-*N*-acetylglucosamines synthesized by cultured Ehrlich carcinoma cells: application of endo- $\beta$ -galactosidase C for analysis of the terminal structure. *J. Biochem.*, 104, 738–741 (1988).
- 115) Nomoto, S., Muramatsu, H., Ozawa, M. et al.: An anti-carbohydrate monoclonal antibody inhibits cell substratum adhesion of F9 embryonal carcinoma cells. *Exp. Cell Res.*, 164, 49–62 (1986).
- 116) Yoshinaga, K., Muramatsu, H. and Muramatsu, T.: Immunohistochemical localization of the carbohydrate antigen 4C9 in the mouse embryo: a reliable marker of mouse primordial germ cells. *Differentiation*, 48, 75–82 (1991).

## CELL SURFACE GLYCOPROTEINS

- 117) Ozawa, M., Sato, M. and Muramatsu, T.: Basement membrane glycoprotein laminin is an agglutinin. *J. Biochem.*, 94, 479–485 (1983).
- 118) Suganuma, T., Muramatsu, H., Muramatsu, T. et al.: Subcellular localization of *N*-acetylglucosaminide  $\beta$ 1-4 galactosyltransferase revealed by immunoelectron microscopy. *J. Histochem. Cytochem.*, 39, 299–309 (1991).
- 119) Ikematsu, S., Kaname, T., Ozawa, M. et al.: Transgenic mouse lines with ectopic expression of  $\alpha$ -1,3-galactosyltransferase: production and characteristics. *Glycobiology*, 3, 575–580 (1993).
- 120) Ozawa, M., Yonezawa, S., Sato, E. et al.: A new glycoprotein antigen common to teratocarcinoma, visceral endoderm and renal tubular brush border. *Dev. Biol.*, 91, 351–359 (1982).
- 121) Ozawa, M., Sato, M., Muramatsu, H. et al.: A membrane glycoprotein involved in teratocarcinoma cell adhesion to substratum. *Exp. Cell Res.*, 158, 127–143 (1985).
- 122) Ozawa, M., Yonezawa, S., Sato, M. et al.: Three groups of teratocarcinoma antigens co-expressed in the visceral endoderm are located in mutually exclusive sites in the adult kidney. *Dev. Growth Differ.*, 29, 455–468 (1987).
- 123) Ozawa, M., Huang, R.-P., Furukawa, T. et al.: A teratocarcinoma glycoprotein carrying a developmentally regulated carbohydrate marker is a member of the immunoglobulin gene superfamily. *J. Biol. Chem.*, 263, 3059–3062 (1988).
- 124) Huang, R.-P., Ozawa, M., Kadomatsu, K. et al.: Developmentally regulated expression of embigin, a member of the immunoglobulin superfamily found in embryonal carcinoma cells. *Differentiation*, 45, 76–83 (1990).
- 125) Huang, R.-P., Ozawa, M., Kadomatsu, K. et al.: Embigin, a member of the immunoglobulin superfamily expressed in embryonic cells, enhances cell-substratum adhesion. *Dev. Biol.*, 155, 307–314 (1993).
- 126) Ogata, S., Kamachi, Y., Arita, Y. et al.: A re-examination of the isolectin components of the fucose-binding proteins of *Lotus tetragonolobus*. *Carbohydr. Res.*, 144, 297–304 (1985).
- 127) Miyauchi, T., Kanekura, T., Yamaoka, A. et al.: Basigin, a new, broadly distributed member of the immunoglobulin superfamily has strong homology with both the immunoglobulin V domain and the  $\beta$ -chain of major histocompatibility complex class II antigen. *J. Biochem.*, 107, 316–323 (1990).
- 128) Miyauchi, T., Masuzawa, Y. and Muramatsu, T.: The basigin group of the immunoglobulin superfamily: complete conservation of a segment in and around transmembrane domains of human and mouse basigin and chicken HT7 antigen. *J. Biochem.*, 110, 770–774 (1991).
- 129) Furukawa, T., Ozawa, M., Huang, R.-P. et al.: A heparin binding protein whose expression increases during differentiation of embryonal carcinoma cells to parietal endoderm cells: cDNA cloning and sequence analysis. *J. Biochem.*, 108, 297–302 (1990).
- 130) Nakamoto, M., Ozawa, M., Jacinto, S.D. et al.: Heparin binding protein-44 (HBP-44) associates with brushin, a high molecular weight glycoprotein antigen common to the kidney and teratocarcinomas. *J. Biochem.*, 114, 344–349 (1993).
- 131) Sato, M., Ozawa, M., Hamada, H. et al.: Cell surface markers to monitor the process of visceral endoderm differentiation from embryonal carcinoma cells: identification of the stage sensitive to high concentration of retinoic acid. *J. Embryol. Exp. Morph.*, 88, 165–182 (1985).
- 132) Ozawa, M. and Muramatsu, T.: Reticulocalbin, a novel endoplasmic reticulum resident  $\text{Ca}^{2+}$ -binding protein HDEL sequence. *J. Biol. Chem.*, 268, 699–705 (1993).
- 133) Miyauchi, T., Muramatsu, H., Ozawa, M. et al.: Receptors for peanut agglutinin isolated from cell lines of human Burkitt lymphoma, lung squamous cell carcinoma and gastric signet ring cell carcinoma. *Gann*, 73, 581–587 (1982).
- 134) Shimoda, N., Muramatsu, H., Osawa, T. et al.: Carbohydrate moieties of peanut agglutinin receptors isolated from human gastric cancer cells. *J. Biochem.*, 102, 657–664 (1987).
- 135) Masuzawa, Y., Miyauchi, T., Hamanoue, M. et al.: A novel core protein as well as polymorphic epithelial mucin carry peanut agglutinin binding sites in human gastric carcinoma cells: sequence analysis and examination of gene expression. *J. Biochem.*, 112, 609–615 (1992).
- 136) Fukuda, K., Matsuyama, H., Fukami, K. et al.: Analysis of glycoproteins from nasal polyps and papillomas by affinity chromatography. *Ann. Otol. Rhinol. Laryngol.*, 93, 85–88 (1984).
- 137) Uehara, F., Muramatsu, T., Sameshima, M. et al.: Identification of peanut agglutinin receptors in the monkey retina. *Exp. Eye Res.*, 37, 303–305 (1983).
- 138) Matsushita, Y., Yonezawa, S., Nakamura, T. et al.: Carcinomaspecific *Ulex europaeus* agglutinin-I binding glycoproteins of human colorectal carcinoma and its relation to carcinoembryonic antigen. *J. Natl. Cancer Inst.*, 75, 219–226 (1985).
- 139) Uehara, F., Muramatsu, T. and Ohba, N.: Two-dimensional gel electrophoretic analysis of lectin receptors from the bovine interphotoreceptor matrix. *Exp. Eye Res.*, 43, 227–234 (1986).

- 140) Uehara, F., Sameshima, M., Muramatsu, T. et al.: Affinity isolation of cone outer segments using a peanut agglutinin-nitrocellulose sheet. *Exp. Eye Res.*, 43, 687–693 (1986).
- 141) Unoki, K., Uehara, F. and Muramatsu, T.: Distribution of glycosyltransferase in bovine eyes. *Ophthalmic Res.*, 22, 342–359 (1990).
- 142) Uehara, F., Ohba, N., Nakashima, Y. et al.: Developmental change of distribution of  $\beta$ -galactoside  $\alpha$ 2,6-sialyltransferase mRNA in rat retina. *Exp. Eye Res.*, 56, 89–93 (1993).
- 143) Uehara, F., Ohba, N., Nakashima, Y. et al.: A fixative suitable for in situ hybridization histochemistry. *J. Histochem. Cytochem.*, 41, 947–953 (1993).
- 144) Uehara, F., Ohba, N., Nakashima, Y. et al.: Distribution of peripherin/rds mRNA in cone-dominant squirrel retina. *Exp. Eye Res.*, 56, 611–613 (1993).
- 145) Matsusako, T., Muramatsu, H., Shirahama, T. et al.: Expression of a carbohydrate signal, sialyl dimeric Le<sup>x</sup> antigen is associated with metastatic potential of transitional cell carcinoma of the human urinary bladder. *Biochem. Biophys. Res. Commun.*, 181, 1218–1222 (1991).
- 146) Matsumoto, H., Muramatsu, H., Muramatsu, T. et al.: Carbohydrate profiles revealed by a lectin and a monoclonal antibody correlate with metastatic potential and prognosis of human lung carcinomas. *Cancer*, 69, 2084–2090 (1992).
- 147) Shirahama, T., Ikoma, M., Muramatsu, H. et al.: Reactivity to fucose-binding proteins of *Lotus tetragonolobus* correlates with metastatic phenotype of transitional cell carcinoma of the bladder. *J. Urol.*, 147, 1659–1664 (1992).
- 148) Shirahama, T., Ikoma, M., Muramatsu, T. et al.: Expression of SSEA-1 carbohydrate antigen correlates with stage, grade and metastatic potential of transitional cell carcinoma of the bladder. *J. Urol.*, 148, 1319–1322 (1992).
- 149) Matsusako, T., Muramatsu, H., Shirahama, T. et al.: A metastasis-associated antigen is present on a 60 KDa glycoprotein in transitional cell carcinoma of the human urinary bladder. *Histochem. J.*, 24, 805–810 (1992).
- 150) Matsumoto, H., Muramatsu, H., Shimotakahara, T. et al.: Correlation of expression of ABH blood group carbohydrate antigens with metastatic potential in human lung carcinoma. *Cancer*, 72, 75–81 (1993).
- 151) Shirahama, T., Ikoma, M., Muramatsu, T. et al.: The binding site for fucose-binding proteins of *Lotus tetragonolobus* is a prognostic marker for transitional cell carcinoma of the human urinary bladder. *Cancer*, 72, 1329–1334 (1993).
- 152) Muramatsu, T.: Carbohydrate signals in metastasis and prognosis of human carcinomas. *Glycobiology*, 3, 291–296 (1993).
- 153) Muramatsu, T.: Les sucres de la membrane cellulaire. *La Recherche*, 20, 624–634 (1989).